OLA SOLLENIUS

EARLY CORRECTION OF UNILATERAL POSTERIOR CROSSBITE

Evidence-based evaluations of oral health related quality
of life, cost-effectiveness and 3D treatment effects

BÄSTA LIVSPLATSEN

Region Halland

MALMÖ
UNIVERSITY

EARLY CORRECTION OF UNILATERAL
POSTERIOR CROSSBITE

Malmö University, Faculty of Odontology
Doctoral Dissertations 2019

© Copyright Ola Sollenius, 2019

Illustrations: Ola Sollenius

ISBN 978-91-7877-018-2 (print)

ISSN 978-91-7877-019-9 (pdf)

Tryck: Holmbergs, Malmö 2019

OLA SOLLENIUS

**EARLY CORRECTION OF
UNILATERAL POSTERIOR
CROSSBITE**

Evidence-based evaluations of oral health related
quality of life, cost-effectiveness and 3D treatment
effects

Department of Orthodontics
Faculty of Odontology, Malmö University, Sweden
Department of Orthodontics, Region Halland
Halmstad, Sweden 2019

This publication is also available at:
<https://muep.mau.se>

To Jenny, Casper and Selma

CONTENTS

PREFACE.....	9
THESIS AT A GLANCE.....	10
ABSTRACT	11
POPULÄRVETENSKAPLIG SAMMANFATTNING	15
Klinisk betydelse.....	16
INTRODUCTION	17
Aetiology	17
Prevalence.....	17
Treatment	17
Why should we treat?	18
When should we treat?	19
What kind of appliance should we use?	19
Three-dimensional scanning	24
Significance.....	25
AIMS	26
HYPOTHESIS	27
MATERIALS AND METHODS.....	28
ITT	31
Cost-minimisation analysis	31
Complementary and new studies.....	35
CPQ/Oral health related quality of life	35
Three-dimensional changes in two groups	35
Statistical analysis Paper II-IV	35
Sample size calculation	36

RESULTS	37
Paper I - Systematic review	37
Type of study	38
Evaluation of study quality and evidence	38
New literature search.....	43
Study design.....	44
Evaluation of study quality and evidence	44
Paper II Quality of life	47
Paper III Cost-minimisation	50
Paper IV Treatment effects.....	52
DISCUSSION.....	58
Methodological considerations.....	59
Intention-to-treat (ITT)	62
The Hawthorne effect	63
Quality of life.....	63
Cost-minimisation analysis	65
How much money could be saved?.....	65
Consequences for the general dentist	66
Three-dimensional analysis	67
Environmental considerations	67
New data.....	68
Ethical considerations	68
Generalisability.....	69
Overall limitations of this thesis	69
Future research	70
CONCLUSIONS.....	71
Clinical implications.....	73
ACKNOWLEDGEMENTS	74
REFERENCES	76
APPENDIX A+B CPQ 8-10 QUESTIONNAIRE	81
PAPERS I-IV.....	93

PREFACE

This thesis is based on the following papers, which are referred to in the text by their Roman numerals I-IV.

- I. Sollenius, O., Petrén, S., Paulsson, L., Norlund, A. and Bondemark L. (2016) Health economic evaluations in orthodontics; a systematic review. *European Journal of Orthodontics*. 38, 259-265.
- II. Kallunki, J., Sollenius, O., Paulsson, L., Petrén, S., Dimberg, L. and Bondemark L. (2019) Oral health related quality of life among children with excessive overjet or unilateral posterior crossbite with functional shift compared to children with no or mild orthodontic treatment need. *European Journal of Orthodontics*. 41, 111-116
- III. Sollenius, O., Petrén, S. and Bondemark L. (2019) An RCT on clinical effectiveness and cost-analysis of correction of unilateral posterior crossbite with functional shift in specialist and general dentistry. *European Journal of Orthodontics*. 11. pii: cjz014. doi: 10.1093/ejo/cjz014
- IV. Sollenius, O., Golež, A., Primožič, J., Ovsenik, M., Bondemark, L. and Petrén S. (2019) Three-dimensional evaluation of forced unilateral posterior crossbite correction in the mixed dentition – a randomised controlled trial. *European Journal of Orthodontics*. 1. pii: cjz054. doi: 10.1093/ejo/cjz054

These papers are reprinted with kind permission from the copyright holders.

THESIS AT A GLANCE

Study	Purpose	Study design	Sample	Main outcome parameter	Main findings
I	Assess the evidence from studies presenting orthodontic treatment outcomes and the related costs.	Systematic review	1,838 articles from seven databases for the period from 1966 to September 2014.	RCTs and CCTs comparing at least two different orthodontic interventions, evaluation of both economic and orthodontic outcomes.	There is currently insufficient evidence available about the health economics of orthodontic interventions. Further studies are warranted.
II	To compare oral health related quality of life (OHRQoL) among 9-year-old children with excessive overjet (EO) to children with unilateral posterior crossbite with functional shift (UPC) and children with normal occlusion (NO).	Clinical controlled trial	The sample sourced from 19 Public Dental Service Clinics in Sweden. The UPC group comprised 93 children, the EO group 71 children, and the NO group 65 children.	Child Perceptions Questionnaire (CPQ8-10) for the evaluation of OHRQoL, overjet and overbite, caries, trauma, enamel defects, and headache.	Children with EO reported significantly lower OHRQoL compared to children with UPC or normal occlusion. The children generally reported low CPQ scores that imply an overall fairly good OHRQoL.
III	To evaluate the costs of quad-helix (QH) and removable expansion plate (EP) treatments performed either in specialist or general dentistry for the correction of unilateral posterior crossbite with functional shift in mixed dentition.	Randomised controlled trial	110 patients randomised into four groups: QH or EP treatment at either specialist orthodontic clinics or general dentistry clinics.	A cost-analysis was performed with reference to intention-to-treat (ITT), regarding direct costs, indirect costs and societal costs.	Treatment of unilateral posterior crossbite in mixed dentition is recommended to be performed by a specialist orthodontist using the QH since the QH treatment performed in specialist orthodontic clinics had the highest costeffectiveness.
IV	To assess the 3D treatment changes of unilateral posterior crossbite correction using either QH or EP appliances in mixed dentition, and to compare the treatments to the 3D changes occurring in age-matched untreated unilateral posterior crossbite patients as well as in subjects with normal occlusion (NO).	Randomised controlled trial	135 patients randomised into five groups, QH or EP treatment at specialist orthodontic clinics or general dentistry clinics, untreated controls and another 25 patients with normal occlusion (NO) who served as controls.	3D palatal surface area, palatal projection area, palatal shell volume; 2D measurements of intermolar and intercanine linear distances.	After treatment, no significant differences were found between the four treatment groups and the normal group, which implies that the surface and projection area together with the shell volume for the four treatment groups and the normal group were equivalent. The natural growth changes were small.

ABSTRACT

Unilateral posterior crossbite (UPC) with functional shift is one of the most common malocclusions in mixed dentition. Left untreated, it may cause long-term effects on the growth and development of the jaws and teeth. Early orthodontic intervention is usually undertaken to correct the condition at the primary or mixed dentition stage. Quad-helix (QH) or removable expansion plates (EP) are common treatment alternatives to correct crossbites and treatment of crossbite may consume a relatively large part of the total resources in orthodontic care. In Sweden the treatment can be performed either in specialist orthodontic clinics or in general dentistry. The role of the orthodontist in general dental care is essential in diagnostics and treatment planning, whereas general dentists may often provide a part of the treatment after consultation with, or under the supervision of, an orthodontist.

Economic evaluations have become an integral component of health services. The main reason is that resources within the health sector (personnel, time, facilities, equipment, and knowledge) are limited. Nevertheless, studies comparing the costs of orthodontic treatments performed in general versus specialist dentistry are virtually non-existent.

To date there are no studies evaluating the oral health related quality of life (OHRQoL) in children with unilateral posterior crossbite. It is thus important to analyse and compare the OHRQoL between children with and without malocclusions (children with normal occlusion).

When treatment effects of unilateral crossbite correction have been assessed, most studies have used two-dimensional evaluations such as linear measurements. However, orthodontic treatment effects including crossbite correction may, if possible, be described in all three planes. There are no studies in the literature that have explicitly investigated the three-dimensional treatment effects of unilateral crossbite correction between different appliances and, in this context, related the treatment changes to growth changes associated with untreated subjects with unilateral posterior crossbite as well as in subjects with normal occlusion and with no or mild orthodontic treatment need.

The research questions addressed in this thesis originate from identified knowledge gaps and clinical needs in orthodontic care, and to provide as high clinical evidence as possible a multi-centre randomised control trial (RCT) has been performed as well as a controlled trial regarding comparisons and the impact different malocclusions may have on OHRQoL.

The results are expected to be beneficial for the patients who will be offered the most widely accepted and effective treatment, which is of importance for the dentists for decisions as to which treatment will give the best outcome, and beneficial for both dental care providers and society in care planning and the allocation of resources.

Therefore, this thesis was based on four studies:

Paper I: a systematic literature review was undertaken to answer the following questions:

- Are there any articles regarding health economics in orthodontics and is it possible to make any conclusions from the articles?

The literature search spanned from January 1966 to September 2014 and was later supplemented and extended to April 2019.

Paper II: the aims were to investigate:

- The OHRQoL using the Child Perceptions Questionnaire (CPQ8-10) in 93 children with unilateral posterior crossbite, 71 children with excessive overjet and 65 children with normal occlusion with no or mild orthodontic treatment need.

Paper III and IV: these two papers originated from a multi-centre RCT. The aims were to investigate:

- Clinical effectiveness and cost-analysis in specialist and general dentistry (Paper III)
- Three-dimensional evaluations of crossbite correction (Paper IV)

Key findings in Paper I and the supplementary search:

- Few orthodontic studies have presented both economic and clinical outcomes. There is currently insufficient evidence available about the health economics of orthodontic interventions.
- Further studies are still warranted and preferably using the same clinical outcomes.

Key findings in Paper II

- Children with excessive overjet reported significantly lower OHRQoL compared to children with unilateral posterior crossbite or normal occlusion.
- The children generally reported low CPQ scores that imply an overall fairly good OHRQoL.

Key findings in Paper III

- Treatment of unilateral posterior crossbite in mixed dentition is recommended to be performed by a specialist orthodontist using the quad-helix appliance since the quad-helix treatment performed in specialist orthodontic clinics had the highest cost-effectiveness.

Key findings in Paper IV

- Crossbite children had, before treatment, significantly smaller palatal surface and volume than normal control children.
- After treatment, there were no significant differences between the treatment groups and the normal group, which implies that the palatal surface and projection area together with the palatal shell volume for the treatment groups and the normal group were equivalent.

Conclusions and clinical implications:

With a superior success rate and cost-effectiveness, it is concluded that treatment of unilateral posterior crossbite in mixed dentition is recommended to be performed by specialist orthodontists using the quad-helix appliance.

The unilateral posterior crossbite correction resulted in a normalisation of the occlusion, palatal area and volume.

POPULÄRVETENSKAPLIG SAMMANFATTNING

Tvångsfört enkelsidigt korsbett är en vanlig bettavvikelse i växelbettet, dvs. i de åldrar då barnets mjölkttänder tappas och ersätts med permanenta tänder. Korsbettet uppstår då överkäken är för smal i förhållande till underkäken och för att bettet skall passa ihop skjuter barnet underkäken åt höger eller vänster. Korsbett behandlas vanligtvis med en fastsittande tandställning, quad-helix, eller en avtagbar expansionsplåt innanför tänderna och i gommen.

I Sverige ser det olika ut om behandlingarna utförs med quad-helix eller expansionsplåt. Det skiljer också om behandlingarna utförs av specialisttandläkare på tandregleringskliniker (specialisttandvård) eller som mer brukligt av allmäntandvårdstandläkare inom allmäntandvården.

I avhandlingens första delarbete genomfördes en systematisk litteraturöversikt från januari 1966 till september 2014 och förnyades sedan till och med april 2019. Målet med översikten var att definiera vilket vetenskapligt underlag som finns för hälsoekonomiskt utfall inom tandreglering.

Det andra delarbetet består av data från en kontrollerad studie där den munhälsorelaterade livskvaliteten undersöktes och jämfördes mellan barn med enkelsidigt korsbett, barn med stort överbett samt barn med normalbett.

För att uppnå ett så högt vetenskapligt underlag som möjligt, utfördes i delarbete tre och fyra randomiserade kontrollerade studier (RCT), vilket innebär att deltagande patienter har lottats till quad-helix eller expansionsplåt respektive behandling i allmän- eller specialisttandvård. För att ytterligare öka det vetenskapliga underlaget användes flera kliniker vilket benämns multicenterstudier. I delarbete tre analyseras och jämförs kostnadseffektiviteten mellan quad-helix och expansionsbehandling samt om kostnadseffekten skiljer sig mellan behandling utförd i allmän- respektive specialisttandvård. I delarbete fyra utvärderas och jämförs de tre- och tvådimensionella behandlingseffekterna i över- och underkäken efter behandling med quad-helix och expansionsplåt samt om behandling sker i allmän- eller i specialisttandvård.

Avhandlingens slutsatser:

- det finns mycket få vetenskapliga studier som utförts avseende kliniskt och ekonomiskt resultat efter tandregleringsbehandling, d.v.s. kunskapsluckorna inom detta område är stora och därför behövs nya hälsoekonomiska studier inom tandreglering.
- barn med stora överbett har sämre munhälsorelaterad livskvalitet jämfört med barn med korsbett och barn med normalbett.
- korsbettsbehandlingar utförda med quad-helix inom specialisttandvården är mer kostnadseffektiva jämfört med behandlingar inom allmäntandvården.
- efter lyckad behandling och oavsett tandställningsmetod eller behandlingsinstans så uppnås likvärdig tre- och tvådimensionell form av överkäken som hos barn med normalbett.

Klinisk betydelse

Behandling av tvångsförda enkelsidiga korsbett i växelbettet rekommenderas att utföras med quad-helix av specialisttandläkare på tandregleringsklinik eftersom detta alternativ är överlägsen expansionsplåten och behandlingar i allmäntandvården med avseende på behandlings- och kostnadseffektivitet. Behandlingen resulterade i en normalisering av bettet och av gommens form.

INTRODUCTION

Aetiology

In primary and mixed dentition, unilateral posterior crossbite with functional shift commonly arises as a result of a narrow maxilla, which forces the mandible to displace or deviate laterally into an abnormal position due to the presence of tooth interferences.^{1,2} The mandibular deviation is not present when the mandible is at rest, but is seen upon closure.

The constricted upper arch can originate from dental problems, skeletal problems or both. The reason for the constricted upper arch/jaw can be caused by a low tongue position. A prolonged use of a pacifier or thumb sucking and also constricted nasal airways results in a low tongue position.

Prevalence

Unilateral posterior crossbite (UPC) with functional shift is a common malocclusion in mixed dentition, with a reported prevalence ranging from 8.5% to 17%.³⁻⁵

Lack of standardisation of definitions, with or without functional shift, subject age, population and sample size can be the cause of the wide range of prevalence.

Treatment

In primary dentition early interceptive grinding seems to be an effective alternative. If the transverse inter-jaw discrepancies were minor, Lindner showed that when grinding was performed, 50% of the children with unilateral posterior crossbite exhibited a normal transverse occlusion evaluated in mixed dentition, at the age of 9.⁶

If the crossbite was left without interceptive intervention only 17% showed a spontaneous correction of the crossbite. In mixed dentition removable expansion plate, quad-helix or rapid maxillary expansion are appliances used to correct unilateral posterior crossbite.

Why should we treat?

If unilateral posterior crossbite with functional shift is left untreated this may lead to asymmetry of both dental arches as well as adverse effects on the temporomandibular joints, pain in the masticatory system, clicking and locking of joints, and altered facial growth including facial asymmetry.⁷⁻⁹ In a recent article using three-dimensional analysis it was investigated where the changes in the mandible occur, and it was concluded that left untreated unilateral posterior crossbites can cause asymmetry of the condylar, angular, and alveolar process plus the mandibular ramus.¹⁰

The relationship between unilateral posterior crossbite and skeletal asymmetry is still unresolved. To date, most of the studies available report a skeletal asymmetric growth. Electromyography activity of masticatory muscles is different between crossbite and non-crossbite sides. Subjects with unilateral posterior crossbite show a smaller bite force than non-crossbite subjects. In a systematic review by Tsanidis et al.¹¹ they found that abnormal masticatory cycle associated with functional posterior unilateral crossbite tends to normalise following early crossbite treatment. Masticatory muscle activity shows an increase after early functional unilateral posterior crossbite treatment, and this activity approaches normal levels after treatment.

There is no consistency in studies reporting masticatory muscle thickness asymmetry in unilateral posterior crossbite subjects.¹² There are also studies that are less conclusive regarding the association between unilateral posterior crossbites and temporomandibular disorder.^{13,14} Thilander and Bjerklin¹³ pointed out that it was uncertain whether early crossbite treatment can prevent future problems in joints and masticatory muscles. Nonetheless, the aims of treatment are to eliminate the occlusal interference, attain good dental arch symmetry, and harmonise the muscular pattern.

When should we treat?

Some sources advocate treatment in primary dentition not only to correct functional asymmetry due to the lateral mandibular shift but also to prevent adverse effects on the maxillary growth, and some recommend treatment in mixed dentition.^{15,16} It is also possible to treat in early permanent dentition using a rapid maxillary expansion (RME) or a surgery-assisted expansion (SARME) if the midpalatal suture is fused later in adult life.

In deciduous dentition, treatment of unilateral posterior crossbite may involve the grinding of interfering teeth or expansion appliances. If the maxillary arch is wide enough, the interfering cusp can be ground off but if the maxillary arch is too narrow it should be expanded by an orthodontic appliance.

In early mixed dentition the grinding of interfering teeth and expansion appliances are also treatment alternatives as well as cross-elastics between the maxillary and mandibular permanent first molars when these are in a single crossbite situation. Expansion of the maxillary arch can be achieved in the deciduous, mixed or early permanent dentition using either removable or fixed appliances.

What kind of appliance should we use?

History

Maxillary expansion treatments have been used for more than a century to correct maxillary transverse deficiency. The earliest commonly cited report is that of E.C. Angell published in *Dental Cosmos* in 1860.¹⁷ He used a removable jack screw to expand the maxilla.

Effects

Three expansion treatment modalities are used today: rapid maxillary expansion (RME), slow maxillary expansion (SME) and surgically assisted maxillary expansion (SARME). Since each treatment modality has its advantages and disadvantages, controversy regarding the use of each exists. Very often practitioners select treatment appliances based on their personal experiences and on the patient's age and malocclusion.

Advocates of rapid maxillary expansion believe that it results in minimum dental movement (tipping) and maximum skeletal movement, it effects not just the maxilla but also other bones in the face and head.¹⁸ Rapid maxillary expansion is indicated in cases with a transverse discrepancy equal to or greater than 4 mm, and where the maxillary molars are already buccally inclined to compensate for the transverse skeletal discrepancy.¹⁹

There is some evidence to suggest that slow maxillary expansion with a quad-helix (QH) appliance may be more successful than removable expansion plates (EP) at correcting posterior crossbites and expanding the inter-molar width for children in mixed dentition.^{16,20}

The removable expansion plate

The removable expansion plate (EP) consists of acrylic with an expansion screw and stainless steel clasps on the primary and permanent first molars (**Figure 1**). The EP can be activated either 0.2 mm once or 0.2 mm twice a week by the patient until a normal transverse relationship is achieved. The dentist thoroughly instructs the patient to use the EP day and night except for meals and tooth brushing. Treatment progression is evaluated every 6 weeks, and the result is retained for approximately 3 months. Adjustment of steel clasps performed when needed. It is also possible to let the acrylic cover the occlusal surfaces of the lateral teeth, to unlock the occlusion.

The quad-helix

The quad-helix (QH) consists of a standard stainless steel arch with stainless steel bands attached with glass ionomer cement on the maxillary first molars (**Figure 1**). The QH can be either removable or soldered to the bands. The four helices increase the flexibility and range of activation. The QH is activated 10 mm before placement and then reactivated every 6 weeks until a normal transverse relationship is achieved. Steel or elastic ligatures are used to secure the QH to the bands. If the occlusion is locked in a severe way, it is possible to unlock the occlusion using glass-ionomer cement on the occlusal surfaces. The treatment result is retained for 3 months.

The rapid maxillary expansion

The rapid maxillary expansion (RME) (**Figure 1**) consists of metal cemented on second primary molars or on first permanent molars, and bonded to the palatal surface of the deciduous maxillary canines. An alternative to the cemented RME is one that is retained by temporary anchorage devices (TADs). The RME will be worn continually for a period of 3-6 months. During the first 2-3 weeks, the RME is activated twice a day until a normal decried expansion of the maxilla is obtained. In the remaining months it is left in the mouth to allow new bone to fill the gap under the gum.

Figure 1 A. Expansion plate. B. Quad-helix. C. RME (Photo with kind permission of Dr Marco Rosa.)

Discomfort

All appliances have their advantages and disadvantages, including pain in the teeth and jaw as well as mechanical wear on mucosa. Using a plate can cause initial speech difficulty due to palatal coverage of the appliance, breakage and loss of appliances also happen due to patients' carelessness. Other disadvantages of removable appliances include limited tooth movement range, appliance bulkiness, and poor oral hygiene maintenance of the appliance. On the other hand, separation ligatures and placement of the bands can also cause discomfort for patients with the quad-helix.

Compliance

Oral hygiene is, of course, essential, but so is good compliance when using a removable appliance such as the expansion plate. Without compliance there would not be any results.

Treatment success rate

There is some evidence to suggest that the quad-helix (fixed) appliance may be more successful than removable expansion plates at correcting posterior crossbites and expanding the first maxillary molars for children in mixed dentition (aged 8-10 years).²⁰

Petrén and Bondemark¹⁶ showed a success rate of 100% using quad-helix appliance and 66% success rate using expansion plates.

Economic evaluation in healthcare

For many years, dental healthcare for children and adolescents up until 19 years of age has been free of charge in Sweden. Recently, this age limit has been raised to 23 years. Approximately one-third of the children and adolescents in Sweden are offered some kind of orthodontic treatment during growth.²¹ Treating patients with a cost-effective alternative is advocated from both a patient and a societal perspective. Economic evaluation has become an integral component of health service research in recent years and is likely to assume increasing importance for decision-making in dental services in the future.²² The main reason is that resources within the health sector (personnel, time, facilities, equipment, and knowledge) are limited.²³ Thus, failure to analyse the economic aspects of dental health services may result in unsustainable over-expenditure or withdrawal or reduction of services or resources in other areas of healthcare.²⁴

Four main types of analysis are available for economic evaluations: cost-effectiveness analysis, cost-minimisation analysis, cost-utility analysis and cost-benefit analysis.²³ Economic evaluations with a societal perspective include calculations of direct and indirect costs. Direct costs comprise material costs and treatment time costs and indirect costs are defined as loss of income (wages plus social security costs) incurred by the patients' parents' absence from work to accompany the patient to the orthodontic appointment, i.e. total time (including waiting time) at the clinic and the travelling duration.

Cost effectiveness between appliances

In a cost-minimisation analysis from Petré et al.²⁵ it was clearly shown that for correction of posterior crossbite, quad-helix (QH) offers significant economic benefits over expansion

plate (EP) treatment. The QH had lower direct and indirect costs and fewer failures needing re-treatment. Even with full co-operation, i.e. when only successful treatments were considered, EP treatment was more expensive than the QH treatment.

Evidence-based health evaluation

The basis for evidence-based health evaluation is the systematic review. The systematic review may identify knowledge gaps, manage volumes of information, and clarify the need for trials. A systematic review is also an excellent tool to provide comprehensive summaries on evidence of a certain scientific field, and, thereby, make the available evidence more accessible to the decision makers as well as for the clinicians since it is almost impossible for them to profit from all the information available. Consequently, from an evidence-based perspective, a systematic review of the literature on cost-effectiveness in orthodontics would increase the understanding of the topic.

A report from the Swedish Agency for Health Technology Assessment of Social Services (SBU) considering orthodontic and health economics was published in 2005²¹ and, in order to be updated a new systematic review has to be carried out on health economic evaluations in orthodontics.²⁶

In a recent systematic review by Dimberg et al.²⁷, they found that malocclusions have negative effects on children's and adolescents' OHRQoL, and that there are no studies that explicitly investigate how a specific malocclusion like unilateral posterior crossbite with functional shift may affect the OHRQoL in young children, and, furthermore, that high-quality studies from Europe are lacking. Therefore, new studies on this topic are needed and specifically to analyse if separate malocclusions have an impact on OHRQoL.

It is well known that scientific assessment in healthcare aims to identify interventions that offer the greatest benefits for the patients while utilising resources in the most effective way. A systematic review have disclosed that further studies providing high level of evidence, i.e. randomised controlled trials (RCTs) are needed to determine which treatment of unilateral posterior crossbite is the most cost-effective and offers the best patient acceptance.²⁸

In Sweden and some other countries the role of orthodontists in general dental care is essential in diagnostics and treatment planning, whereas general dentists may often provide a part of the treatment after consultation with, or under the supervision of, an orthodontist. Examples of treatments performed by general dentists are the correction of posterior or anterior crossbites, or the correction of overjet, often performed in mixed dentition. However, there are very few studies that compare orthodontic treatment outcome in comparison with economic outcome, especially in unilateral posterior crossbite and in general no comparisons between care performed in general and specialist dental care.

Three-dimensional scanning

Most of the previous studies on crossbite correction have concentrated on dental changes. Many of the claims about orthopaedic change are based on the analysis on two-dimensional (2D) radiographs, sometimes involving case series with an absence of untreated controls. In addition, it is important to use new technology and well-structured study design to address the deficiency of knowledge on this topic. In recent years, three-dimensional (3D) scanners are gaining increasing popularity and, moreover, the evaluation can now be performed in 3D. Still evaluations in 3D are scarce particularly regarding crossbite corrections in mixed dentition.

Plaster casts have traditionally been used for orthodontic diagnosis and treatment planning in a 2D technique. Tooth size, arch length discrepancies, overjet, and overbite are routinely measured and recorded. Plaster casts have been used for many decades and have proven to work effectively. However, they have several disadvantages, such as considerable space requirements for storage and the time needed to retrieve specific casts and also the risk of losing med casts. Three-dimensional digital model scanners and intraoral scanners are increasingly incorporated into orthodontic offices and/or in orthodontic research. They are used to construct a digital three-dimensional image of maxillary and mandibular arches from existing plaster models or impressions, thus making them a practical replacement for traditional plaster models.²⁹ With an intraoral scanner it is also possible to directly take an intraoral digital impression. These Three-dimensional digital models are used

for several purposes such as orthodontic measurements, along with treatment planning, and prosthetic and surgery planning. Intraoral scanning, computed tomography and extraoral scanning can all be used together for complete digital planning. Furthermore, three-dimensional digital model scanning is an indirect imaging technique where the physical plaster model or impression is scanned by a laser scanner and subsequently reconstructed as a digital file. In the last few decades the use of three-dimensional scanning has increased steadily. With the ‘new’ technology the orthodontist is able to digitise the patient’s casts and have an entirely computer-based patient record.

Significance

The series of studies on which this thesis is based was designed to evaluate the treatment effects, cost-effectiveness and impact on OHRQoL of posterior crossbite correction in mixed dentition, using an evidence-based approach. The results of these studies are expected to be beneficial for the patients who will be offered the most widely accepted and effective treatment, which is of importance for the dentists for decisions as to which treatment will give the best outcome, and beneficial for the dental-care providers and society in care planning and the allocation of resources.

AIMS

Paper I

To conduct a systematic review of published studies of health economics in orthodontics, assessing the evidence from studies presenting an analysis of costs associated with treatment outcome.

Paper II

To investigate the OHRQoL among 9-year-old children in mixed dentition and to compare the OHRQoL of children with unilateral posterior crossbite (UPC) to children with excessive overjet (EO) as well as with children with normal (NO) with no or mild orthodontic treatment need.

Paper III

To compare the costs of the quad-helix and removable expansion plate for the treatment of unilateral posterior crossbite with functional shift in mixed dentition as well as to evaluate the costs of such crossbite treatments performed either in specialist or general dentistry.

Paper IV

To assess the three-dimensional treatment changes (specifically palatal surface area and volume) of forced unilateral posterior crossbite correction using either quad-helix or removable expansion plate appliances in mixed dentition, and to compare the treatment changes with the three-dimensional changes occurring in age-matched untreated unilateral posterior crossbite patients as well as in subjects with normal occlusion and with no or mild orthodontic treatment need.

HYPOTHESIS

Paper I

Economic evaluation has become an integral component of health services research in recent years, despite this there is only limited evidence on health economic evaluations in orthodontics.

Paper II

The hypothesis was that children with excessive overjet have lower OHRQoL than children with unilateral posterior crossbite with functional shift or children with no or mild orthodontic treatment need since excessive overjet occurs in the aesthetic zone.

Paper III

It was hypothesised that both treatment methods are equal regarding costs as well as for the costs of treatment in general and specialist dentistry.

Paper IV

It was hypothesised that there are no significant differences regarding palatal surface area and volume between patients treated with the quad-helix and expansion plate. Furthermore, it was hypothesised that there were no significant morphological changes in the untreated control group. It was also hypothesised that the palatal surface area and volume in the treated groups would not reach the dimensions of the normal group.

MATERIALS AND METHODS

Paper I was a systematic review conducted in four steps, according to Goodman's model, in order to identify all studies evaluating economic aspects of orthodontic interventions. The search covered the databases Medline (<http://www.ncbi.nlm.nih.gov>), CINAHL, Cochrane, Embase, Google Scholar, National Health Service Economic Evaluation Database, and SCOPUS, for the period of 1966 to September 2014. The inclusion criteria were: RCTs or controlled clinical trials comparing at least two different orthodontic interventions, evaluation of both economic and orthodontic outcomes, and study populations of all ages.

In **Paper II** the patients were sourced from 19 Public Dental Service Clinics in Sweden. The sample comprised baseline data from children that originate from two different controlled trials, one regarding unilateral posterior crossbite with functional shift (UPC) and the other focusing on excessive overjet combined with Class II malocclusion (EO). Both trials also include children with normal occlusion and with no or mild orthodontic treatment need (NO).

The UPC group comprised 93 children (45 boys and 48 girls), the EO group 71 children (36 boys and 35 girls), and the NO group 65 children (32 boys and 33 girls). In conjunction to the clinical examination, all children completed the Child Perceptions Questionnaire (CPQ8-10) for evaluation of OHRQoL. The CPQ8-10 comprises 25 questions grouped into four domains:

oral symptoms, functional limitations, emotional, and social well-being. (Appendix A) Validated questions about pain in the jaws and face were also included (Appendix B).

The four treated groups in the UPC group; QH treatments in specialist orthodontic clinics (QHS), QH treatments in general dentistry (QHG), EP treatments in specialist orthodontic clinics (EPS), and EP treatments in general dentistry (EPG), were asked to fill out the CPQ questionnaire 6 weeks in treatment and one month after debonding. The untreated crossbite group (UC) and the normal group (NO) filled out the CPQ questionnaire before treatment and one year after the initial recording.

The ethics committee of Lund University, Lund, Sweden (Dnr: 2012/650), which follows the guidelines of the Declaration of Helsinki, approved the informed consent form and protocol for Paper II-IV. In Paper II the trial protocol and informed consent were also approved by the Regional Ethical Review Board in Linköping (Dno: 2012/302-31), Sweden.

Thus, the patients and parents received oral and written information about the trial and then signed the consent form.

This trial was registered at “FoU I Sverige” (<https://www.researchweb.org/is/sverige>), registration number 220751.

For **Paper III** the subjects were recruited from 10 different Public Dental Health Services, In Halland County Council, Sweden.

Figure 2. Flowchart Paper III.

All patients met the following criteria; mixed dentition (all incisors and first molars erupted), unilateral posterior crossbite with functional shift including the maxillary first molar, no sucking habits or sucking habit ceased at least one year before the trial. The exclusion criteria were: craniofacial syndromes and children who had previously undergone orthodontic treatment. A total of 110 patients, 59 girls and 51 boys, mean age 9.2 years (SD 1.02), entered the study and were randomised into four groups (Figure 2). The randomisation procedures were prepared and carried out by an independent person not involved in the study and the randomisation used blocks of 20. Thus, the orthodontists carried boxes of pre-prepared envelopes. Six opaque envelopes were prepared with 20 sealed notes in each (five notes for each group) and, for every new patient in the study, a note was extracted from the first envelope. When the envelope was empty, the second envelope was opened, and the 20 new notes were extracted as the patients were recruited to the trial. This procedure was then repeated four more times. After enrolment and randomisation,

all patients were treated according to a present standard concept, detailed descriptions of the design and handling of the appliances (see the quad-helix and the removable expansion plate), and also a form to fill out, for example, time for ordinary and emergency visits, parents' travelling time and absence from work.

Accordingly, the quad-helix treatments at the specialist orthodontic clinics (QHS) included 11 boys and 17 girls (mean age 9.3, SD 1.14), the quad-helix treatments at the general dentistry clinics (QHG) involved 13 boys and 14 girls (mean age 9.6, SD 1.05), the removable expansion plates treated at the specialist orthodontic clinics (EPS) comprised 14 boys and 13 girls (mean age 8.8, SD 0.85) while the removable expansion plates treated at the general dentistry clinics (EPG) consisted of 13 boys and 15 girls (mean age 9.0, SD 0.87).

ITT

In Paper III and IV we used the intention-to-treat principle (ITT), i.e. data on all patients who were randomly assigned to different groups, whether successful or not as well as dropouts, were included in the final analysis. If the crossbite was not corrected within one year, the outcome was defined as unsuccessful. In addition, any drop-outs were considered to be unsuccessful.

Cost-minimisation analysis

A cost-analysis was performed with reference to the intention-to-treat (ITT) principle, regarding direct costs, indirect costs and societal costs. A cost-minimisation analysis was used to describe the situation since the consequences of the two treatments were broadly equivalent, and thus the difference between them was reduced to a comparison of costs. The sum of direct and indirect costs was defined as 'societal costs' and the cost minimisation was calculated as follows:

1. The mean societal costs of successful cases only on completion of treatment within one year in all groups:
 - the mean societal costs for all successful QH treatments at the specialist orthodontic clinic divided by the number of successful patients, i.e. societal costs for 28 treatments divided by 28 patients.

- the mean societal costs for all successful QH treatments at the general clinic divided by the number of successful patients, i.e. societal costs for 23 treatments divided by 23 patients.
 - the mean societal costs for all successful EP treatments at the specialist orthodontic clinic divided by the number of successful patients, i.e. societal costs for 17 treatments divided by 17 patients.
 - the mean societal costs for all successful EP treatments at the general clinic divided by the number of successful patients, i.e. societal costs for 18 treatments divided by 18 patients.
2. The mean societal costs of the successful and retreated cases (cases that had to be retreated since no correction was achieved during one year) on completion of treatment in all groups:
- the mean societal costs for all successful treatments and retreatments of QH at the specialist orthodontic clinic divided by the number of successful patients, i.e. 28 treatments divided by 28 patients, thus no retreatments and unaltered costs.
 - the mean societal costs for all successful treatments and retreatments of QH at the general clinic divided by the number of successful patients, i.e. 27 treatments divided by 23 patients, thus 4 retreatments and an increase in costs.
 - the mean societal costs for all successful treatments and retreatments of EP at the specialist orthodontic clinic divided by the number of successful patients, i.e. 27 treatments divided by 17 patients, thus 10 retreatments and an increase in costs.
 - the mean societal costs for all successful treatments and retreatments of EP at the general clinic divided by the number of successful patients, i.e. 28 treatments divided by 18 patients, thus 10 retreatments and an increase in costs.

Paper IV consisted of the patients from Paper III and a further 50 patients who served as controls, i.e. 25 patients with untreated crossbite patients and 25 age-matched children in mixed dentition

with normal occlusion and with no or mild orthodontic treatment need. The crossbite children were randomised into one of five groups: treatment in general dentistry clinics with quad-helix (QHG) or removable expansion plate (EPG), or in specialist orthodontic clinics with quad-helix (QHS), or removable expansion plate (EPS) or into an untreated control group (UC) where the treatment of posterior crossbite was postponed for one year (Figure 3).

Figure 3. Flowchart Paper IV

During the one-year trial period, the three-dimensional treatment changes (**Figure 4**) (palatal surface and projection area and surface volume), and two dimensional transverse linear (**Figure 5**) changes of forced unilateral posterior crossbite correction using either quad-helix or removable expansion plate appliances in mixed dentition were assessed, and then compared to the treatment changes with the three-dimensional changes occurring in age-matched untreated unilateral posterior crossbite patients as well as in subjects with normal occlusion.

Figure 4. Illustrations of surface area, projection area and surface volume on three- dimensional scanings in Paper IV

Figure 5. Two dimensional transverse linear measurements on study casts in Paper IV.

Complementary and new studies

New literature search

To update the systematic review a new survey was added spanning from October 2014 to April 2019, using the six databases described above on page 27. The same MeSH terms as in the original search were used in this updated version. The reference list of retrieved articles was also hand-searched with the same inclusion criteria, data collection and analysis as described in **Paper I**.

CPQ/Oral health related quality of life

To investigate if there were any differences in OHRQoL between the appliances of quad-helix and expansion plate pre-treatment, during treatment and posttreatment the patients were asked to fill in the CPQ questionnaire before treatment, after 6 weeks of treatment and one month after having completed treatment. The same CPQ8-10 questionnaire used in Paper II was utilised for this supplementary investigation.

Three-dimensional changes in two groups

In order to evaluate if there existed any three-dimensional differences totally and without regard to whether treatment was performed in general or specialist dental care between quad-helix and expansion plate treatments, the two groups QHS and QHG were combined to one quad-helix group, QH, and the same was done with EPS and EPG, which were merged in to one group, EP.

Statistical analysis Paper II-IV

The statistical analysis was carried out using SPSS (IMB SPSS® Version 24.0 IBM Corp., Armonk, USA). A *P* value of less than 5% ($P < 0.05\%$) was considered statistically significant. Means between groups were compared using ANOVA with Tukey's post hoc test in Paper II-IV. In Paper II, a general linear model in combination with Tukey's test was performed when corrections were made for background/confounding variables: gender, caries, enamel defects, headache and trauma. In Paper II-IV, chi-square analysis and Fisher's exact test were used to determine differences between groups regarding categorical data, e.g. to calculate the success rate of crossbite correction.

Sample size calculation

In **Paper II**, the test was two-tailed, which means that an effect in either direction was interpreted. With a proposed sample size of 60 in each group, the trial had a power of 81.6% ($1 - \beta$) to yield a statistically significant result. This computation assumed that the difference in proportions was 0.25 (25% difference of quality of life between the groups). The difference of 25% was selected as the smallest difference in the sense that any smaller difference would not be of clinical or substantive significance. It was also assumed that this effect size was reasonable in the sense that an effect of this magnitude could be anticipated in this field of research.

In **Paper III**, and to detect a relevant cost difference of 1000 SEK/€96 standard deviation (SD): 100/96), and for a power of 90% at a significance level of $P > 0.05$, it was calculated that the sample size needed to be 21 patients per group. To compensate for possible dropouts of approximately 20% per group the number of patients per group was increased to at least 25.

In **Paper IV**, as Primožič et al.³⁰ are the only authors who have evaluated three- dimensional palatal vault changes in assessing treatment of constricted maxilla in growing subjects, we used their sample size calculation as a base for clinical relevant changes. With the assumption of means in variances for different groups taken from Primožič et al.³⁰, and using multiple testing adjustment suggested by Dunnett and Tamhane³¹, with standard parameters of 80% power and α 0.05, a number of 23 children per group was needed, and as we expected some dropouts to occur, the required sample size was estimated to be 25 in each of the groups.

RESULTS

Paper I - Systematic review

The applied terms for searches yielded 1,838 studies, of which 989 were excluded as duplicates. Application of the inclusion and exclusion criteria identified 26 eligible studies for which the full-text versions were retrieved and scrutinised.

Reason for exclusion

- Not following the objective of the review (648)
- Descriptive (18)
- Editorial (64)
- Opinion (42)
- Review (45)
- Case report (6)

At the final analysis, eight studies remained^{25,32-38}. Out of these eight studies, four came from Sweden^{25,32,33,38}, two from the Netherlands^{36,37} and two from the UK^{34,35}. Of these studies there were four of different orthodontic appliances^{25,33,34,38}, two on retainers^{32, 35} and one on infant orthopaedic treatment³⁶, and, finally, one study on cleft palate treatment³⁷. Of the four studies on appliances^{25, 33,34,38}, only two included comparable forms of appliance treatment, mainly crossbite treatment using quad-helix or expansion plates.^{25,38}

Type of study

Three studies were based on cost-effectiveness analyses³⁵⁻³⁷ and the other five^{25,32-34,38} on cost-minimisation analysis. Two of the cost-minimisation studies^{25,32} included a societal perspective, i.e. the sum of direct and indirect costs. The aims of most of the studies varied widely and of studies comparing equivalent treatment methods few were of sufficiently high study quality.

Evaluation of study quality and evidence

The quality of the studies was graded as limited, medium or high according to the GRADE system. The quality of four^{33, 36-38} of the eight studies was assessed as limited, three^{25,32,34} as moderate, and one as high³⁵. The reason for grading four articles as limited was primarily due to inconsistency/lack of transparency in how the patients were selected and if the number of patients needed was sufficient and the fact that two had a retrospective study design. The RCT study from the UK³⁵ presented high transparency and included almost 400 patients, and was assessed as being of high quality. Thus, the literature provides an inadequate evidence base for the economic aspects of orthodontic treatment.

Table 1. Summary of studies included in the quality assessment; listed in descending order, from the latest publication.

Author, year of publication and country	Aim of the study	Study design Follow up	Study population	Economic and orthodontic outcomes	Study quality
Edman Tynelius et al. 2014 (32) Sweden	Compare costs for three retention methods; vacuum formed retainer + canine to canine retainer (VCTC), vacuum formed retainer + stripping retainer (V-S) and positioner (P)	Costs-based piggy-back on one earlier published RCT 2 years follow-up	Adolescents 17-19 years of age. Group VCTC: N = 24, 17 girls and 7 boys. 1 dropout. Group V-S: N = 23, 14 girls and 9 boys. 2 dropouts. Group P: N = 22, 13 girls and 9 boys. 3 dropouts.	Cost-minimisation analysis. Direct, indirect and societal costs. Equally efficient in retaining orthodontic treatment results Costs in Euro per patient: VCTC 457 V-S 451 Positioner 429 CTC retainer was the least cost effective.	Estimated treatment times. Average times used for calculations. Societal perspective. Intention-to-treat. Low attrition. Moderate study quality.
Petrén et al. 2011 (25) Sweden	Compare costs of unilateral posterior crossbite correction using quad-helix (QH) or expansion plate (EP)	Costs-based piggy-back on one earlier published RCT 3 years follow-up	Group QH: N = 20, mean age 9.0 yrs. (range 7.8-10.2 yrs.), 11 girls and 9 boys. Group EP: N = 20, mean age 8.5 yrs. (range 7.5-9.5 yrs.), 13 girls and 7 boys.	Cost-minimisation analysis. Direct, indirect and societal costs. Success rate of correction, transverse expansion. QH and EP had equally successful crossbite corrections. QH offers significant economic benefits over EP. QH had lower direct and indirect costs, also fewer failures needing retreatment. Costs in Euro QH 1031 vs. EP 1395 (p<0.01)	Unclear about variability of some costs compared to treatment time. Societal perspective. Intention-to-treat analysis No attrition. Moderate study quality.

(Table 1 continues on next page)

(Table 1 cont.)

Laag et al. 2009 (33) Sweden	Compare results and costs at a general dentist clinic (GDC) and a special orthodontic clinic (SOC) for treatment of crowding, increased overjet, lateral crossbite and dental displacement.	Retrospective CCT. Cases at GDC and controls at SOC Follow-up unknown	Removable appliance at GDC and fixed appliance at SOC. Group GDC: N = 31, mean age 11.4 yrs. (range 8.2-14.2 yrs.), 20 girls and 11 boys. Group SOC: N = 19, mean age 13.1 yrs. (11.1-14.2 yrs.), 16 girls and 3 boys.	Cost-minimisation analysis. Weighted Peer Assessment Rating (WPAP-index), GDC 69.0 (+/-10.6) vs. SOC 84.1 (+/- 7.1) (p<0.001) Costs SEK GDC 3.525 vs. SOC 5.510 Trained general dentists may achieve sufficiently qualitative results at a reasonably low cost. The results tend to be better at the SOC but more expensive than at the GDC.	Simplified calculations of costs. Retrospective CCT. Non-cooperative patients were not registered, thus only successful treatment was counted. Limited study quality.
O'Brien et al. 2009 (34) United Kingdom	Evaluate the effectiveness and costs of early vs. adolescent orthodontic treatment of Class II division 1 malocclusion.	Multi-centre RCT 10 years follow-up	Group early twin-block appliance: N = 89, mean age at start 9.7 yrs. (SD 0.98), 41 girls and 48 boys. Removable or fixed appliance for the adolescent group: N = 85, mean age 9.8 yrs. (SD 0.98), 39 girls and 46 boys. Early treatment group: mean age 12.4 yrs. (SD 1.16) at second treatment phase. Adolescent treatment group: mean age 12.1 yrs. (SD 1.1).	Cost-minimisation analysis. No difference with respect to skeletal pattern, extraction rate, patient self-esteem. Early twin-block treatment incurred higher costs 3,913 vs. 3,018 \$/patient. Early treatment with removable appliance followed by further treatment in adolescence at the appropriate time does not result in any meaningful long-term benefit over one course of treatment started in the late mixed dentition.	No sensitivity analysis. Intention-to-treat analysis. 19% drop-out rate. Moderate study quality.

Hichens et al. 2007 (35) United Kingdom	Compare cost-effectiveness of Hawley retainers (HR) and vacuum-formed retainers (VFR).	RCT 6 months follow-up	Group HR: N = 196, mean age 14.8 yrs., 124 girls 92 boys. Group VFR: N = 201, mean age 15.0 yrs., 119 girls and 82 boys.	Cost-effectiveness analysis. VFR was more cost-effective than HR from all perspectives, i.e. National Health Service, the practices and the patients.	Well-performed and good transparency of calculations. Power analysis. Drop-outs presented. High study quality.
Konst et al. 2003 (36) The Netherlands	Investigate the cost-effectiveness of infant orthopaedic treatment (IO) of unilateral cleft lip and palate compared with no such treatment (NT) on speech development.	Three-centre RCT 2 years follow-up	20 patients randomised to two groups. Group IO: N = 10, age 2.5 yrs., 2 girls and 8 boys. Group NT: N = 10, age 2.5 yrs., 1 girl and 9 boys.	Cost-effectiveness analysis. IO: Cost of treatment by orthodontists per patient €1,460 NT: €419. ICER Incremental cost effectiveness ratio, €1,041 per 1.34-point speech quality improvement. Regarding speech in cleft patients, IO group had a statistically significant higher rating than the NT group.	Selection of subjects unclear, i.e. unknown if consecutive patients. High attrition rate. Interpretation unclear Number of patients not corresponding to the power analysis in original study. Limited study quality.
Severens et al. 1997 (37) The Netherlands	Investigate cost-effectiveness of complete unilateral cleft palate treatment from presurgical orthopaedic treatment (PSOT).	Three-centre RCT PSOT compared with non-PSOT 18 weeks follow up	Group PSOT: N = 23, age 2 weeks, gender distribution not presented. Group non-PSOT: N = 20, age 2 weeks, gender distribution not presented.	Cost-effectiveness analysis. Surgical lip closure, no significant difference. PSOT US\$852 versus non-PSOT US\$304. Regarding the incremental cost-effectiveness ratio, PSOT was more expensive than non-PSOT and had an insignificantly lower effect.	No information about attrition. Number of patients not corresponding to the power analysis in original study. Limited study quality.

(Table 1 continues on next page)

(Table 1 cont.)

Hermanson et al. 1985(38) Sweden	Compare effects, complications, discomfort and costs of unilateral crossbite correction using quad-helix appliance (QH) and Removable plate appliance (RP).	Retrospective CCT 2 years follow-up	Group QH: N = 27, mean age 9.8 yrs. (range 7.3 – 13.0 yrs.), 14 girls and 13 boys. Group RP: N = 17, mean age 9.5 yrs. (range 6.8 – 14.3 yrs.), 12 girls and 5 boys.	Cost-minimisation analysis Successful treatment: QH 26 of 27 and RP 16 of 17 Complications equal. Costs QH: 1,223 SEK versus RP 1,714 SEK. The cost analysis showed a 40% higher treatment cost with RP than QH. Treatment with RP takes longer than QH treatment.	Selection of subjects unclear, i.e. unknown if consecutive patients. Economic analysis not a cost-benefit analysis but cost- minimisation, thus the authors have labelled the economic analysis incorrectly. Retrospective CCT. Limited study quality.
---	--	--	---	---	---

New literature search

In May 2019, a supplementary literature search was performed in PubMed, CINAHL, Cochrane, Google Scholar and the National Health Service Economic Evaluation Database. The search was conducted from October 2014 (the original search ended in September 2014) and up to April 2019. This complementary search yielded a further 159 articles but only four³⁹⁻⁴² studies fulfilled the inclusion criteria. The reasons for exclusion and the number of excluded articles are listed in **Figure 6**.

Figure 6. Flow of information through the different phases of the updated systematic review.

Study design

All studies were cost-minimisation analyses comparing two different appliances to correct a malocclusion. Out of the four studies, one concerned treatment of unilateral posterior crossbite³⁹, one study was on anchorage loss⁴⁰, one study on large overjet reduction⁴¹, and one study on anterior crossbite treatment⁴². All four studies came from Sweden and all studies were RCTs.³⁹⁻⁴² A quality analysis of the four studies is shown in Table 2.

Evaluation of study quality and evidence

Several well-structured and well-performed articles in this field have been published since September 2014, most of which are from Sweden. There were different aims in all the studies evaluated except for the articles from Petrén et al.²⁵ and Sollenius et al.³⁹. Although the studies derive from the same university and research group there is strong evidence that the quad-helix is more cost-effective than the expansion plate in treating crossbites in mixed dentition.

Table 2. Summary of studies included in the quality assessment; listed in descending order, from the latest publication.

Author, year of publication and country	Aim of the study	Study design Follow-up	Study population	Economic and orthodontic outcomes	Study quality
Sollenius et al. 2019 (39) Sweden	To evaluate the costs of quad-helix (QH) and removable expansion plate (EP) treatments performed either in specialist or general dentistry for the correction of unilateral posterior crossbite with functional shift in mixed dentition.	Costs based on one RCT	Children in early mixed dentition 8-10 years of age. Group QHS (quad-helix in specialist dentistry): N = 28, 17 girls and 11 boys. Group QHG (quad-helix in general dentistry): N = 27, 14 girls and 13 boys. Group EPS (expansion plate in specialist dentistry): N = 27, 13 girls and 14 boys. Group EPG (expansion plate in general dentistry): N = 28, 15 girls and 13 boys. 0 dropout	Cost-minimisation analysis. Direct, indirect and societal costs. Success rate of correction. QHS had 100% successful crossbite correction. QHS offers significant economic benefits over the other three groups.	Well-performed and good transparency of calculations. Power analysis. Drop-outs presented. ITT. Societal perspective. High study quality.
Ganzer et al. 2019 (40) Sweden	To analyse the cost-effectiveness of anchorage reinforcement with buccal miniscrews (Group A) and with molar blocks (Group B).	Costs-based piggy-back on one earlier published RCT.	Adolescents 11-19 years of age. Group A: N = 40, mean age 16.4 yrs. (SD 1.7), 26 girls and 14 boys. Group B: N = 40, mean age 15.0 yrs. (SD 1.8), 26 girls and 14 boys.	Cost-minimisation analysis. Direct, indirect and societal costs. The difference in anchorage loss primarily occurred during space closure and was statistically significant.	Well-performed and good transparency of calculations. Power analysis. Drop-outs presented. ITT. Only question mark was that the calculation for loss of income was based on the assumption that the adults accompanying the patient took time off from work. High study quality.

(Table 2 continues on next page)

(Table 2 cont.)

Čirgić et al. 2018 (41) Sweden	Compare results and assess and relate the societal costs of reducing large overjet with a prefabricated functional appliance (PFA), or a slightly modified Andersen activator (AA), using a cost-minimisation analysis (CMA).	Cost-based piggy-back? on one earlier published RCT Follow-up	Removable appliance at GDC. Children at 6-14 years of age. >6 mm overjet, c II. Mean age 10.3 years, 44 girls, 53 boys Group PFA: N = 57. Group AA: N = 40. At each clinic, two envelopes were available, one for girls and one for boys with five AA and five PFA notes for both genders.	Cost-minimisation analysis. Direct, indirect and societal costs. Overjet ≤ 3 mm = success. Low success in both groups.	ITT. Drop-outs presented. Retreatment not included in cost-minimisation. Power analysis Individual numbers on indirect costs. High study quality.
Wriedel et al. 2015 (42) Sweden	Evaluate and compare the costs of correcting anterior crossbite with functional shift, using fixed appliance (FA) or removable appliances (RA). Cost-minimisation	Two-centre RCT 2 years follow-up. Blocks of 10	62 patients randomised to two groups. 37 boys and 25 girls, mean age 9.8 years Group FA: N = 31, age 2.5 yrs., 2 girls and 8 boys. Group RA: N = 31, age 2.5 yrs., 1 girl and 9 boys.	Cost-minimisation. FA: Cost of treatment by orthodontists per patient € 630 (success), € 678 (including retreat). RA: € 945 (success), € 1,031 (including retreat). Significant economic benefits for FA over RA. Even when only successful outcomes were assessed, treatment with RA was more expensive.	No power analysis. Drop-outs presented. ITT. Only question mark was that the calculation for loss of income was based on the assumption that the adults accompanying the patient took time off from work. High study quality.

Paper II Quality of life

In total, 229 patients were included in the three groups. All children in the unilateral posterior crossbite group (UPC) and normal occlusion group (NO) had Class I molar relation while all in the excessive overjet group (EO) showed Class II division 1 malocclusion. A significantly larger overjet ($P < 0.001$) and overbite ($P < 0.001$) was found in the EO group compared to the UPC group. There was also a significantly larger overjet ($P < 0.001$) in the EO compared to the NO group. There was a significantly larger number of children without lip closure ($P < 0.001$) in the EO group compared to the other two groups. All children in the UPC group had moderate to severe treatment need (IOTN 3-4) indicating that it was a forced crossbite with a shift of the mandible to the crossbite side. In the EO group, all patients had severe to extreme treatment need (IOTN 4-5) due to the large overjet, while in the NO group the patients showed no or mild orthodontic treatment need (IOTN 1-2). The prevalence of trauma, enamel defects, and headache was low in all three groups, and without any significant inter-group difference.

Oral health related quality of life (OHRQoL)

The results show that the total CPQ mean score for EO was significantly larger ($P = 0.048$) compared to UPC and also compared to the NO ($P = 0.012$). No gender differences were found.

Emotional and social well-being

The children with EO reported significantly higher scores in the domains of emotional well-being compared to UPC ($P = 0.039$) and social well-being compared to NO ($P = 0.012$).

Confounders

Different background variables may have a negative impact on the OHRQoL, and, therefore, such variables that could have an effect, e.g. gender, caries, trauma, enamel defects and headache were checked. When corrections were made for these variables/possible confounders, the significant differences between the groups still remained. Furthermore, headache was the only variable that showed a significant negative impact on the domain of oral symptoms ($P = 0.05$).

Results from the complementary and new study of CPQ values

The results from the CPQ questionnaire, before T0, during T1 and after treatment T2 in the unilateral posterior crossbite group have not yet been published in any peer-reviewed journal. This is so far a preliminary reporting of achieved treatments and the number of participants is not sufficient to reach the power that we have wanted, but still it is interesting to evaluate how treatment per se and different appliances may affect the crossbite children.

The total CPQ mean score was 5.3 at T0 for all the treated groups (QHS, QHG, EPS, EPG), at T1 the total mean score was 7.5 and at T2 it was 4.9. The total CPQ mean score for UC (untreated control) was 5.4 at T0 and 5.7 at T2, and for NC (normal control) the mean score was 3.3 at T0 and 3.5 at T2.

In the treated crossbite groups there was a significant difference between T0 and T1 and between T1 and T2 ($P=0.000$). There was no significant difference between T0 and T2.

The quad-helix treatments in general dentistry QHG had 6 weeks in treatment (T1), with a significantly higher total mean CPQ score than the expansion plate treatments in specialist orthodontic clinics EPS ($P=0.032$), (Table 3). For the QHG the mean subscale score at T1 was 3.9 for functional limitation (FL), for quad-helix treatments in specialist orthodontic clinics (QHS) QHS 2.1, for EPS it was 0.9 and for expansion plate treatments in general dentistry EPG 0.8. The patients in the QHG group reported significantly higher scores in the domains FL compared to QHS ($P=0.046$), EPS ($P=0.000$) and EPG ($P=0.001$). For the domains OS, EW and SW, no significant differences were detected between the groups.

At T0 the total mean CPQ score was significantly higher for the treated crossbite group (QHS+QHG+EPS+EPG) compared to the normal group (NC) ($P=0.001$), after treatment T2 there was no significant difference found between the total mean CPQ score for the crossbite group and for the NC group.

At T0, T1 and T2 there are no significant differences in total mean CPQ scores between treatments performed in specialist orthodontic clinics compared to general dentistry.

At T1 there is a significantly higher total mean CPQ score (P=0.003) in the QH group (QHS+QHG) compared to the expansion plate group (EPS+EPG).

Regarding CPQ scores there seem to be small differences regardless of which appliance is used or at which treatment instance the treatment is performed; however, somewhat higher CPQ scores are seen 6 weeks in treatment (T1) using the quad-helix compared to expansion plate.

The crossbite appliances are effective in reducing the CPQ score since after crossbite correction the patients' CPQ value is reduced to the same level as the normal patients.

Table 3. Showing CPQ₈₋₁₀ scores for T0, T1 and T2.

	N		CPQ Total		Oral symptoms (OS)		Functional limitations (FL)		Emotional well-being (EW)		Social well-being (SW)		Difference
			Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
T0	28	QHS	4.6	4.25	3.1	2.71	0.8	1.62	0.5	1.37	0.6	1.21	N.S
	20	QHG	5.6	4.06	3.7	2.83	1.3	2.02	0.6	1.00	0.2	0.48	
	27	EPS	5.0	5.30	3.4	2.76	0.7	1.46	0.3	1.18	1.1	1.68	
	27	EPG	5.7	3.38	3.8	2.52	0.9	1.53	1.0	2.66	0.5	1.25	
	19	UC	5.4	6.14	3.0	2.24	1.0	1.65	0.7	1.46	1.0	1.93	
	28	NC	3.3	2.66	2.4	2.97	0.5	0.93	0.4	1.06	0.0	0.00	
T1	28	QHS	7.9	5.52	4.4	3.00	2.1	2.51	0.6	0.92	0.8	1.12	QHGS>EPS = 0.032 FL: QHGS>QHS = 0.046 QHGS>EPS = 0.000 QHGS>EPG = 0.001
	17	QHG	11.1	10.15	4.5	3.86	3.9	3.28	1.3	1.86	1.4	2.23	
	27	EPS	5.7	4.84	3.4	3.06	0.9	1.50	0.2	0.43	1.0	1.06	
	27	EPG	6.4	4.49	2.6	2.29	0.8	1.21	0.2	0.50	0.5	0.84	
		UC	-	-	-	-	-	-	-	-	-	-	
		NC	-	-	-	-	-	-	-	-	-	-	
T2	28	QHS	4.1	3.96	2.4	2.24	0.5	0.79	0.6	1.77	0.5	0.88	N.S
	16	QHG	4.6	3.83	2.8	2.57	1.0	2.07	1.1	1.39	0.8	1.38	
	27	EPS	4.9	4.76	2.4	1.63	0.6	1.16	0.1	0.44	0.6	1.08	
	27	EPG	5.9	4.53	2.6	2.45	0.7	1.42	0.2	0.38	0.2	0.50	
	19	UC	5.7	5.60	3.0	2.15	0.7	1.06	0.2	0.71	0.5	1.43	
	8	NC	3.5	1.69	2.8	1.28	0.5	0.76	0.1	0.35	0.0	0.00	

Paper III Cost-minimisation

Successful treatments

In the quad-helix specialist group (QHS), 28 of 28 patients were successfully corrected compared to 23 of 27 in the quad-helix general (QHG) group. Treatment with expansion plate was less successful: 18 of 27 in the expansion plate specialist (EPS) group and 18 of 28 in the expansion plate general (EPG) group. Consequently, QHS was significantly more successful compared to EPS ($P = 0.001$) and EPG ($P = 0.000$).

Number of appointments

When including scheduled and emergency appointments as well as retreatments there were significantly lower numbers of appointments found in the QHS group compared with EPS ($P = 0.007$) and EPG ($P = 0.000$), also QHG had significantly fewer appointments compared with the EPG ($P = 0.032$).

Not a single quad-helix was lost in the two QH groups; however, 33 expansion plates had to be reproduced due to lost or broken appliances. Thus, the reversal of 13 new plates in the EPS and 20 in the EPG group implied to increased material cost for these groups, only the increased material costs for broken/lost plates were for EPS € 1875 and for EPG € 2884.

Societal costs - successful treatments

The mean societal costs, i.e. direct and indirect costs per patient, with successful quad-helix treatment were significantly lower for QHS than QHG ($P = 0.000$). For the cost in the expansion plate groups the EPS was significantly lower compared with EPG ($P = 0.015$). (Table 4)

Material costs - successful treatments

The mean material cost per patient was significantly lower for the quad-helix compared to expansion plate treatments in special as well as in general dentistry ($P = 0.000$). In addition, the material costs of expansion plate treatments in special orthodontic clinics were lower than plate treatments in general dentistry ($P = 0.013$).

Treatment time costs - successful treatments

The QHG costs were significantly higher compared to both QHS and EPS ($P=0.000$). Furthermore, the treatments' time costs were significantly higher than EPG ($P=0.000$).

Indirect costs - successful treatments

The indirect cost per patient was significantly lower for the QHS compared with QHG ($P = 0.008$). For all the groups the indirect cost per patient was 5% to 10% of the societal costs.

Costs including retreatments

The lower success rate for expansion plate treatments performed in general and specialist dentistry as well as for the quad-helix in general dentistry caused the increased costs on all levels for these groups. Since quad-helix treatments in the specialist orthodontic clinic had a success rate of 100%, we added this cost to the unsuccessful treatments when the costs including retreatment were calculated.

Mean societal, direct and indirect costs (Euros) for the quad-helix treatment in the specialist dentistry (QHS), quad-helix in the general dentistry (QHG), removable expansion plate treatment in specialist dentistry (EPS), and removable expansion plate treatment in general dentistry (EPG). Mean total costs per patient for successful patients only and after retreatment. There were no retreatments in the QHS group while there were four in the QHG and 10 retreatments in each of the EPS and EPG groups. Thus explaining the increased costs in the QHG, EPS and EPG groups for including retreatments compared to the successful treatments.

Table 4. Mean costs in Euros for the QHS, QHG, EPS and EPG for successful patients only and after re-treatment, QHS = 0 patients, QHG = 5 patients, EPS = 10 patients and EPG = 10 patients. The added costs for retreatments are the cost for one successful quad-helix treatment in specialist clinics (QHS).

	Successful only	Including re-treatment
Quad-helix in specialist clinics (QHS)		
Societal costs	765	765
Direct costs - material	120	120
Direct costs - time	601	601
Indirect costs	44	44
Quad-helix general dentistry (QHG)		
Societal costs	1025	1139
Direct costs - material	120	138
Direct costs - time	788	877
Indirect costs	117	124
Expansion plate specialist clinics (EPS)		
Societal costs	844	1099
Direct costs - material	185	226
Direct costs - time	575	775
Indirect costs	83	98
Expansion plate general dentistry (EPG)		
Societal costs	878	1151
Direct costs - material	229	272
Direct costs - time	569	784
Indirect costs	80	95

Paper IV Treatment effects

A total of 160 children entered the study, and, of these, 135 children with crossbite were randomised into five groups; quad-helix treatments in specialist orthodontic clinics (QHS), quad-helix treatments in general dentistry (QHG), removable expansion plate treatments in specialist orthodontic clinics (EPS), removable expansion plate treatments in general dentistry (EPG) and untreated crossbite group (UC). Parallel to these five groups, 25 children with normal occlusion were followed and constituted the normal group (NC) (Table 5).

Table 5. Baseline/pre-treatment demographic characteristics. QHS: quad-helix treatments in specialist orthodontist clinics, QHG: quad-helix treatments in general dentist clinics, EPS: removable expansion plate treatments in specialist orthodontist clinics, EPG: removable expansion plate treatments in general dental clinics. UC: untreated posterior crossbite control group, NC: normal control group with normal occlusion and no or mild orthodontic treatment need.

Group	Baseline/pre-treatment					
	Number			Age (year)		SD
	Total	Boys	Girls	Mean	Range	
QHS	28	11	17	9.3	4.1	1.09
QHG	27	14	13	9.5	4.2	1.04
EPS	27	15	12	8.7	3.1	0.78
EPG	28	11	17	9.2	5.9	1.11
UC	25	16	9	8.5	2.4	0.57
NC	25	17	8	9.3	3.4	1.06

Three-dimensional baseline changes

At baseline or before treatment, all five crossbite groups had significantly less palatal surface and projection area as well as palatal shell volume compared to the normal group.

Three-dimensional expansion effect on the maxilla

After treatment, the surface and projection area and shell volume increased in the four treatment groups (QHS, QHG, EPS, EPG). However, in the untreated crossbite (UC) group and the normal occlusion (NO) group the surface and projection area along with the shell volume were more or less unaltered in size after one year (Table 6).

Table 6. Mean differences in surface area, projection area, and shell volume within and between the groups. QHS: quad-helix treatments in specialist orthodontist clinics, QHG: quad-helix treatments in general dentist clinics, EPS: removable expansion plate treatments in specialist orthodontist clinics, EPG: removable expansion plate treatments in general dental clinics. UC: untreated posterior crossbite control group, NC: normal control group with normal occlusion and no or mild orthodontic treatment need. One-way analysis of variance with Tukey's *post hoc* test.

	QHS	QHG	EPS	EPG	UC	NC	Difference
Maxilla Surface area (mm ²)	107.4	69.7	91.4	52.6	13.6	8.1	QHS, QHG, EPS>UC and NC $P<0.015$. QHS>EPG $P=0.012$.
Projection area (mm ²)	112.7	88.2	76.5	44.4	8.4	0.7	QHS, QHG, EPS>UC and NC $P<0.000$. QHS>EPG $P=0.000$. QHG>EPG $P=0.042$. EPS>EPG $P=0.045$.
Shell volume (mm ³)	599.9	440.1	697.3	404.7	161.2	147.0	QHS, EPS>UC and NC $P<0.06$.

After treatment, the four treatment groups reached the normal group values considering the surface and projection area together with the shell volume (Table 7).

Table 7. Surface area, projection, area, and shell volume at post-treatment. QHS, quad-helix treatments in specialist orthodontist clinics; QHG, quad-helix treatments in general dentist clinics; EPS, removable expansion plate treatments in specialist orthodontist clinics; EPG, removable expansion plate treatments in general dental clinics; UC, untreated posterior crossbite control group; NC, normal control group with normal occlusion and no or mild orthodontic treatment need. One-way analysis of variance with Tukey's *post hoc* test.

	QHS	QHG	EPS	EPG	UC	NC	Difference
Maxilla Post-treatment Surface area (mm ²)	1261	1276	1275	1220	1184	1300	QHS > UC $P=0.02$, EPS > UC $P=0.02$, NC > UC $P=0.002$.
Projection area (mm ²)	922	870	933	894	843	940	EPS > UC $P=0.02$. NC > UC $P=0.012$.
Shell volume (mm ³)	5423	5618	5758	5355	4976	5823	EPS > UC $P=0.011$, NC > UC $P=0.005$.

Three-dimensional expansion effect on the mandible

In the mandible, small or negligible and insignificant changes within and between the groups were noted considering surface and projection area as well as shell volume.

Two-dimensional baseline linear measurements

At baseline or before treatment, there were significantly smaller maxillary intermolar and intercanine distances except for distances between the intercanine cusp tips when the five crossbite groups were compared to the normal group. However, no significant differences were found between the crossbite groups.

Two-dimensional expansion effect on the maxilla

After treatment, the intermolar and intercanine distances in the maxilla significantly increased within all four treatment groups. The increase was mostly pronounced in the QHS and QHG groups which exceeded some of the values of the normal group (Table 8).

Table 8. Intermolar and intercanine distances (mm) after treatment. QHS, quad-helix treatments in specialist orthodontist clinics; QHG, quad-helix treatments in general dentist clinics; EPS, removable expansion plate treatments in specialist orthodontist clinics; EPG, removable expansion plate treatments in general dental clinics; UC, untreated posterior crossbite control group; NC, normal control group with normal occlusion and no or mild orthodontic treatment need. One-way analysis of variance with Tukey's *post hoc* test.

Post-treatment	QHS	QHG	EPS	EPG	UC	NC	Difference
Maxilla	Mean	Mean	Mean	Mean	Mean	Mean	
Intermolar distance, mesiobuccal cusp tips	52.2	52.3	52.0	51.5	47.9	50.8	UC<QHS, QHG, EPS, EPG, NC P<0.003
Intermolar distance, gingival margin	35.1	35.2	34.2	33.7	31.0	33.4	UC<QHS, QHG, EPS, EPG, NC P<0.002 NC<QHS, QHG P<0.05
Intercanine distance, cusp tips	33.5	33.2	33.4	32.5	29.9	31.7	UC<QHS, QHG, EPS, EPG P<0.01
Intercanine distance, gingival margin	25.8	25.5	25.6	24.9	23.1	25.6	UC<QHS, QHG, EPS, NC P<0.005

Two-dimensional expansion effect on the mandible

Small or negligible and insignificant changes were found in the mandible regarding changes of the intermolar and intercanine distances.

Complementary new results

When the results from the two quad-helix groups (QHS, QHG) and from the two expansion plate groups (EPS, EPG) were merged into one QH group and one EP group, some new significant differences and findings could be discovered. (Table 9).

Table 9. Mean differences in surface area, projection area, and shell volume within and between the groups. QH: quad-helix treatments in specialist orthodontist clinics and in general dentist clinics, EP: removable expansion plate treatments in specialist orthodontist clinics and in general dental clinics. UC: untreated posterior crossbite control group. NC: normal control group with normal occlusion and no or mild orthodontic treatment need. One-way analysis of variance with Tukey's *post hoc* test. SD indicates standard deviation and CI the confidence interval.

	Group QH			Group EP			Group UC			Group NC			Difference
	Mean	SD	CI%	Mean	SD	CI%	Mean	SD	CI%	Mean	SD	CI%	
Maxilla Surface area (mm ²)	88.9	59.96	72.7-105.1	71.6	66.72	53.6-89.7	13.6	54.96	-9.1-36.3	8.1	66.19	-19.2-35.5	QH>UC and NC $P=0.000$. EP> UC $P=0.001$. EP> NC $P=0.000$.
Projection area (mm ²)	100.7	67.69	82.4-119.0	60.1	56.09	45.0-75.3	8.4	31.66	-4.6-21.5	0.7	45.65	-18.1-19.6	QH> EP $P<0.001$. QH>UC and NC $P=0.000$. EP> UC $P=0.001$. EP> NC $P=0.000$.
Shell volume (mm ³)	521.4	370.43	421.3-621.6	548.3	506.64	411.4-685.3	161.2	402.86	-5.1-327.5	147.0	529.29	-71.5-365.5	QH>UC and NC $P<0.07$. EP> UC and NC $P<0.02$.

Consequently, for the three-dimensional variables' surface area, projection area and shell volume all increased significantly more in the treated groups compared to the untreated groups. Furthermore, the projection area increased significantly more in the merged quad-helix group compared to the merged expansion plate group. The increased projection area was also seen in Paper IV, where the biggest

increase was seen in the QHS and QHG groups, but in Paper IV there was also an increased palatal surface area, which was not seen when the groups were merged together. In contrast, the effect on the surface area and shell volume were more similar in the two treated groups without any significant differences, see Table 9. This is in contrast to the results from Paper IV where the most shell volume expansion was seen in the QHS and EPG groups.

The two-dimensional variables, maxillary intermolar and intercanine distances, increased significantly more in the quad-helix and the expansion plate groups compared to the untreated group untreated groups, this was also found in Paper IV, where QHS and QHG had the largest increase. The quad-helix group also had a larger expansion effect on the molars, whereas the effect on the canines was more similar between the groups and without any significant differences (Table 10). In Paper IV the effect on intercanine expansion was highest in QHS compared to the other treated groups.

Table 10. Mean differences in intermolar and intercanine distances (mm) within and between the groups. QH: quad-helix treatments in specialist orthodontist clinics or in general dentist clinics, EP: removable expansion plate treatments in specialist orthodontist clinics or in general dentist clinics, UC: untreated posterior crossbite control group, NC: normal control group with normal occlusion and no or mild orthodontic treatment need. One-way analysis of variance with Tukey’s *post hoc* test. SD indicates standard deviation and CI the confidence interval.

Maxilla	Group QH			Group EP			Group UC			Group NC			Difference
	Mean	SD	CI%	Mean	SD	CI%	Mean	SD	CI%	Mean	SD	CI%	
Intermolar distance, mesiobuccal cusp tips	4.5	2.04	4.00-5.07	3.0	1.94	2.41-3.51	0.5	0.36	0.34-0.65	0.2	0.67	-0.07-0.49	QH>EP <i>P</i> =0.000. QH>UC and NC <i>P</i> = 0.000. EP>UC and NC <i>P</i> = 0.000.
Intermolar distance, gingival margin	4.1	1.85	3.63-4.63	2.6	1.66	2.10-3.04	0.3	0.37	0.16-0.48	0.2	0.57	-0.04-0.43	QH>EP <i>P</i> =0.000. QH>UC and NC <i>P</i> = 0.000. EP>UC and NC <i>P</i> = 0.000.
Intercanine distance, cusp tips	3.4	1.89	2.88-3.99	2.6	1.74	2.12-3.18	0.3	0.77	-0.03-0.68	0.6	0.90	0.06-1.20	QH>UC and NC <i>P</i> = 0.000. EP>UC and NC <i>P</i> = 0.000.
Intercanine distance, gingival margin	2.5	1.44	2.10-2.94	2.0	1.46	1.58-2.46	0.4	0.56	0.13-0.64	0.5	0.81	-0.02-1.02	QH>UC and NC <i>P</i> = 0.000. EP>UC and NC <i>P</i> = 0.000.

DISCUSSION

The major aims of this thesis were to investigate how patients with unilateral posterior crossbite with functional shift should be treated in best possible way, both from the patients' perspective and from a societal and cost-effective point of view.

A high level of evidence was of high priority; therefore, for this purpose, the first step was to undertake a systematic review of the literature to evaluate the evidence on cost-effectiveness of posterior crossbite correction. Subsequently a study of patients' OHRQoL was performed, and furthermore, an RCT to analyse the effectiveness and costs. Moreover, by using digital technology, a three-dimensional evaluation of morphological changes of the jaws was also performed.

The main findings from the systematic review disclosed that few orthodontic studies have presented both economic and clinical outcomes. The malocclusions and appliances varied in the different studies; thus, accordingly, there was insufficient evidence available about the health economics of orthodontic interventions. The updated systematic review revealed that some new high-quality studies on clinical and economic outcomes were to be found in the literature. With the results from Paper III³⁹ on cost-effectiveness it can now be concluded that there is high evidence for treating unilateral posterior crossbite with functional shift in mixed dentition with quad-helix.

In the study of OHRQoL, the main finding was that patients with large overjet report lower quality of life compared to patients with unilateral posterior crossbite or patients with

normal occlusion with no or mild orthodontic treatment need. This study also showed that unilateral posterior crossbite with functional shift in mixed dentition has little effect on quality of life.

In addition, Paper III showed that treatment of unilateral posterior crossbite in mixed dentition is recommended to be performed by a specialist orthodontist using the quad-helix appliance, since the crossbite is corrected in a shorter treatment time, with fewer appointments, and thereby, is more cost-effective. It was also showed according to Paper IV that in unilateral posterior crossbite patients it was possible to correct and reach the arch width and palatal volume of the maxilla to the same level as the arch and palatal volume of a patient with normal occlusion without crossbite.

Methodological considerations

The James Lind alliance (JLA)⁴³ discusses “What are evidence uncertainties?”

The JLA definition of an evidence uncertainty is that:

- no up-to-date, reliable systematic reviews of research evidence addressing the uncertainty exist.
- up-to-date systematic reviews of research evidence show that uncertainty exists.

The systematic review (Paper I), including the supplementary search up to April 2019 was conducted in four steps, according to Goodman’s model⁴⁴ in order to identify all studies evaluating the economic aspects of orthodontic interventions. To minimise the risk of overlooking important information, highly regarded, comprehensive databases were searched and the reference lists of retrieved articles were hand-searched.

The quality assessment was performed according to protocols independently by the researchers and disagreement over assessment was resolved by discussion to achieve consensus. Study quality was denoted as limited, moderate, or high, according to the checklist proposed by Drummond et al.²³ The strength of the scientific evidence based on study quality from included studies was undertaken according to the GRADE system.⁴⁵

As mentioned above, the systematic review revealed insufficient evidence regarding health economics and orthodontics, but the question has arisen as to whether it is at all possible to achieve high levels of evidence regarding economic evaluations of orthodontic interventions. It has been stated that it is difficult to compare the costs in one country to another, primarily because problems of transferability: the costs incurred in studies from one country are seldom representative of other country, i.e. monetary variables are influenced by local factors such as insurance system, staff salaries, etc.

What could be done to enhance comparisons between economic studies of orthodontic interventions undertaken in different countries? One way would be to include specific data not only on economic results, but also on resource utilisation. For instance, this could entail specifications of time input by dentists and dental nurses, respectively, and specifications of how overhead costs were calculated, as a percentage or as absolute added costs. Studies from different countries could then be compared more easily. Despite the issue of transferability, assessment of studies identified in systematic reviews is important as the evidence may serve as a guideline for the planning and management of orthodontic interventions.

The supplementary search from October 2014 up to April 2019 showed that during the last 5 years a series of new studies has been undertaken. It should be noted that the new studies primarily come from Sweden and this may be seen in the context of the Swedish system of subsidised dental care including publicly funded orthodontic treatment for children and adolescents. Of these four new studies, three were considered to represent high-quality studies with low bias. The cost-minimisation analysis from Petré et al.²⁵ 2013 together with the new study in Paper III revealed that the quad-helix appliance was less expensive, primarily because there were fewer failures needing re-treatment. However, there is still insufficient scientific evidence whether the treatment should be performed at specialist orthodontic or at general dentistry clinics since only one study on this topic has been performed. Thus, to reach a higher level of evidence more high-quality studies are required.

The results from Paper II show that patients with excessive overjet reported a lower self-perceived OHRQoL when compared with children with unilateral posterior crossbite.

Nonetheless, the scores for all the children participating in this study are generally low. This indicated that the children, even with the presence of a malocclusion with objective treatment need, perceive themselves of having a good OHRQoL. What could be the reason for this? The age factor might be one reason, the participants in this study were fairly young, 8-10 years old, and the CPQ scores shows that many of the crossbite patients are unaware of the malocclusion and this is in contrast to the excessive overjet group whose malocclusion is in the aesthetic zone.

It has often been described that patients seldom consider the degree of malocclusion and treatment need to be as high as the orthodontist will do.^{46,47} Other contributing factors to the high level of OHRQoL presented in Paper II could be the low caries prevalence and risk as well as the low numbers of patients showing dental fear and dental pain. All young children had, together with their parents, voluntarily accepted to participate in an additional clinical examination including impressions for study casts and all the children fulfilled the examinations and impressions without any problems and did not show any signs of dental anxiety. It can be assumed that patients with dental anxiety would probably decline to participate in such a study.

In Papers II-IV there was an even gender distribution while in many other studies there were more girls than boys receiving treatment or wanted treatment.^{48,49} O'Brien et al.⁵⁰ showed a strong association between gender and uptake of orthodontic treatment: being female increases the odds of receiving orthodontic treatment.

One might also wonder if it is necessary to treat in this early age since the child does not seem to be affected by the malocclusion. The correlation between future temporomandibular joint problems and posterior crossbites has been discussed in many studies^{8,9} but the evidence to support this is questionable. Whether the treatment is to be performed in early-, mixed- or permanent dentition has not been

an aim of this thesis, but it would be of great interest to analyse the eventual future facial asymmetries and temporomandibular disorders if the treatment was postponed until later in adolescence. In the light of early versus late treatment start, Berset et al.⁵¹ and Lagerström et al.⁵²) reported that, when 19-year-olds leave the free Public Dental Service in Sweden, some have persisting functional malocclusions or poor dental aesthetics. There were several explanations for this: One reason was that in early adolescence, the malocclusion traits had been assessed as mild deviations from the norm, for which no free orthodontic treatment was offered according to the Scandinavian model for treatment need. Another reason may be that at the time treatment was offered, in early adolescence, the individual was not concerned about the appearance and function of the dentition and declined the offer of orthodontic treatment. A third reason may be relapse of earlier treatment.

In Paper IV untreated control groups were used since crossbite correction is undertaken in the growing child. Hence, treatment changes must be evaluated in the context of changes associated with normal growth in children who have had no orthodontic treatment. Therefore, it was important to use an untreated control group as well as a normal group to relate to normal growth in Paper IV.

Intention-to-treat (ITT)

The advantages of using the RCT methodology include minimising the bias in treatment assignment, specifically selection bias and the confounding variables, hidden or out of control, will be of equal value in the groups. However, there are still several types of bias that can affect the trial.⁵³ The ITT approach was used to evaluate every patient record, regardless of whether the treatment was successful or not. Using the ITT approach instead of per-protocol will avoid overoptimistic estimates of efficacy of an intervention resulting from the removal of unsuccessfully treated subjects. Dropouts from the study, regardless if they moved to another location, if they did not finish the treatment because of poor compliance or if the treatment goal was not reached in a specific time, were all registered as non-successful. Thus, in this thesis, all cases, whether successful or not, as well as dropouts, were included in the final analysis. If the crossbite

was not corrected within the trial period of one year, the result was denoted as unsuccessful. In the cost-minimisation analysis we also calculated the costs for retreatments of unsuccessful cases.

The Hawthorne effect

One aspect to bear in mind is that when patients or therapists are involved in a trial, they tend to behave better than they would normally. This could be due to extra interest or knowledge or that they simply know that they are under supervision. This could, in particular, have an effect on the cooperation of the expansion plate patients, since cooperation is a key to successful treatments using removable appliances, resulting in a false-positive bias.⁵⁴ Therefore, the difference between quad-helix and expansion plate could be even greater, in favour of the quad-helix. However, the Hawthorne effect seems to be primarily visible during the first 6 months and there were no treatment periods shorter than 6 months in Paper III or Paper IV.

Quality of life

The child perception questionnaire (CPQ) protocol was used before, during and after treatment, to record discomfort, pain and other difficulties associated with both appliances, to gain a valuable insight on how the orthodontic treatments affects the OHRQoL.

The new CPQ data revealed that the CPQ level was fairly low from start (T0), in all four treated crossbite groups. After 6 weeks in treatment (T1) the values had increased, showing that the patients OHRQoL were reduced. After treatment (T2) the values had returned to the same as before.

Regarding CPQ, the scores there seem to be small differences regardless of which appliance was used or at which treatment instance the treatment was performed. However, somewhat higher CPQ scores were found 6 weeks of treatment (T1) using the quad-helix compared to expansion plates. One reason for this could be the increased number of emergency visits in the quad-helix general dentistry group. Although a high number of emergency visits was also seen in the expansion plate groups, this was mostly due to lost or broken plates, which might not have such a negative impact on

the patients compared to loose bands, or trauma from helices to either the tongue or palatal soft tissue or partly embedded in the palate. Wiedel and Bondemark⁵⁵ have also shown that there were small differences in patients' perceptions on pain and discomfort between removable plates and fixed appliances in anterior crossbite correction. Patients treated with a fixed appliance had a higher level of pain and discomfort the first 3 days and reported more difficulties eating different kinds of hard foods compared to the removable appliance group. On the other hand, the patients treated with removable appliances were more affected with speech problems and adverse effects on school and leisure activities.

The results from this new analysis show the impact different orthodontic appliances, quad-helix and expansion plate, have on young patients' OHRQoL, treated to correct unilateral posterior crossbite with functional shift. In addition, comparisons were made between treatments at specialist orthodontic clinics and general dentistry. Comparisons were also made with patients with untreated crossbite (UC) and to a normal control group (NC) i.e. patients with no or mild orthodontic treatment need. The main findings were that during treatment the CPQ values increased and that after treatment the CPQ values were reduced to the same level as before or even slightly lower. However, patients treated with quad-helix in general dentistry reported six weeks in treatment significantly higher CPQ values compared to the other treated groups. The negative impact was evident in the domain functional limitation; in this domain questions regarding eating, speaking and sleeping are investigated. One explanation for this could be the fact that the patients have the option to remove the appliance when speaking and eating compared to the fixed quad-helix. The difference between quad-helix in specialist orthodontic clinics and the quad-helix in general dentistry was perhaps not an unexpected finding since there were a larger number of emergency visits when treating patients with quad-helix at the general dentistry, and loose ligatures and broken bands could be a causation factor compared to fewer emergency visits when treated at the specialist orthodontic clinics. Irrespective of this, the treatment seems to have little impact on the OHRQoL since they were more or less unaware of the malocclusion from the start. Thus, it could not

really be expected to gain any major changes from the treatment, at the same time the patients do not seem to be particularly bothered by the treatment.

Cost-minimisation analysis

This study is the first to assess the costs of crossbite treatments performed in either specialist or general dentistry. It was clearly found that quad-helix treatment performed in specialist orthodontic clinics had lower costs than quad-helix or expansion plate treatments accomplished in general dentistry as well as expansion plate treatments in specialist dentistry. Consequently, the results rejected the hypotheses that the crossbite treatment methods are equal regarding costs and treatment efficiency in general and specialist dentistry.

One reason for the lower costs of quad-helix treatment was that this treatment is less dependent on patient cooperation compared to expansion plate treatment. The lower costs of treatments in specialist orthodontic clinics can be explained by trained specialists in orthodontics being employed in specialist clinics and that these orthodontists have greater experience working with fixed appliances compared with general dentists who may be more familiar with using removable appliances. Furthermore, there is some variety among the general dentists in clinical orthodontic skills. One could always argue that it could be better to intensify the orthodontic training for the dentists in general dentistry in how to manage the quad-helix treatment. However, as the situation currently stands in Sweden, there is a shortage of dentists and the turnover of general practitioners is high in many parts of the country, which results in an inconsistency of the treatments. Thus, it is often recognised that experienced dentists move on to other clinics and to other duties while the recruitment of new dentists is often restricted to less experienced ones. This constant flow of new co-workers at the clinics may affect the treatments in a negative way.

How much money could be saved?

The mean cost differences in the present study were clearly significant, and extrapolation of the data to a national level further highlights the differences between the economics of the two treatment modalities. As

of 2018, there were 123,866 9-year-old children living in Sweden and the prevalence of posterior crossbite is estimated to be around 10%. Treating all these children for posterior crossbite, the differences in societal costs between quad-helix treatment in specialist clinics and expansion plate treatments performed in general dentistry would be approximately € 3.8 million.

An RCT from 2013 on cost-effectiveness by Petrén et al.²⁵ showed that differences in societal costs, between treatments performed in general dentistry with the quad-helix or expansion plate, would save 3.5 million Euros if treatments were performed with the quad-helix instead of the expansion plate.

Looking at healthcare planning and allocation of resources there are now two independent studies^{25,39} that show that the quad-helix should be used instead of the expansion plate to reduce the social costs for crossbite corrections in mixed dentition.

Consequences for the general dentist

Will the challenge for general dentistry be less stimulating if the number of duties is reduced, if all that is left is to drill and fill? Another aspect may be, is it possible for the specialist dentistry to treat all crossbites? Currently, there is a shortage of orthodontists in many parts of Sweden and an increased workload would cause the queues to be further extended. Also, a none-too-bold guess is that treating all crossbite patients at the specialist clinics might contribute to a negative impact on environmental emissions, since there are fewer specialist clinics and this will result in more travelling for the patients. At the same time shorter treatment time and fewer dental appointments were seen in patients treated at specialist clinics, so ultimately this might be a zero-sum game. However, if the crossbite corrections are to be performed in general dentistry, the dentists must be trained at a specialist clinic to improve their skills and, if the circumstances do not allow this, then the treatment should be performed by an orthodontist at the specialist clinic.

Three-dimensional analysis

Unilateral posterior crossbites can be corrected in many ways, but do any of them change the morphology of the palate? It can be pointed out that Paper IV is the first RCT that has used and analysed the three-dimensional changes after unilateral posterior crossbite correction in mixed dentition and simultaneously compared the treatment results with those changes that occurred in two control groups; one untreated crossbite control group and another control group with normal occlusion. The use of an untreated crossbite group was to analyse how the growth developed compared both to the treated crossbite patients and to the subjects with normal occlusion. The reason for using a normal control group was to also evaluate the maxillary dimensions in subjects with normal occlusion, i.e. if the treated crossbite patients reached the same dimension values as the normal children. The normal material was collected at the same time by the general practitioners or at the specialist clinics when siblings or friends of patients participating in the crossbite study attended the appointment.

Both the 2D and 3D evaluations in this study revealed that the maxillary variables of the crossbite children had smaller values compared to children with normal occlusion at baseline. After treatment with the quad-helix or expansion plate, the maxillary expansion of the crossbite patients significantly improved, and both the 2D and 3D evaluations showed that the maxillary parameters reached the size of the parameters found in the normal group with normal occlusion. It can now be concluded that the morphology of the palate can be normalised with an expansion treatment. These post-treatment results contrasted with the treatment outcomes presented by Petrén and Bondemark.¹⁶ The reason for these divergent results is unknown and future research on this topic may be justified.

Environmental considerations

Plaster casts have been used for many decades and have been proven to work effectively. However, they have several disadvantages which have consequences for the environment, such as plaster- and impression material production, transportation of impressions and casts from the orthodontic office to a laboratory technician and then

back again, considerable space requirements for storage and the time needed to retrieve specific casts and also the risk of losing casts.

Three-dimensional scanners are increasingly incorporated into orthodontic offices. With an intraoral scanner it is possible to directly take an intraoral digital impression. The three-dimensional digital models are then used for several purposes such as orthodontic measurements, and treatment planning including prosthetic and surgery planning. Intraoral scanning, computed tomography and extraoral scanning can all be used together for a complete digital planning. These files can then be sent digitally to different clinics and therapists.

New data

To reveal if there were any specific treatment differences between the appliances the four groups were reduced to two. When the results from the two quad-helix groups (QHS, QHG) and from the two expansion plate groups (EPS, EPG) were merged into one QH group and into one EP group, some new significant differences and findings could be revealed. Surface area, projection area and shell volume all increased significantly more in the treated groups compared to the untreated groups. Projection area increased significantly more in the quad-helix group compared to the expansion plate group, whereas the effect on the surface area and shell volume were more similar in the two treated groups (Table 9)

The effect on shell volume was significantly larger in the treated groups but not to the same extent as surface- and projection area. One reason for this could be that, as Haas reported, as a result of expanding the maxilla, the palatal vault of the maxilla was lowered as a result of outward tilting of the maxillary halves.⁵⁶

Ethical considerations

Over the last few decades there has been a shift towards research with children and the adoption of the concept of child-centred research. However, the majority of oral health research is conducted on children, rather than with them. The position of children in society has changed, with increasing recognition of children's rights and the need to involve them in decisions about their education, and social and healthcare including oral health. In orthodontics, it is the child who undergoes the treatment and endures the appliance and the

consequences of whether or not they choose treatment. It is therefore important to consider their perspectives, desires, and expectations in decision making about their orthodontic care. With this in mind it is important to conduct research with the children at the centre, with due respect for ethical principles. In Sweden, ethical regulations and international recommendations apply to research upon children, in order to protect the child. Informed consent must be obtained from the parents/legal guardian if the child is younger than 15 years. However, it is the child who has the final decision and is excluded from the research trial if the child disagrees.

Generalisability

This thesis was based on a sample consisting of children in mixed dentition, with a mean age of 8-10 years of age and with an even gender distribution. When the multicentre RCT was undertaken, attempts were made to reduce the risk of bias and confounding factors. The prospective design and protocol ensured that the baseline characteristics and treatment progress were easily observed. Furthermore, a treatment protocol as close as possible to the normal treatment procedure was used to ensure that the results could be implemented in everyday work. Moreover, the multicentre design resulted in treatment outcomes from a large number of operators, and this made the patient-related outcome closer to reality compared with if only a few operators were involved in the study.

Overall limitations of this thesis

It is important to stress that it is sometimes difficult to compare the results of quality of life studies and also of cost-effectiveness studies. A potential demographic impact is displayed when comparing children in this study with children from other countries, for example Brazil, where the Swedish children scored lower on the CPQ. Caries have a negative impact on quality of life, and the Swedish patients had a very low frequency, and, thereby, this makes it difficult to compare to the Brazilian patients. Another factor that may affect the result is that children and adolescents from wealthy families have a higher oral health related quality of life.⁵⁷

In a cost-effectiveness study it is also important to point out that monetary variables are influenced by local factors such as insurance systems, staff salaries, rental costs, taxes, urban versus rural areas, etc., and, consequently, the figures shown in this thesis could not be directly extrapolated to other locations but should be seen in relation to the conditions of different societies and countries.

Future research

The quality of life results from Paper II where comparisons were made between large overjet, unilateral posterior crossbite and normal controls before treatment could be followed up after treatment to compare if there are any differences in the long term.

Another new area of research could be to compare TAD-supported expanders to conventional expanders like the one used in Paper III and IV. Finally, it may also be worth exploring how the expansion treatments will affect sleep-disordered breathing.

CONCLUSIONS

The systematic review (Paper I) led to the following conclusions:

- Few orthodontic studies have presented both economic and clinical outcomes.
- There is currently insufficient evidence available about the health economics of orthodontic interventions. Further investigation is warranted.

In the updated supplementary literature search the following conclusions were made:

- In the last few years there has been an increase in the number of articles, especially from Sweden, presenting both economic and clinical outcomes.
- There is now high evidence for using the quad-helix to effectively treat unilateral posterior crossbite in mixed dentition.

In the controlled trial (Paper II) comparing oral health related quality of life (OHRQoL) between excessive overjet (EO), unilateral posterior crossbite with functional shift (UPC) and normal occlusion (NO) it was concluded that:

- Young children with EO reported significantly lower self-perceived OHRQoL compared to children with UPC with functional shift or children with NO and with no or mild orthodontic treatment need.
- No difference regarding self-perceived OHRQoL could be detected between children with UPC with functional shift when compared with children with NO and with no or mild orthodontic treatment need.

- Despite the presence of malocclusions, the children generally reported low CPQ scores which implies an overall fairly good self-perceived OHRQoL.

The new CPQ results from before, during and after treatment showed:

- The CPQ values increased during treatment but returned to the same level after treatment as before.
- There seems to be small differences in CPQ values regardless of which appliance are used or at which treatment instance the treatment is performed.

In the RCT on clinical effectiveness and cost-analysis (Paper III) it was concluded:

- Quad-helix treatment had a higher success rate compared to expansion plate.
- It was more cost effective to perform correction of unilateral crossbite in mixed dentition if the correction was made with quad-helix.

Treatment of unilateral posterior crossbite in mixed dentition is recommended to be performed by a specialist in orthodontics using the quad-helix appliance.

In the RCT on three-dimensional evaluation of crossbite correction (Paper IV) it was concluded that:

- Children with unilateral posterior crossbite with functional shift in mixed dentition revealed before treatment a significantly smaller palatal surface and projection area as well as smaller palatal shell volume than normal control children.
- Unilateral posterior crossbite correction resulted in a normalisation of the palatal structures in terms of palatal area and volume.
- Quad-helix treatments in orthodontic specialist clinics but also in general dentistry as well as removable expansion plate treatments in specialist orthodontic clinics revealed a significant increase of palatal area and volume.
- The natural growth changes during the one-year trial were small.

The new results from when the two quad-helix groups (QHS, QHG) and the two expansion plate groups (EPS, EPG) were merged into one QH group and into one EP group:

- Projection area increased significantly more in the quad-helix group compared to the expansion plate group,
- The effect on the surface area and shell volume were more similar in the two treated groups.

Clinical implications

Quad-helix treatment is superior to expansion plate treatment in terms of success rate and cost-effectiveness, when crossbite corrections are planned in mixed dentition.

Treatment of unilateral posterior crossbite in mixed dentition should be performed by a specialist in orthodontics using the quad-helix appliance.

Unilateral posterior crossbite correction resulted in a normalisation of the occlusion as well as the palatal structures in terms of palatal area and volume.

Despite the presence of malocclusions, the children generally reported low CPQ scores which implies an overall fairly good self-perceived oral health related quality of life.

ACKNOWLEDGEMENTS

This project has been an amazing experience for me and made me realise that it is possible to do more than just moving teeth. However, the completion of this thesis would not have been achieved without the support of many people, to whom I wish to express my gratitude, especially to all the children who participated in the studies.

I also wish to acknowledge the following people for their invaluable contributions:

Professor *Lars Bondemark*, my main supervisor and co-author, for sharing with me your great knowledge in the field of research, for your steadfast belief in this project and never-ending positive attitude. My co-supervisor and co-author Odont. dr. and Associate Professor *Sofia Petrén*, for excellent advice in health economy and methodological discussion in a very structured way. Yet also with a very positive attitude and always with a smile on your lips. Thank you to Lars and Sofia for all the help with this process of me becoming a researcher. What a team you are!

Odont. dr. Associate Professor *Liselotte Paulsson* my co-author on Paper I and II for stimulating discussion and constructive criticism of the manuscripts.

Co-author PhD *Anders Norlund* on Paper I for your great knowledge in the health economy.

Senior consultant orthodontist *Jenny Kallunki* my co-author on Paper II and Odont dr. *Lillemor Dimberg* co-author on Paper II for your expert knowledge in the field of Oral Health Related Quality of Life.

Dr. dent. med. *Aljaž Golež*, Associate Prof. dr. *Jasmina Primožič* and Prof. dr. *Maja Ovsenik* my co-authors on Paper IV, for excellent support with the 3D analysis.

Per Erik Isberg, for excellent help with statistical analysis on Paper II-IV.

Anders Nielsen for excellent revision of the English text

My friend and colleague senior consultant orthodontist *Seifi Esmaili* who, without personal gain, helped me treat patients and collect the material for the studies. Your valuable support and inspiring interest are greatly appreciated.

Staff members at the public dental clinics all around Varberg, Falkenberg, Hylte and Halmstad, Halland County Council, for their valuable help with the clinical procedures during patient treatments.

My colleagues, co-workers and friends at the Department of Orthodontics in Halmstad and the Maxillofacial unit in Halmstad.

My friend and colleague Dr. *Lennart Lagerström*, who encouraged me and guided me in the right directions at the start of my research career.

My dearest *Jenny, Casper and Selma*, for your patience and understanding during these years.

My parents *Lisbeth* and *Thomas*, and my sister *Åsa*, thank you for your constant support and encouragement, and for being there for my family and me.

I would like to dedicate this to my dear friend and colleague senior consultant orthodontist *Lovisa Bjerke* who sadly passed away earlier this year.

The research was supported by grants from the FoU County Council, Halland Sweden, Södra sjukvårdsregionen, Faculty of Odontology, Malmö University, Sweden, the Swedish Dental Society, and the European Orthodontic Society.

REFERENCES

1. Malandris, M. and Mahoney, EK. (2004) Aetiology, diagnosis and treatment of posterior cross-bites in the primary dentition. *International Journal of Paediatric Dentistry*, 14, 155-166.
2. Primožič, J., Ovsenik, M., Richmond S., Kau, C.H. and Zhurov A. (2009) Early crossbite correction: a three-dimensional evaluation. *European Journal of Orthodontics*, 31, 352–356.
3. Thilander, B. and Myrberg, N. (1973) The prevalence of malocclusion in Swedish schoolchildren. *Scandinavian Journal of Dental Research*, 81, 12–21.
4. Heikinheimo, K. (1978) Need of orthodontic treatment in 7-year-old Finnish children. *Community Dentistry and Oral Epidemiology*, 6, 129–134
5. Bassler-Zeltmann, S., Kretschmer, I. and Goz, G. (1998) Malocclusion and the need for orthodontic treatment in 9-year-old children. Survey based on the Swedish National Board of Health and Welfare Scale. *Journal of Orofacial Orthopedics*, 59, 193-201
6. Lindner A. (1989) Longitudinal study on the effect of early interceptive treatment in 4-year-old children with unilateral cross-bite. *Scandinavian Journal of Dental Research*, 97, 432-438.
7. Pinto, A.S., Buschang, P.H., Throckmorton, G.S. and Chen, P. (2001) Morphological and positional asymmetries of young children with functional unilateral posterior crossbite. *American Journal of Orthodontics and Dentofacial Orthopedics*, 120, 513-520.
8. Egermark, I., Magnusson, T. and Carlsson, G.E. (2003) A 20-year follow-up of signs and symptoms of temporomandibular disorders and malocclusions in subjects with and without orthodontic treatment in childhood. *The Angle Orthodontist*, 73, 109–115.
9. Andrade, A.S., Gameiro G.H., Derossi, M. and Gaviao, M.B. (2009) Posterior crossbite and functional changes. A systematic review. *The Angle Orthodontist*, 79, 380-386.

10. Leonardi, R., Muraglie, S., Bennici, O., Cavallini, C. and Spampinato C. (2019) Three-dimensional analysis of mandibular function units in adult patients with unilateral posterior crossbite: A cone beam study with the use of mirroring and surface-to-surface matching techniques. *The Angle Orthodontist*, 89, 590-596
11. Tsanidis, N., Antonarakis, GS. and Kiliaridis S. (2016) Functional changes after early treatment of unilateral posterior cross-bite associated with mandibular shift: a systematic review. *Journal of Oral Rehabilitation* 43, 59-68.
12. Iodice, G., Danzi, G., Cimino, R., Paduano, S. and Michelotti, A. (2016) Association between posterior crossbite, skeletal, and muscle asymmetry: a systematic review. *European Journal of Orthodontics*, 38, 638–651
13. Thilander B. and Bjerklín K. (2012). Posterior crossbite and temporomandibular disorders (TMDs): need for orthodontic treatment? *European Journal of Orthodontics*, 4, 667-673.
14. Gesch, D., Bernhardt, O., Kocher, T., John, U., Hensel, E. and Alte D. (2004). Association of malocclusion and functional occlusion with signs of temporomandibular disorders in adults: results of the population-based study of health in Pomerania. *The Angle Orthodontist* 74, 512-520.
15. Primoži , J., Perinetti, G., Richmond, S. and Ovsenik M. (2013). Three-dimensional evaluation of facial asymmetry in association with unilateral functional crossbite in the primary, early, and late mixed dentition phases. *The Angle Orthodontist* 83, 253-258.
16. Petrén, S. and Bondemark, L. (2008) Correction of unilateral posterior crossbite in the mixed dentition: a randomized controlled trial. *American Journal of Orthodontics and Dentofacial Orthopedics*, 133:790.e7-13.
17. Timms DJ. The dawn of rapid maxillary expansion. (1999). *The Angle Orthodontist*. 69, 247–250.
18. Bell RA. (1982) A review of maxillary expansion in relation to rate of expansion and patient's age. *American Journal of Orthodontics and Dentofacial Orthopedics*, 81, 32–37
19. Gill, D., Naini, F., McNally, M. and Jones, A. (2004). The Management of Transverse Maxillary Deficiency. *Dental Update*, 31, 516–523
20. Agostino, P., Ugolini, A., Signori, A., Silvestrini-Biavati, A., Harrison, JE. and Riley, P. (2014) Orthodontic treatment for posterior crossbites. *Cochrane Database of Systematic Reviews* 8;(8):CD000979. doi: 10.1002/14651858.CD000979
21. SBU. Malocclusions and Orthodontic Treatment in a Health Perspective. A systematic review. The Swedish Council on Technology Assessment in Health Care. SBU-Report 2005, Nr 176. ISBN: 91-85413-06-2. (In Swedish)

22. Kumar, S., Williams, A.C. and Sandy, J.R. (2006). How do we evaluate the economics of health care? *European Journal of Orthodontics*, 28, 513–519.
23. Drummond, M.F., Scholpher, M.J., Torrence, G.W., O'Brian, B. and Stoddart, G.L. (2005) *Methods for the Economical Evaluation of Healthcare Programs*. Oxford University Press, Oxford, 3rd edn.
24. Tickle, M. (1997) Clinical effectiveness and primary dental care. 2. The influence of health economics. *Primary Dental Care*, 4, 85–87.
25. Petrén, S., Bjerklín, K., Marké, L.-Å. and Bondemark, L. (2011). Early Treatment of Posterior Crossbite – an RCT Study on Cost-Minimization. *European Journal of Orthodontics*, 35, 14-21.
26. Sollenius, O., Petrén, S., Björnsson, L., Norlund, A. and Bondemark, L. (2016) Health economic evaluations in orthodontics: a systematic review. *European Journal of Orthodontics*, 38, 259–265.
27. Dimberg, L., Arnrup, K. and Bondemark, L. (2015) The impact of malocclusion on the quality of life among children and adolescents: a systematic review of quantitative studies. *European Journal of Orthodontics*, 37, 238–247.
28. Harrison, J.E. and Ashby, D. (2001) Orthodontic treatment for posterior crossbites. *Cochrane Database of Systematic Reviews Issue 1*. Art. No.: CD000979. Published Online: 8 OCT 2008.
29. Martin, C.B., Chalmers, E.V., McIntyre, G.T., Cochrane, H. and Mossey P.A. (2015) Orthodontic scanners: what is available? *Journal of Orthodontics*, 42, 136–143.
30. Primožic, J., Perinetti, G., Contardo, L. and Ovsenik, M. (2013) Diagnostic performance of 3-dimensional evaluation of palatal vault changes in assessing successful treatment of constricted maxilla in growing subjects. *American Journal of Orthodontics and Dentofacial Orthopedics*, 143, 42–49
31. Dunnett, C.W. and Tamhane, A.C. (1992) A step-up multiple test procedure. *Journal of the American Statistical Association*, 87, 162–170.
32. Tynelius Edman, G., Lilja-Karlander, E. and Petrén, S. (2014) A cost-minimization analysis of an RCT of three retention methods. *The European Journal of Orthodontics*, 36, 436–441.
33. Laag, B. and Ström, C. (2009) Comparisons of similar patients treated by general dental clinicians and orthodontic specialists. Outcome and economical considerations. *Swedish Dental Journal*, 33, 67–73.
34. O'Brien, K., et al. (2009) Early treatment for Class II Division 1 malocclusion with the Twin-block appliance: a multi-centre, randomized, controlled trial. *American Journal of Orthodontics & Dentofacial Orthopedics*, 135, 573–579.

35. Hichens, L., Rowland, H., Williams, A., Hollinghurst, S., Ewings, P., Clark, S., Ireland, A. and Sandy, J. (2007) Cost-effectiveness and patient satisfaction: Hawley and vacuum-formed retainers. *The European Journal of Orthodontics*, 29, 372–378.
36. Konst, E.M., Prah, C., Weersink-Braks, H., De Boo, T., Prah-Andersen, B., Kuijpers-Jagtman, A.M. and Severens, J.L. (2004) Cost-effectiveness of infant orthopedic treatment regarding speech in patients with complete unilateral cleft lip and palate: a randomized three-center trial in the Netherlands (Dutchcleft). *Cleft Palate Craniofacial Journal*, 41, 71–77.
37. Severens, J.L., Prah, C., Kuijpers-Jagtman, A.M. and Prah-Andersen, B. (1998) Short-term cost-effectiveness analysis of presurgical orthopaedic treatment in children with complete unilateral cleft lip and palate. *Cleft Palate Craniofacial Journal*, 35, 222–226.
38. Hermanson, H., Kurol, J. and Rönnerman, A. (1985) Treatment of unilateral posterior crossbite with quad-helix and removable plates. A retrospective study. *The European Journal of Orthodontics*, 7, 97–102.
39. Sollenius, O., Petré, S. and Bondemark, L. (2019) An RCT on clinical effectiveness and cost analysis of correction of unilateral posterior crossbite with functional shift in specialist and general dentistry. *European Journal of Orthodontics*, Apr 11 pii, cjz014. doi:10.1093/ejo/cjz014. [Epub ahead of print].
40. Ganzer N., Feldmann, I., Petré, S. and Bondemark, L. (2019) A cost-effectiveness analysis of anchorage reinforcement with miniscrews and molar blocks in adolescents: a randomized controlled trial. *The European Journal of Orthodontics*, 29, 180-187.
41. Cirgic, E., Kjellberg, H., Petzold, M. and Hansen, K. (2018) A cost-minimization analysis of large overjet reduction with two removable functional appliances based on a randomized controlled trial. *The European Journal of Orthodontics*, 27, 437-443.
42. Wiedel, A.P., Norlund, A., Petré, S. and Bondemark, L. (2016) A cost-minimization analysis of early correction of anterior crossbite - a randomized controlled trial. *European Journal of Orthodontics*, 38,140-145.
43. www.jla.nihr.ac.uk The James Lind Alliance.
44. Goodman, C.S. (2014) HTA 101), HTA 101. Introduction to health technology assessment. http://www.nlm.nih.gov/nichsr/hta101/HTA_101_FINAL_7-23-14.pdf.
45. Guyatt, G.H., et al. (2011) GRADE guidelines: 4. Rating the quality of Evidence-study limitations (risk of bias). *Journal of Clinical Epidemiology*, 64, 407–415.
46. Mandall, N.A., McCord, J.F., Blinkhorn, A.S., Worthington, H.V. and O'Brien K. (2000) Perceived aesthetic impact of malocclusion and oral self-perceptions in 14-15-year-old Asian and Caucasian children in greater Manchester. *European Journal of Orthodontics*, 22, 175-183.

47. Espeland, LV. and Stenvik, A. (1991) Perception of personal dental appearance in young adults: relationship between occlusion, awareness, and satisfaction. *American Journal of Orthodontics and Dentofacial Orthopedics*, 100, 234-241.
48. Wheeler, T., McGorray, S., Yurjiewicz, L., Keeling, S. and King, G. (1994) Orthodontic treatment demand and need in third and fourth grade schoolchildren. *American Journal of Dentofacial Orthopedics*, 106, 22-33.
49. Christopherson, E.A., Briskie, D. and Inglehart, M.R. (2009) Objective, subjective, and self-assessment of preadolescent orthodontic treatment need - a function of age, gender, and ethnic/racial background? *Journal of Public Health Dentistry*, 69, 9-17.
50. O'Brien, K., McComb, J.L., Fox, N. and Wright, J. (1996) Factors influencing the uptake of orthodontic treatment. *British Journal of Orthodontics*, 23, 331-334.
51. Berset, G.P., Eilertsen, I.M., Lagerström, L., Espeland, L. and Stenvik, A. (2000). Outcome of a scheme for specialist orthodontic care. *Swedish Dental Journal*, 24, 39-48.
52. Lagerström L., Stenvik A., Espeland L. and Hallgren A. (2000). Outcome of a scheme for orthodontic care: a comparison of untreated and treated 19-year-olds. *Swedish Dental Journal*, 24, 49-57.
53. Bondemark L. and Abdulraheem S. (2018). Intention to treat (ITT) analysis as reported in orthodontic randomized controlled trials-evaluations of methodology and recommendations for the accurate use of ITT analysis and handling dropouts. *European Journal of Orthodontics*, 40, 409-413.
54. Abdulraheem S. and Bondemark L. (2018). Hawthorne effect reporting in orthodontic randomized controlled trials: truth or myth? Blessing or curse? *European Journal of Orthodontics*. 40, 475-479.
55. Wiedel A.P. and Bondemark L. (2016) A randomized controlled trial of self-perceived pain, discomfort and impairment of jaw function in children undergoing orthodontic treatment with fixed or removable appliance. *The Angle Orthodontist*, 86, 324-330.
56. Haas AJ. (1961) Rapid expansion of the maxillary dental arch and nasal cavity by opening the midpalatal suture. *The Angle Orthodontist*. 31, 73-90.
57. Kavaliuskiene, A., Šidlauskas, A. and Zaborskis, A. (2018) Relationship Between Orthodontic Treatment Need and Oral Health-Related Quality of Life among 11-18-Year-Old Adolescents in Lithuania. *International Journal of Environmental Research and Public Health*, Open access 15, 1012.

CPQ₈₋₁₀

BARNVERSION

Dagens datum: ____/____/____
dag månad år

1. Är du pojke eller flicka?

Pojke

Flicka

2. När är du född? ____/____/____ Ålder ____
dag månad år

3. När du tänker på dina tänder eller din mun, skulle du säga att de är:

Mycket bra

Bra

Okej

Dåliga

4. Hur mycket stör dina tänder eller din mun dig varje dag?

Inte alls

Lite grann

Något

Mycket

Nu några frågor om dina tänder och din mun

5. Hur ofta har du under de senaste 4 veckorna haft **ont i tänderna eller munnen?**

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

6. Hur ofta har du under de senaste 4 veckorna haft **sår i munnen?**

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

7. Hur ofta har du under de senaste 4 veckorna haft **ont i tänderna när du dricker kalla drycker eller äter varm mat?**

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

8. Hur ofta har du under den senaste 4 veckorna haft **mat som fastnat mellan tänderna?**

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

9. Hur ofta har du under de senaste 4 veckorna haft **dålig andedräkt**?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

10. Hur ofta har du under de senaste 4 veckorna på grund av dina tänder eller din mun **behövt mer tid än andra för att äta din mat**?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

11. Hur ofta har du under de senaste 4 veckorna på grund av dina tänder eller din mun **haft svårt att bita eller tugga äpplen, morötter eller kött**?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

12. Hur ofta har du under de senaste 4 veckorna på grund av dina tänder eller din mun **haft svårt att äta vad du vill**?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

- 13.** Hur ofta har du under de senaste 4 veckorna **haft svårt att säga några ord** på grund av dina tänder eller din mun?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

- 14.** Hur ofta har du under de senaste 4 veckorna **haft problem med att sova på natten** på grund av dina tänder eller din mun?

Aldrig

En eller två gånger

Ibland

Alltid

Varje dag, eller nästan varje dag

Nu några frågor om dina känslor

15. Hur ofta har du under de senaste fyra 4 veckorna **varit upprörd eller irriterad** på grund av dina tänder eller din mun?
- Aldrig
- En eller två gånger
- Ibland
- Alltid
- Varje dag, eller nästan varje dag
16. Hur ofta har du under de senaste fyra 4 veckorna **känt dig besviken eller ledsen** på grund av dina tänder eller din mun?
- Aldrig
- En eller två gånger
- Ibland
- Ofta
- Varje dag, eller nästan varje dag
17. Hur ofta har du under de senaste 4 veckorna **varit blyg** på grund av dina tänder eller din mun?
- Aldrig
- En eller två gånger
- Ibland
- Ofta
- Varje dag, eller nästan varje dag

18. Hur ofta har du under de senaste 4 veckorna **varit oroad** över vad andra tycker om dina tänder eller din mun?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

19. Hur ofta har du under de senaste 4 veckorna varit oroad över om du på grund av dina tänder eller din mun **inte ser lika bra ut som andras?**

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

Frågor om din skola

20. Hur ofta har du under de senaste 4 veckorna **varit borta från skolan** på grund av dina tänder eller din mun?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

21. Hur ofta har du under de senaste 4 veckorna **haft svårt att göra läxor** på grund av dina tänder eller din mun?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

22. Hur ofta har du under de senaste 4 veckorna **haft svårt att följa med i skolan** på grund av dina tänder eller din mun?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

23. Hur ofta har du under de senaste 4 veckorna **inte velat tala eller läsa högt i klassen** på grund av dina tänder eller din mun?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

Frågor om dina relationer med andra personer

24. Hur ofta har du under de senaste 4 veckorna, på grund av dina tänder eller din mun, **försökt låta bli att skratta eller le när du varit tillsammans med andra barn?**

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

25. Hur ofta har du under de senaste 4 veckorna **inte haft lust att prata med andra barn** på grund av dina tänder eller din mun?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

26. Hur ofta har du under de senaste 4 veckorna **inte haft lust att vara tillsammans med andra barn** på grund av dina tänder eller din mun?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

27. Hur ofta har du under de senaste 4 veckorna **undvikit sport eller andra aktiviteter** på grund av dina tänder eller din mun?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

28. Hur ofta under de senaste 4 veckorna har **andra barn retat dig eller kallat dig öknamn** på grund av dina tänder eller din mun?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

29. Hur ofta under de senaste 4 veckorna har **andra barn frågat dig** om dina tänder eller din mun?

Aldrig

En eller två gånger

Ibland

Ofta

Varje dag, eller nästan varje dag

Allra sist; två extrafrågor:

Gör det ont i tinning, ansikte käke eller käkled en gång i veckan eller oftare?

Ja

Nej

Gör det ont när du gapar eller tuggar, en gång i veckan eller oftare?

Ja

Nej

Stort tack för din medverkan!

I

Systematic Review

Health economic evaluations in orthodontics: a systematic review

Ola Sollenius*, Sofia Petrén**, Liselotte Björnsson**,
Anders Norlund*** and Lars Bondemark**

*Department of Orthodontics, County Council, Halland, Halmstad, **Department of Orthodontics, Faculty of Odontology, Malmö University, ***Section of Insurance Medicine, Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden.

Correspondence to: Ola Sollenius, Department of Orthodontics, County Council, Halland, SE-302 46 Halmstad, Sweden. E-mail: ola.sollenius@regionhalland.se

Summary

Background: Economic evaluation is assuming increasing importance as an integral component of health services research.

Aim: To conduct a systematic review of the literature and assess the evidence from studies presenting orthodontic treatment outcomes and the related costs.

Materials/methods: The literature review was conducted in four steps, according to Goodman's model, in order to identify all studies evaluating economic aspects of orthodontic interventions. The search covered the databases Medline, Cinahl, Cochrane, Embase, Google Scholar, National Health Service Economic Evaluation Database, and SCOPUS, for the period from 1966 to September 2014. The inclusion criteria were as follows: randomized controlled trials or controlled clinical trials comparing at least two different orthodontic interventions, evaluation of both economic and orthodontic outcomes, and study populations of all ages. The quality of each included study was assessed as limited, moderate, or high. The overall evidence was assessed according to the GRADE system (The Grading of Recommendations Assessment, Development and Evaluation).

Results: The applied terms for searches yielded 1838 studies, of which 989 were excluded as duplicates. Application of the inclusion and exclusion criteria identified 26 eligible studies for which the full-text versions were retrieved and scrutinized. At the final analysis, eight studies remained. Three studies were based on cost-effectiveness analyses and the other five on cost-minimization analysis. Two of the cost-minimization studies included a societal perspective, i.e. the sum of direct and indirect costs. The aims of most of the studies varied widely and of studies comparing equivalent treatment methods, few were of sufficiently high study quality. Thus, the literature to date provides an inadequate evidence base for economic aspects of orthodontic treatment.

Conclusion: This systematic review disclosed that few orthodontic studies have presented both economic and clinical outcomes. There is currently insufficient evidence available about the health economics of orthodontic interventions. Further investigation is warranted.

Introduction

Economic evaluation has become an integral component of health services research in recent years and is likely to assume increasing importance for decision making in dental services in the

future (1). The main reason is that resources within the health sector (personnel, time, facilities, equipment, and knowledge) are limited (2). Thus, failure to analyse economic aspects of dental health services may result in unsustainable over expenditure or

withdrawal or reduction of services or resources in other areas of health care (3).

Economic evaluations involve techniques that were developed in order to support decision making. In health care settings, cost evaluations provide only limited information for this process. In allocating resources, including dental care and orthodontics, health services purchasers need to take into account not only evidence of clinical effectiveness of treatment procedures but also relative costs, i.e. 'value for money' (4). Two features characterize health economic evaluation: it concerns the relationship between costs (input) and consequences (output) and preferential selection of diagnostic or therapeutic options, implying alternative ways of allocating resources.

Four main analyses are available for economic evaluations (2):

A cost-effectiveness analysis is characterized by analysis of both costs and outcomes, where the magnitude of the outcomes of the various methods might differ. These studies express effectiveness in a single dimension in order to enable direct comparison of costs.

A cost-minimization analysis, which is a type of cost-effectiveness analysis, applicable when two or more treatment methods achieve the same outcomes (e.g. correction of crossbite) and the aim is to identify the least expensive method.

A cost-utility analysis, focusing on both costs and quality of the health outcome produced by the treatments: used, for example, in health-related quality-of-life studies.

A cost-benefit analysis, which is characterized by the fact that health outcome is also expressed in monetary units. This kind of analysis is seldom used but has been implemented, for example, when evaluating vaccination programs.

Economic evaluations with a societal perspective include calculations of direct and indirect costs. Direct costs are those directly associated with diagnostic or treatment interventions, i.e. clinical interventions, costs of material, costs for the premises and equipment, maintenance, cleaning, and staff costs. Indirect costs arise as consequences of absence from work for treatment, often defined as corresponding to loss of income incurred by the patient, or the patient's parents, in taking time off from work for clinical appointments and travel to and from the clinic. The societal costs are defined as the sum of direct and indirect costs.

Delivery of equitable health care requires assessment of the economic implications of different interventions (1). In orthodontics, economic analysis is also important as a basis for decision making in planning and management of dental care.

The aim of this study was to conduct a systematic review of published studies of health economics in orthodontics, assessing the evidence from studies presenting an analysis of costs associated with the treatment outcomes.

Materials and methods

The literature review was conducted in four steps, according to Goodman's model (5).

Definition of the research question

The question to be addressed was 'What evidence is available from studies combining costs and orthodontic treatment outcomes?'

Formulation of a plan for the literature search

A literature search was conducted to identify all studies evaluating economic aspects of orthodontic interventions. The period to be

covered was from 1966 to September 2014. The following databases were searched: Medline, Cinahl, Cochrane, Embase, Google Scholar, National Health Service Economic Evaluation Database, and SCOPUS.

The computerized search was conducted with the assistance of a specialist in informatics at the Library, Faculty of Odontology, Malmö University, Sweden. The following search syntax was used:

```
'economics, dental'[MeSH Terms] OR ('economics'[All Fields]
AND 'dental'[All Fields]) OR 'dental economics'[All Fields] OR
('economics'[All Fields] AND 'dental'[All Fields]) OR 'economics,
dental'[All Fields]) AND ('orthodontics'[MeSH Terms] OR
'orthodontics'[All Fields]) OR (('cost-benefit analysis'[MeSH
Terms] OR ('cost-benefit'[All Fields] AND 'analysis'[All Fields])
OR 'cost-benefit analysis'[All Fields] OR ('cost'[All Fields] AND
'effectiveness'[All Fields]) OR 'cost effectiveness'[All Fields]) AND
('orthodontics'[MeSH Terms] OR 'orthodontics'[All Fields]) OR
(cost-utility[All Fields] AND ('orthodontics'[MeSH Terms] OR
'orthodontics'[All Fields]) OR (cost-minimization[All Fields]
AND ('orthodontics'[MeSH Terms] OR 'orthodontics'[All
Fields])) OR (('cost-benefit analysis'[MeSH Terms] OR ('cost-
benefit'[All Fields] AND 'analysis'[All Fields]) OR 'cost-benefit
analysis'[All Fields] OR ('cost'[All Fields] AND 'benefit'[All
Fields]) OR 'cost benefit'[All Fields]) AND ('orthodontics'[MeSH
Terms] OR 'orthodontics'[All Fields])).
```

Literature search and retrieval of publications

At the start of the review, consensus was reached on the criteria presented in Table 1.

The potential eligibility of studies was determined independently by five researchers. The titles and abstracts of all potentially relevant studies were independently reviewed. The full-text version of an article was retrieved if at least one of the five reviewers considered the study to be relevant or if the title and abstract did not provide sufficient information. Using a preset protocol and the initial inclusion and exclusion criteria, the full-text versions of the articles were independently analysed and evaluated by the five researchers. In cases of inter-examiner disagreement, each article was reread and discussed until consensus was reached. The reference lists of articles deemed eligible were also hand-searched for additional studies.

Data extraction, quality assessment, interpretation, and evaluation of evidence

Data from studies were tabulated as follows: Reference, country, aim of the study, study design, study population, economic and orthodontic outcomes, and finally, study quality (Table 2).

The quality assessment was performed according to protocols independently by the researchers and disagreement over assessment was resolved by discussion to achieve consensus. Study quality was denoted as limited, moderate, or high, according to the checklist proposed by Drummond *et al.* (2). The strength of the scientific evidence based on study quality from included studies was undertaken according to the GRADE system (14).

Results

The PRISMA-compliant selection process (15) is presented in Figure 1. Initially, 1838 articles were identified, of which 989 were deleted as duplicates. After application of the inclusion and exclusion criteria, the full-text versions of 26 studies remained (Figure 1).

Table 1. Inclusion and exclusion criteria used in this systematic review.

Inclusion criteria	Exclusion criteria
Studies comparing at least two different orthodontic interventions, including evaluation of economic and orthodontic outcomes.	Incomplete economic analysis
Randomized controlled trials (RCTs) or controlled clinical trials (CCTs)	Practice economic evaluations
Study populations of all ages	Case reports and case series
Full-text articles written in all languages	Descriptive articles, opinions, editorials and reviews

A further 18 of these studies were excluded for various reasons (Figure 1). Eight studies then remained for final assessment, four from Sweden (6–8,13) and two each from the Netherlands (11,12) and UK (9,10). Of these studies, there were four on different orthodontic appliances (7–9,13), two on retainers (6,10), one on infant orthopaedic treatment (11), and finally, one study on cleft palate treatment (12). The studies by Konst *et al.* (11) and Severens *et al.* (12) concerned the same cohort from the Dutch-cleft study (16), with cost data based on resource utilization from the latter randomized controlled trial (RCT) (16). Of the four studies on appliances, only two included comparable forms of appliance treatment, namely crossbite treatment using Quad Helix or expansion plates (7,13).

Five studies were based on cost-minimization analyses (6–9,13), although in the study by Hermanson *et al.* (13), the cost-minimization analysis was erroneously described as a cost-benefit analysis. Three studies used a cost-effectiveness analysis (10–12). Only two studies (6,7) included a societal perspective.

Evaluation of study quality and evidence

The quality of four (8,11–13) of the eight studies was assessed as limited, three (6,7,9) as moderate, and one as high (10). The reason for assessing the articles from Konst *et al.* (11) and Severens *et al.* (12) as limited were that the number of included patients did not correspond to the number of patients included in the original RCT. Thus, in the Dutch-cleft study by Pahl *et al.* (16), which formed the basis for the calculations, 23 patients per group was estimated to be needed. However, in the study by Severens *et al.* (12), 23 and 20 patients per group were used for the economic evaluation. The other two studies (8,13) of limited study quality both had a retrospective design. One RCT from UK (10), presented with high transparency and included almost 400 patients, was assessed as high study quality.

Under the GRADE system for grading the evidence, study quality is denoted as high, moderate, or limited (Table 2). Because of the broad range of aims among most of the studies (6,8–12), and the few studies of adequate quality covering the same comparison of appliances (7,13), no evidence base for economic aspects of orthodontic treatment emerged.

Discussion

No evidence of cost effectiveness of orthodontic interventions emerged from this review, primarily because so few relevant studies could be included and all but two concerned different methods. Of the two studies investigating the same orthodontic methods (7,13), only one (7) was assessed as having moderate study quality. However, when there are few relevant published studies, as in the present case, it is important to bear in mind that lack of evidence does not necessarily imply lack of effect, including economic outcome. Thus, there is an urgent need for further economic analysis of orthodontic interventions. It can also be noted that our results coincide with a recent systematic review regarding results for economic

evaluation of diagnostic methods used in dentistry (17). That review highlighted a paucity of economic evaluations of diagnostic methods and a need for studies from various perspectives with well-defined research questions and measures of the cost and effectiveness (17).

Of the eight studies remained for the final analysis, four studies considered different orthodontic appliances (7–9,13) and of these only two included comparable forms of appliance treatment, namely crossbite treatment using Quad Helix or expansion plates (7,13). In addition, two studies concerned retainers (6,10), one on infant orthopaedic treatment (11) and finally one study on cleft palate treatment (12). Since few studies have been undertaken and methodologically there was between-study heterogeneity, no cumulative evidence or meta-analysis could be assessed regarding the health economics of orthodontic interventions. Consequently, much of what we practice lacks sufficient evidence about the health economics of orthodontic interventions.

The studies assessed in this review were predominantly from Europe and especially from Sweden. Thus, there were few or no studies from other parts of the world. It seems that Swedish researchers have taken the initiative in presenting combined economic and orthodontic outcomes in studies of health economics. This may be seen in the context of the Swedish system of subsidized dental care, including publicly funded orthodontic treatment for children and adolescents.

The question has arisen as to whether lack of evidence based on economic evaluations of orthodontic interventions presents a problem. Some authors, for instance Anderson (18), claim that systematic reviews of economic studies are of limited relevance, primarily because of problems of transferability: as the costs incurred in studies from one country are seldom representative of another country (18). Thus, studies of cost effectiveness are seldom relevant for other countries and an example of this may partly be mirrored in a Finnish study (19) where the factors affecting the costs of children's free orthodontic treatment were studied and significant differences in treatment patterns and costs between different health centres were found.

What could be done to enhance comparisons between economic studies of orthodontic interventions undertaken in different countries? One way would be to include specific data not only on economic results but also on resource utilization. For instance, specifications of time input by dentists and dental nurses, respectively, and specifications of how overhead costs were calculated, as a percentage or as absolute added costs. Studies from different countries could then be compared more easily. Despite the issue of transferability, assessment of studies identified in systematic reviews is important, as the evidence may serve as guidelines for planning and management of orthodontic interventions.

One attempt to find ways of comparisons between studies of cost effectiveness in orthodontic care is described in a study of exploratory nature (20). There were large differences of cost effectiveness found between included practitioners from seven countries in Europe. However, the differences were mainly due to differences in

Table 2. Summary of studies included in the quality assessment, listed in descending order, from the latest publication. RCT = randomized controlled trial.

Reference, country	Aim of the study	Study design follow-up	Study population	Economic and orthodontic outcomes	Study quality
Tynelius Edman <i>et al.</i> (6), Sweden	Compare costs for three retention methods; Vacuum formed retainer + canine to canine retainer (V-CTC), vacuum formed retainer + stripping (V-S), and positioner (P)	Costs based piggy-back on one earlier published RCT. 2 years follow-up	Adolescents 17–19 years of age. Group V-CTC: N = 24 (17 girls and 7 boys). 1 dropout. Group V-S: N = 23 (14 girls and 9 boys). 2 dropouts. Group P: N = 22 (13 girls and 9 boys). 3 dropouts	Cost-minimization analysis. Direct, indirect, and societal costs. Equally efficient in retaining orthodontic treatment results. Costs in Euro per patient: V-CTC 457; V-S 451; Positioner 429; V-CTC retainer was the least cost effective	Estimated treatment times. Average times used for calculations. Societal perspective. Intention to treat. Low attrition. Moderate study quality
Petrén <i>et al.</i> (7), Sweden	Compare costs of unilateral posterior crossbite correction using Quad Helix (QH) or expansion plate (EP)	Costs based piggy-back on one earlier published RCT. 3 years follow-up	Group QH: N = 20, mean age 9.0 years (range 7.8–10.2 years), 11 girls and 9 boys. Group EP: N = 20, mean age 8.5 years (range 7.5–9.5 years), 13 girls and 7 boys	Cost-minimization analysis. Direct, indirect, and societal costs. Success rate of correction, transverse expansion. QH and EP had equally successful crossbite corrections. QH offers significant economic benefits over EP. QH had lower direct and indirect costs, also fewer failures needing retreatment. Costs in Euro QH 1031 versus EP 1395 ($P < 0.01$)	Unclear about variability of some costs compared to treatment time. Societal perspective. Intention to treat analysis. No attrition. Moderate study quality
Laag and Ström (8), Sweden	Compare results and costs at a general dentist clinic (GDC) and a special orthodontic clinic (SOC) for treatment of crowding, increased overjet, lateral crossbite, and dental displacement	Retrospective CCT. Cases at GDC and controls at SOC. Follow-up unknown	Removable appliance at GDC and fixed appliance at SOC. Group GDC: N = 31, mean age 11.4 years (range 8.2–14.2 years), 20 girls and 11 boys. Group SOC: N = 19, mean age 13.1 years (11.1–14.2 years), 16 girls and 3 boys	Cost-minimization analysis. Weighted Peer Assessment Rating (WPAP-index). GDC 69.0 (± 10.6) versus SOC 84.1 (± 7.1) ($P < 0.001$). Costs SEK GDC 3.525 versus SOC 5.510. Trained general dentists may achieve sufficiently qualitative results at a reasonably low cost. The results tend to be better at the SOC but more expensive than at the GDC	Simplified calculations of costs. Retrospective CCT. Non-cooperative patients were not registered, thus only successful treatment was counted. Limited study quality
O'Brien <i>et al.</i> (9), UK	Evaluate the effectiveness and costs of early versus adolescent orthodontic treatment of Class II division 1 malocclusion	Multicentre. RCT. 10 years follow-up	Group early Twin-block appliance: N = 89, mean age at start 9.7 years (SD 0.98), 41 girls and 48 boys. Removable or fixed appliance for the adolescent group: N = 85, mean age 9.8 years (SD 0.98), 39 girls and 46 boys. Early treatment group: mean age 12.4 years (SD 1.16) at second treatment phase. Adolescent treatment group: mean age 12.1 years (SD 1.1)	Cost-minimization analysis. No difference with respect to skeletal pattern, extraction rate, patient self-esteem. Early twin-block treatment incurred higher costs 3913 versus 3018 \$/patient. Early treatment with removable appliance followed by further treatment in adolescence at the appropriate time does not result in any meaningful long-term benefit over one course of treatment started in the late mixed dentition	No sensitivity analysis. Intention to treat analysis. 19% dropout rate. Moderate study quality

Table 2. Continued

Reference, country	Aim of the study	Study design follow-up	Study population	Economic and orthodontic outcomes	Study quality
Hichens <i>et al.</i> (10), UK	Compare cost-effectiveness of Hawley retainers (HR) and vacuum-formed retainers (VFR)	RCT. 6 months follow-up	Group HR: N = 196, mean age 14.8 years, 124 girls and 92 boys. Group VFR: N = 201, mean age 15.0 years, 119 girls and 82 boys	Cost-effectiveness analysis. VFR was more cost effective than HR from all perspectives, i.e. National Health Service, the practices and the patients	Well performed and good transparency of calculations. Power analysis. Drop-outs presented. High study quality
Konst <i>et al.</i> (11), The Netherlands	Investigate the cost-effectiveness of infant orthopaedic treatment (IO) of unilateral cleft lip and palate compared with no such treatment (NT) on speech development	Three-centre RCT. 2 years follow-up	20 patients randomized to two groups. Group IO: N = 10, age 2.5 years, 2 girls and 8 boys. Group NT: N = 10, age 2.5 years, 1 girl and 9 boys	Cost-effectiveness analysis. IO: Cost of treatment by orthodontists per patient €1460. NT: €419. ICER: incremental cost-effectiveness ratio, €1041 per 1.34 point speech quality improvement. Regarding speech in cleft patients, IO group had a statistically significant higher rating than the NT group	Selection of subjects unclear, i.e. unknown if consecutive patients. High attrition rate. Interpretation unclear. Number of patients not corresponding to the power analysis in original study. Limited study quality
Severens <i>et al.</i> (12), The Netherlands	Investigate cost effectiveness of complete unilateral cleft palate treatment from pre-surgical orthopaedic treatment (PSOT)	Three-centre RCT. PSOT compared with none-PSOT. 18 weeks follow-up	Group PSOT: N = 23, age 2 weeks, gender distribution not presented. Group none-PSOT: N = 20, age 2 weeks, gender distribution not presented	Cost-effectiveness analysis. Surgical lip closure, no significant difference. PSOT US\$ 852 versus none-PSOT US\$ 304. Regarding the incremental cost-effectiveness ratio, PSOT was more expensive than none-PSOT and had an insignificantly lower effect	No information about attrition. Number of patients not corresponding to the power analysis in original study. Limited study quality
Hermanson <i>et al.</i> (13), Sweden	Compare effects, complications, discomfort, and costs of unilateral crossbite correction using Quad-helix appliance (QH) and removable plate appliance (RP)	Retrospective CCT. 2 years follow-up	Group QH: N = 27, mean age 9.8 years (range 7.3–13.0 years), 14 girls and 13 boys. Group RP: N = 17, mean age 9.5 years (range 6.8–14.3 years), 12 girls and 5 boys	Cost-minimization analysis. Successful treatment: QH 26 of 27 and RP 16 of 17. Complications equal. Costs QH: 1223 SEK versus RP 1714 SEK. The cost analysis showed a 40% higher treatment cost with RP than QH. Treatment with RP takes longer than QH treatment	Selection of subjects unclear, i.e. unknown if consecutive patients. Economic analysis not a cost-benefit analysis but cost minimization, thus the authors have labelled the economic analysis incorrectly. Retrospective CCT. Limited study quality

costs, not in effects. The authors stressed the importance of including cost-effectiveness analyses in orthodontics in order to achieve ‘value for money’ (20). Moreover, a retrospective multicentre study from the UK had a focus on costs of direct health services. Costs were assessed of combined orthodontic and surgical treatment of dentofacial deformity. It was demonstrated that orthodontic costs on average comprised 25 per cent of the total treatment cost and this is stated to be inexpensive. Unfortunately, the effectiveness of orthodontic care was not considered in this study (21).

This systematic literature review was conducted according to Goodman’s model (5) and comprised four steps: 1. definition of

research question, 2. formulation of a plan for literature search, 3. literature search and retrieval of publications, and 4. data extraction, interpretation, and evaluation of evidence from the literature retrieved. This model fulfills the criteria of repeatability and minimizes the risk that the conclusions will be affected by chance or arbitrariness. In a further step, the study quality of the articles was classified as low, moderate, or high according to criteria suggested by Drummond *et al.* (2). In order to avoid subjectivity in the assessments, the independent observers evaluated the quality of separate studies and the overall level of evidence. In 90 per cent of cases, the different reviewers’ evaluation and grading of the full-text articles

Figure 1. Flow of information through the different phases of the systematic review.

were in agreement; consensus on the remainder was reached after rereading and discussion.

One challenge encountered in this systematic review was that although the literature search identified as many as 1838 studies, 989 were deleted as duplicates. Furthermore, 823 articles did not meet the stated objective of the review and only 26 articles remained for the final analysis. This is not, however, an unusual outcome for systematic reviews: the initial literature search was intentionally designed to be broad enough to ensure that no article would be overlooked or disregarded.

Conclusions

This systematic review disclosed that few orthodontic studies have presented both economic and clinical outcomes. There is currently insufficient evidence available about the health economics of orthodontic interventions. Further investigation is warranted.

Funding

Halland County Council (Dnr HALLAND-457871); Swedish Dental Society (Dnr 2012/368).

References

1. Kumar, S., Williams, A.C. and Sandy, J.R. (2006) How do we evaluate the economics of health care? *European Journal of Orthodontics*, 28, 513–519.
2. Drummond, M.F., Scholpher, M.J., Torrence, G.W., O'Brien, B. and Stoddart, G.L. (2005) *Methods for the Economical Evaluation of Healthcare Programs*. Oxford University Press, Oxford, 3rd edn.
3. Tickle, M. (1997) Clinical effectiveness and primary dental care. 2. The influence of health economics. *Primary Dental Care*, 4, 85–87.
4. Buck, D. (2000) Economic evaluation and dentistry. *Dental Update*, 27, 66–73.
5. Goodman, C.S. (2014) HTA 101. (2014) Introduction to health technology assessment. http://www.nlm.nih.gov/nichsr/hta101/HTA_101_FINAL_7-23-14.pdf.
6. Tynelius Edman, G., Lilja-Karlander, E. and Petré, S. (2014) A cost-minimization analysis of an RCT of three retention methods. *The European Journal of Orthodontics*, 36, 436–441.
7. Petré, S., Bjerklin, K. and Bondemark, L. (2011) Stability of unilateral posterior crossbite correction in the mixed dentition: a randomized clinical trial with a 3-year follow-up. *American Journal of Orthodontics & Dentofacial Orthopedics*, 139, e73–e81.
8. Laag, B. and Ström, C. (2009) Comparisons of similar patients treated by general dental clinicians and orthodontic specialists. Outcome and economic considerations. *Swedish Dental Journal*, 33, 67–73.

9. O'Brien, K., et al. (2009) Early treatment for Class II Division 1 malocclusion with the Twin-block appliance: a multi-center, randomized, controlled trial. *American Journal of Orthodontics & Dentofacial Orthopedics*, 135, 573–579.
10. Hichens, L., Rowland, H., Williams, A., Hollinghurst, S., Ewings, P., Clark, S., Ireland, A. and Sandy, J. (2007) Cost-effectiveness and patient satisfaction: Hawley and vacuum-formed retainers. *The European Journal of Orthodontics*, 29, 372–378.
11. Konst, E.M., Prahl, C., Weersink-Braks, H., De Boo, T., Prahl-Andersen, B., Kuijpers-Jagtman, A.M. and Severens, J.L. (2004) Cost-effectiveness of infant orthopedic treatment regarding speech in patients with complete unilateral cleft lip and palate: a randomized three-center trial in the Netherlands (Dutchcleft). *Cleft Palate Craniofacial Journal*, 41, 71–77.
12. Severens, J.L., Prahl, C., Kuijpers-Jagtman, A.M. and Prahl-Andersen, B. (1998) Short-term cost-effectiveness analysis of presurgical orthopedic treatment in children with complete unilateral cleft lip and palate. *Cleft Palate Craniofacial Journal*, 35, 222–226.
13. Hermanson, H., Kuroi, J. and Rönnerman, A. (1985) Treatment of unilateral posterior crossbite with quad-helix and removable plates. A retrospective study. *The European Journal of Orthodontics*, 7, 97–102.
14. Guyatt, G.H., et al. (2011) GRADE guidelines: 4. Rating the quality of evidence—study limitations (risk of bias). *Journal of Clinical Epidemiology*, 64, 407–415.
15. Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G. and The PRISMA Group (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Medicine*, 6, e1000097.
16. Prahl, C., Kuijpers-Jagtman, A.M., van't Hof, M.A. and Prahl-Andersen, B. (2001) A randomized prospective clinical trial into the effect of infant orthopaedics on maxillary arch dimensions in unilateral cleft lip and palate (Dutchcleft). *European Journal of Oral Science*, 109, 297–305.
17. Christell, H., Birch, S., Horner, K., Lindh, C., Rohlin, M. and SEDENTEXCT Consortium (2014) Economic evaluation of diagnostic methods used in dentistry. A systematic review. *Journal of Dentistry*, 42, 1361–1371.
18. Anderson, R. (2010) Systematic reviews of economic evaluations: utility or futility? *Health Economy*, 19, 350–364.
19. Järvinen, S. and Widström, E. (2002) Determinants of costs of orthodontic treatment in the Finnish public health service. *Swedish Dental Journal*, 26, 41–49.
20. Deans, J., Playle, R., Durning, P. and Richmond, S. (2009) An exploratory study of the cost-effectiveness of orthodontic care in seven European countries. *The European Journal of Orthodontics*, 31, 90–94.
21. Kumar, S., Williams, A.C. and Sandy, J.R. (2006) Orthognathic treatment: how much does it cost? *The European Journal of Orthodontics*, 28, 520–528.

II

Original article

Oral health-related quality of life among children with excessive overjet or unilateral posterior crossbite with functional shift compared to children with no or mild orthodontic treatment need

Jenny Kallunki¹, Ola Sollenius², Liselotte Paulsson³, Sofia Petré³, Lillemor Dimberg⁴ and Lars Bondemark⁵

¹The Center for Orthodontics and Pedodontics, County Council Östergötland, Linköping, Sweden, ²Department of Orthodontics, County Council Halland, Halmstad, Sweden, ³Department of Orthodontics, Faculty of Odontology, Malmö University, Sweden, ⁴Orthodontic Department, Eastman institute, County Council Stockholm, Sweden and ⁵Department of Orthodontics, Faculty of Odontology, Malmö University, Sweden

Correspondence to: Jenny Kallunki, The Center for Orthodontics and Pedodontics, S-601 82 Norrköping, Sweden. E-mail: jenny.kallunki@regionostergotland.se

Summary

Objectives: To compare Oral health-related quality of life (OHRQoL) among 9-year-old children with excessive overjet (EO) to children with unilateral posterior crossbite (UPC) and children with normal occlusion (NO).

Materials and methods: The study sample sourced from 19 Public Dental Service Clinics in Sweden. Reported are baseline data originating from two controlled trials, one regarding UPC and the other focusing on EO. The NO children derive from the same trials. The UPC group comprised 93 children (45 boys and 48 girls), the EO group 71 children (36 boys and 35 girls), and the NO group 65 children (32 boys and 33 girls). In conjunction to a clinical examination, all children completed the Child Perceptions Questionnaire (CPQ₈₋₁₀) for evaluation of OHRQoL. The CPQ₈₋₁₀ comprises 25 questions grouped into four domains: oral symptoms, functional limitations, emotional, and social well-being. Validated questions about pain in the jaws and face were also included.

Results: The total mean CPQ score was 5.1 for the UPC, 7.4 for the EO, and 4.4 for the NO group, showing a significant difference between the UPC and EO ($P = 0.048$) and between EO and NO group ($P = 0.012$). These differences remained when adjusted for the confounders' caries, trauma, enamel defects, and headache. No difference between UPC and NO was found. The EO children also reported significantly higher scores in the domains emotional and social well-being ($P = 0.039$ and $P = 0.012$).

Limitations: The study would be strengthened if a longitudinal design had been performed.

Conclusion: Children with EO reported significantly lower OHRQoL compared to children with UPC or NO. The children generally reported low CPQ scores that imply an overall fairly good OHRQoL.

Introduction

The World Health Organization (WHO) states that oral health is essential to both general health and quality of life. Good oral health implies being free from diseases and disorders that could limit a person's capacity in biting, chewing, smiling, speaking, or negatively affect psychosocial well-being (1). One of the most frequent oral conditions is malocclusions and frequency ranges between 39 and 93%, whereas some of the most common specific malocclusions are excessive overjet (EO) with Class II malocclusion and unilateral posterior crossbite (UPC) (2–5).

Several indices exist to assess orthodontic treatment need in terms of the severity of a malocclusion. However, a person's self-perception of the malocclusion may not always be related to malocclusion severity, and outcomes related to patients' values or perceptions of malocclusions are often lacking. A recent systematic review confirmed that orthodontic research seems to have overlooked patient values among children while there have been many studies measuring morphological differences (6). Therefore, to include the patient's perspective is as important as clinical registrations when making decisions regarding treatments or performing research (6,7). Consequently, to incorporate patient's perspectives enables a wider evaluation of treatment need, and furthermore, treatment success (8).

The term used when evaluating patient's perceptions of psychosocial well-being and oral health is oral health-related quality of life (OHRQoL). Self-report questionnaires are commonly used to register and evaluate patients' perspective. A common instrument used in child populations is the Child Perceptions Questionnaire (CPQ) (9), and this instrument enables assessment of the OHRQoL of children in order to provide additional information when making clinical decisions or evaluating the success of treatments or interventions.

In a recent systematic review (10), it was found that malocclusions have negative effects on children's and adolescent's OHRQoL, and especially if the malocclusions were present in the aesthetic zone. However, a majority of the high-quality studies originated from Brazil, and an additional finding was the lack of OHRQoL studies with European origin. It is known that OHRQoL may differ between countries and societies, since a self-perceived measure can be dependent upon the context of the child (11, 12). Furthermore, there are no studies that explicitly investigate how the specific malocclusions UPC with functional shift and EO may affect the OHRQoL in young children. It has been reported that if UPC with functional shift, i.e. a forced guidance of the mandible where the mandible deviates to the crossbite side, is left untreated adverse effects on the temporomandibular joints, pain in the masticatory system, and altered facial growth including facial asymmetry can occur (13–15). In addition, EO, especially without complete lip seal, has greater risk of dental trauma (5, 16) and can also be associated with bullying (17). Since pain, abnormal facial appearance, and bullying are related to specific malocclusions, it is important to investigate what influence specific different malocclusions may have on the OHRQoL in children. Likewise, it is important in this context to include children with normal occlusion (NO) and with no or mild orthodontic treatment need as controls in the evaluations. Therefore, the aim of this study was to investigate the OHRQoL among 9-year-old children in mixed dentition and compare the OHRQoL of children with UPC to children with EO as well as with children with NO with no or mild orthodontic treatment need. The hypothesis was that children with EO have lower OHRQoL than children with UPC with functional shift or children with no or mild orthodontic treatment need since EO occur in the aesthetic zone.

Subjects and methods

The study sample of the children was sourced from 19 Public Dental Service Clinics in Sweden. The sample was comprised of baseline data from children that origin from two different controlled trials, one regarding UPC with functional shift and the other focusing on EO combined with Class II malocclusion. Both trials also include children with NO and with no or mild orthodontic treatment need. To ensure the representativeness of the sample, all children were selected from clinics located in different socioeconomically areas, and the clinics were located in communities ranging from 10 000 up to 140 000 inhabitants.

The inclusion criteria considering the UPC group were consecutive children, boys and girls, in the mixed dentition (between 8 and 10 years of age), with UPC with functional shift of more than 1 mm, including the first permanent molars and a Class I occlusion. In addition, no sucking habits or ceased sucking habits should have been evident at least 1 year before the trial was started.

The inclusion criteria for the group of patients with EO were consecutive children, boys, and girls, in the mixed dentition (between 8 and 10 years of age), with EO (≥ 6 mm) and first permanent molars in Class II malocclusion. Furthermore, no sucking habits or ceased sucking habits should have been evident at least 1 year before the trial was started.

The inclusion criteria for the normal group were consecutive children, boys and girls, in the mixed dentition (between 8 and 10 years of age), with NO and with no or mild orthodontic treatment need, i.e. Index of Orthodontic Treatment Need – Dental health Component (IOTN-DHC) (18) of 1 or 2.

The exclusion criteria were craniofacial syndromes, children who had previously undergone orthodontic treatment, or severe crowding of teeth (extraction of teeth necessary).

All children and their parents were invited to attend and were asked to give their informed consent before entering the trial. The trial protocol and informed consent form were approved by the Regional Ethical Review Board in Lund (Dno: 2012/650) and Linköping (Dno: 2012/302–31), Sweden. Data collection took place from December 2013 to April 2017.

CPQ₈₋₁₀ questionnaire

To evaluate the OHRQoL among the children in the three groups, the CPQ₈₋₁₀ (19), which is validated for children 8–10 years of age, was used. The CPQ₈₋₁₀ comprise 25 questions grouped into four domains: oral symptoms (OS), functional limitations (FL), emotional well-being (EW), and finally social well-being (SW). Each question has five answering possibilities, i.e. 'never' (scoring 0), 'once or twice' (scoring 1), 'sometimes' (scoring 2), 'often' (scoring 3), and 'every day or almost every day' (scoring 4). The domains were added in order to produce a total score. The questionnaire also contained two global self-ratings of oral health using a four-point scale ranging from 'excellent' to 'poor' and overall well-being using a four-point scale ranging from 'not affected at all' to 'very much affected'.

The total score ranged between 0 and 100, with subscale scores range of each domain from 0 to 20 (OS, FL, EW) or 40 (SW). If answer alternatives were not applicable for the child, the question was given the median input of the total score. The higher scores corresponded to poorer quality of life status.

Finally, the children also answered validated questions about pain in the jaws and face (20).

The questionnaire was filled out in conjunction to the clinical examination and the children were instructed to answer the questions without any support from their parents. However, all children had the

opportunity to ask questions about the questionnaires, and if needed, to get support with reading the questions by clinical staff personnel.

Clinical and dental record measures

The clinical measures and the determinations were performed by two experienced orthodontists (JK, OS) following a prespecified protocol that was used to collect data about malocclusions, orthodontic treatment need, caries, enamel defects, incisors with or without lip closure, and if any trauma had occurred. Alginate impressions were taken of each child who participated in the trial and study casts were made and trimmed after index in central occlusion. Study casts were evaluated according to the method of Bjoerk *et al.* (21) in order to determine malocclusions in sagittal, vertical, and transversal relations as well as determination of crowding, spacing, tooth rotations, overjet, and overbite.

Measurements of overjet and overbite were made with a stainless-steel ruler while crowding and spacing was determined by a digital sliding calipers (Digital 6, 8M007906, Mauser-Messzeug GmbH, Oberndorf/Neckar, Germany). All measurements were made to the nearest 0.5 mm.

The dental health component of the index of orthodontic treatment need (IOTN-DHC) was used to rank the treatment need (18). This index classifies the treatment need into five grades: no need 1; mild/little need 2; moderate/borderline need 3; severe need 4; and extreme need 5.

In combination with the clinical examination, dental records of the patients were used to support the determination of caries of the anterior teeth (the aesthetic zone, i.e. canine to canine in both arches). A caries lesion was characterized as caries identified by probing.

Enamel defects were registered for permanent teeth and in the evaluation dichotomized into the presence of quantitative decay of enamel (hypoplasia) or qualitative alteration in the transparency (opacity) of the enamel, and no enamel defect.

Reporting of perceived trauma was affirmed by the dental records and in the investigations dichotomized into trauma or not.

Calibration of operators

Before diagnostics and study cast evaluations in this trial, 20 study casts from patients not included in this study were separately examined by the observers. Ten patients with UPC and 10 patients with EO were evaluated. In each patient, 26 decisions had to be performed according to a pre-set protocol regarding diagnostic parameters, malocclusion traits, measurements of overjet, and overbite as well as determination of treatment need according to IOTN-DHC. The two independent observers (JK, OS), made 26 decisions in each of the 20 study casts. Consequently, each observer had to carry out 520 decisions, and concordance between the observers was reached in 97.1% of the decisions. The inter-examiner disagreements of 2.9% were resolved by discussions to reach consensus. Most of the disagreements dealt with determination of treatment need according to IOTN-DHC.

Sample size estimation and statistical analysis

The criterion for significance (α) was set at 0.05. The test was two-tailed, which means that an effect in either direction was interpreted. With a proposed sample size of 60 in each group, the trial had a power of 81.6 % ($1 - \beta$) to yield a statistically significant result. This computation assumed that the difference in proportions was 0.25 (25 % difference of quality of life between the groups). The difference of 25% was selected as the smallest difference in the sense that any smaller difference would not be of clinical or substantive significance. It was also assumed that this effect size was reasonable

in the sense that an effect of this magnitude could be anticipated in this field of research.

The data were statistically analyzed by using the IBM SPSS Statistics version 24. Means between groups were compared using ANOVA. Chi-square analysis and Fisher's exact test were used to determine difference between groups regarding categorical data. General linear model in combination with Tukey's test were performed when corrections were made for the following background/confounding variables: gender, caries, enamel defects, headache, and trauma. Differences with probabilities of less than 5% ($P < 0.05$) were considered to be statistically significant.

Results

Sample characteristics

In the UPC group, there were 93 children, 45 boys and 48 girls, with a mean age of 9.1 years (SD 1.02) whereas in the EO group 71 children, 36 boys and 35 girls, with a mean age of 9.5 years (SD 0.53) participated. The NO group consisted of 65 children, 32 boys and 33 girls, with a mean age of 9.4 years (SD 1.01).

Thus, the gender distribution between the groups was similar but a small albeit significant difference between the UPC and EO groups regarding age was found ($P = 0.025$).

All children in the UPC and NO group had Class I molar relation while all in the EO group showed Class II division 1 malocclusion. A significant larger overjet ($P < 0.001$) and overbite ($P < 0.001$) was found in the EO group compared to the UPC group, 8.2 (SD 1.50) and 3.7 mm (SD 1.40) versus 3.1 (SD 1.29) and 2.3 mm (SD 1.43). The overjet and overbite were 3.2 mm (SD 1.11) and 3.0 mm (SD 1.12) in the NO group. There was also a significant larger overjet ($P < 0.001$) in the EO compared to the NO group. Consequently, significantly more children without lip closure ($P < 0.001$) were found in the EO group (Table 1). No significant differences were revealed in overjet and overbite between the UPC and NO group.

The prevalence of trauma, enamel defects, and headache were low in all three groups, and without any significant inter-group differences (Table 1).

All children in the UPC group had moderate to severe treatment need indicating that the crossbite was of functional disturbing nature, i.e. a forced guidance of the mandible deviating the midline of the mandible to the crossbite side. In the EO group, all children had severe to extreme treatment need due to the EO, while in the NO group the children showed no or mild orthodontic treatment need (IOTN-DHC 1 or 2), Table 1. Furthermore, in each group there were few children with mild to moderate crowding or spacing in the aesthetic zone and with no inter-group differences.

Oral health-related quality of life (OHRQoL)

There were no dropouts, all questionnaires were completely filled in, and none had to be excluded due to missed data.

The total CPQ mean score was 5.1 for the UPC group, 7.4 for the EO group, and 4.4 for the NO group. The EO group had significant higher total CPQ mean score than the UPC group ($P = 0.048$) and the EO group also had significantly higher total CPQ mean score than the normal group ($P = 0.012$), see Table 2 and Figure 1. No gender differences were found.

For the UPC group, the mean subscale score was 3.3 for OSs, 0.9 for FLs, and 0.5 for emotional and 0.5 for SW (Table 2). The corresponding data for the EO group was 3.9, 1.2, 1.2, 1.2, and for the NO group 3.0, 0.7, 0.5, and 0.2. The children with EO reported significantly higher scores in the domains EW compared to UPC-group

Table 1. Descriptive data for the three groups of children: excessive overjet (EO), unilateral posterior crossbite (UPC), and normal occlusion (NO).

	EO N = 71		UPC N = 93		NO N = 65		Group differences
	N	%	N	%	N	%	
Gender							
Boys	36	50.7	45	48.4	32	49.2	No group differences
Girls	35	49.3	48	51.6	33	50.8	$P = 0.985$ (C)
IOTN							
5	15	21.1					EO versus UPC $P < 0.001$ (F)
4	56	78.9	18	19.4			EO versus NO $P < 0.001$ (F)
3			75	80.6			UPC versus NO $P < 0.001$ (F)
2					7	10.8	
1					58	89.2	
Lips							
With closure	50	70.4	92	99	65	100	EO versus UPC $P < 0.001$ (F)
Without closure	21	29.6	1	1	0	0	EO versus NO $P < 0.001$ (F)
Caries	0	0	0	0	1	1.5	UPC versus NO $P = 1.00$ (F)
Trauma	5	7	3	3.2	2	3.1	No group differences
Enamel defects	5	7	1	1.1	2	3.1	$P = 0.284$ (F)
Headache	3	4.2	8	8.6	4	6.2	No group differences
							$P = 0.558$ (F)

C: Chi-squared test; F: Fisher's exact test.

Table 2. CPQ8-10 scores for the three groups; excessive overjet (EO), unilateral posterior crossbite (UPC), and normal occlusion (NO).

	CPQ total		Oral symptoms (OS)		Functional limitations (FL)		Emotional well-being (EW)		Social well-being (SW)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
EO (N = 71)	7.4	8.2	3.9	2.7	1.2	1.8	1.2	2.4	1.2	3.2
UPC (N = 93)	5.1	5.1	3.3	2.7	0.9	1.6	0.5	1.2	0.5	0.8
NO (N = 65)	4.4	5.2	3.0	2.7	0.7	1.3	0.5	1.8	0.2	0.8
p-level when adjusted for pos- sible confounders (caries, trauma, MIH, headache) ^a	EO versus UPC $P = 0.048^b$ EO versus NO $P = 0.012^b$						EO versus UPC $P = 0.039^b$		EO versus NO $P = 0.012^b$	

^aGeneral linear model.^bTukey's test.

($P = 0.039$) and SW compared to NO-group ($P = 0.012$) (Table 2). For the domains OS and FL, no significant differences were detected between the groups.

When corrections were made for the background variables/possible confounders; gender, caries, trauma, enamel defects, and headache, the significant difference between the groups still remained (Table 2). Headache was the only variable that showed significant negative impact on the OHRQoL and that in the domain OS ($P < 0.05$).

Discussion

This study analyzed the impact a UPC with functional shift or an EO may have on young children's OHRQoL and comparisons

were also made to children with no or mild orthodontic treatment need. The main finding was that children with EO reported a lower self-perceived OHRQoL when compared with children with unilateral posterior cross-bite or children with no or mild orthodontic treatment need. In addition, no difference in OHRQoL could be detected between children with UPC and children with no or mild orthodontic treatment need. These findings support our hypothesis and are accordance with earlier research stating that malocclusions in the aesthetic zone have a negative impact on OHRQoL (10). The negative impact was evident in the domains of SW and EW. Perhaps not an unexpected finding, but an important finding seen in the context of the young age of the patients.

A somewhat surprising result was that children with UPC did not show reduced OHRQoL compared to children with no or mild

Figure 1. CPQ8-10 total score for children with unilateral posterior crossbite (UPC), excessive overjet (EO), and normal occlusion (NO). A significant difference ($P = 0.048$) in mean total CPQ score between EO and UPC was found, there was also a difference between EO and NO ($P = 0.012$) but not for UPC and NO.

orthodontic treatment need. The explanation may be that children with crossbite are not yet aware of their malocclusion, or that the functional disturbance associated has not yet caused any problems that have reflected in reduced OHRQoL among the children with UPC.

Although the primary aim of this study was to compare children with EO and children with UPC as well as with children with NO, the children in the EO group also had significant increased overbite which may influence the OHRQoL. In addition, few children in each group without any inter-group differences showed small amount of crowding and spacing, having no influence on OHRQoL.

Assessing OHRQoL, by the use of CPQ₈₋₁₀, revealed generally low scores for the children participating in this study. This indicates that the children, even though the presence of malocclusions with severe to moderate orthodontic treatment need, perceive themselves of having an overall fairly good OHRQoL. It can also be underlined that there is no clear correlation or pattern between high IOTN scores and low OHRQoL (22).

One possible explanation for the generally low CPQ scores found in this study is the low prevalence of caries among the children. Children with caries are more likely to show a reduced OHRQoL and when compared with dental trauma or malocclusions, caries may have as much as twice the negative impact on OHRQoL (23). A target has been set by WHO, namely by year 2020, the average amount of caries measured as decade filled teeth in European 12-year-olds should not exceed 1.5. This ambition has already been achieved in Sweden since the year 1995 (24). Accordingly, no caries was found in the aesthetic zone in any of the children participating in this study.

Other important confounders that may impair OHRQoL are dental fear and dental pain (12). Age presents a negative correlation, as the risk for dental fear and potential behaviour management problems increases as the child's age decreases (25). In this study, all young children had together with their parents voluntarily accepted to participate in an additional clinical examination including impressions for study casts and all children fulfilled the examinations

and impressions without any problems. Thus, although not objectively verified, the children in this study did not show any signs of dental fear.

As shown in Figure 1, one child in the EO group and one in the normal group reported obvious high CPQ scores. To prevent misinterpretations of the results, the statistical analyses were repeated without these outliers. The result prevailed, implying that the group with EO still without the outlier presented significantly higher total CPQ score and higher scores in the domains of emotional and functional well-being when compared with the other groups.

There were rather few children reporting headache and there was no difference between the groups regarding the prevalence of headache. However, the only confounder in this study that exhibit to have a significant negative impact was headache. Hence, headache reduced OHRQoL, and especially for the domain OS. The impact of headache on OHRQoL is in accordance with other studies using the CPQ, however performed with older children (22). Migraine and tension-type headache are the most common types of headache in children, and can have a profound effect on the child, nevertheless the aetiology is usually medical or psychosocial (26).

Self-perceived orthodontic treatment need differs for children from different cultures (27). OHRQoL might also be affected by confounders that may be dependent upon cultural or social context (11, 12). The need for studies from different countries has been advocated since most available research regarding children and their self-perceived OHRQoL has been conducted in Brazil (10). A potential demographic impact is displayed when the children in this study scored lower on the CPQ, i.e. reported better OHRQoL than Brazilian peers with no need for orthodontic treatment (28). Whether this is due to differences regarding confounders is unclear.

The strengths of this study were the number of children participating and the well-defined groups including a NO group with no or mild orthodontic treatment need. Moreover, there were no drop-outs in this study, and clinical routines ensured that the children were able to respond to the questionnaire without the involvement of parents. In addition, the children were recruited from several clinics located in different communities to ensure an appropriate socioeconomic scope. Finally, the examiners were calibrated before evaluating clinical data and the influence of possible confounders was low.

The study would further be strengthened if psychological characteristics such as psychological well-being and self-esteem were incorporated. These dimensions have been shown to influence the OHRQoL for adolescents (29). Finally, a longitudinal study design would also strengthen the study. However, evaluating changes over time would mean postponing interventions for children with severe orthodontic treatment need.

Conclusions

Young children with EO reported significantly lower self-perceived OHRQoL compared to children with UPC with functional shift or children with NO and with no or mild orthodontic treatment need. No difference regarding self-perceived OHRQoL could be detected between children with UPC with functional shift when compared with children with NO and with no or mild orthodontic treatment need. However, despite the presence of malocclusions, the children generally reported low CPQ scores which imply an overall fairly good self-perceived OHRQoL.

Funding

This study was supported by the European Orthodontic Society Research Grant (2015) and the Region Halland and Region Östergötland, Sweden.

Acknowledgements

Our gratitude to the patients and their parents for participating in this study.

Conflict of interest

None to declare.

References

- http://www.who.int/oral_health/en//www.who.int/oral_health/en/ (24 March 2017, date last accessed).
- Helm, S. (1968) Malocclusion in Danish children with adolescent dentition: an epidemiologic study. *American Journal of Orthodontics*, 54, 352–366.
- Thilander, B. and Myrberg, N. (1973) The prevalence of malocclusion in Swedish schoolchildren. *Scandinavian Journal of Dental Research*, 81, 12–21.
- Thilander, B., Pena, L., Infante, C., Parada, S.S. and de Mayorga, C. (2001) Prevalence of malocclusion and orthodontic treatment need in children and adolescents in Bogota, Colombia. An epidemiological study related to different stages of dental development. *European Journal of Orthodontics*, 23, 153–167.
- Dimberg, L., Lennartsson, B., Arnrup, K. and Bondemark, L. (2015) Prevalence and change of malocclusions from primary to early permanent dentition: a longitudinal study. *The Angle Orthodontist*, 85, 728–734.
- Tsichlaki, A. and O'Brien, K. (2014) Do orthodontic research outcomes reflect patient values? A systematic review of randomized controlled trials involving children. *American Journal of Orthodontics and Dentofacial Orthopedics*, 146, 279–285.
- Gilchrist, F., Rodd, H., Deery, C. and Marshman, Z. (2014) Assessment of the quality of measures of child oral health-related quality of life. *BMC Oral Health*, 14, 40.
- Sheiham, A., Maizels, J.E. and Cushing, A.M. (1982) The concept of need in dental care. *International Dental Journal*, 32, 265–270.
- Jokovic, A., Locker, D., Stephens, M., Kenny, D., Tompson, B. and Guyatt, G. (2002) Validity and reliability of a questionnaire for measuring child oral-health-related quality of life. *Journal of Dental Research*, 81, 459–463.
- Dimberg, L., Arnrup, K. and Bondemark, L. (2015) The impact of malocclusion on the quality of life among children and adolescents: a systematic review of quantitative studies. *European Journal of Orthodontics*, 37, 238–247.
- Kragt, L., Dhano, B., Wolvius, E.B. and Ongkosuwo, E.M. (2016) The impact of malocclusions on oral health-related quality of life in children: a systematic review and meta-analysis. *Clinical Oral Investigations*, 20, 1881–1894.
- Merdad, L. and El-Housseiny, A.A. (2017) Do children's previous dental experience and fear affect their perceived oral health-related quality of life (OHRQoL)? *BMC Oral Health*, 17, 47.
- Pinto, A.S., Buschang, P.H., Throckmorton, G.S. and Chen, P. (2001) Morphological and positional asymmetries of young children with functional unilateral posterior crossbite. *American Journal of Orthodontics and Dentofacial Orthopedics*, 120, 513–520.
- Egermark, I., Magnusson, T. and Carlsson, G.E. (2003) A 20-year follow-up of signs and symptoms of temporomandibular disorders and malocclusions in subjects with and without orthodontic treatment in childhood. *The Angle Orthodontist*, 73, 109–115.
- Andrade, A.d.a.S., Gameiro, G.H., Derossi, M. and Gavião, M.B. (2009) Posterior crossbite and functional changes. A systematic review. *The Angle Orthodontist*, 79, 380–386.
- Thiruvengkatchari, B., Harrison, J.E., Worthington, H.V. and O'Brien, K.D. (2013) Orthodontic treatment for prominent upper front teeth (Class II malocclusion) in children. *Cochrane Database Systematic Review*, 11:CD003452.
- de Oliveira, C.M. and Sheiham, A. (2003) The relationship between normative orthodontic treatment need and oral health-related quality of life. *Community Dentistry and Oral Epidemiology*, 31, 426–436.
- Brook, P.H. and Shaw, W.C. (1989) The development of an index of orthodontic treatment priority. *European Journal of Orthodontics*, 11, 309–320.
- Jokovic, A., Locker, D., Tompson, B. and Guyatt, G. (2004) Questionnaire for measuring oral health-related quality of life in eight- to ten-year-old children. *Pediatric Dentistry*, 26, 512–518.
- Nilsson, I.M., List, T. and Drangsholt, M. (2006) The reliability and validity of self-reported temporomandibular disorder pain in adolescents. *Journal of Orofacial Pain*, 20, 138–144.
- Bjoerk, A., Krebs, A. and Solow, B. (1964) A method for epidemiological registration of malocclusion. *Acta Odontologica Scandinavica*, 22, 27–41.
- Dimberg, L., Lennartsson, B., Bondemark, L. and Arnrup, K. (2016) Oral health-related quality-of-life among children in Swedish dental care: the impact from malocclusions or orthodontic treatment need. *Acta Odontologica Scandinavica*, 74, 127–133.
- Martins, M.T., Sardenberg, F., Bendo, C.B., Abreu, M.H., Vale, M.P., Paiva, S.M. and Pordeus, I.A. (2017) Dental caries remains as the main oral condition with the greatest impact on children's quality of life. *PLoS One*, 12, e0185365.
- Socialstyrelsen (2010) Karies hos barn och ungdomar. *En lägesrapport för år 2008*. 2010-3-5, 12–13.
- Wogelius, P., Poulsen, S. and Sørensen, H.T. (2003) Prevalence of dental anxiety and behavior management problems among six to eight years old Danish children. *Acta Odontologica Scandinavica*, 61, 178–183.
- Gofsteyn, J.S. and Stephenson, D.J. (2016) Diagnosis and management of childhood headache. *Current Problems in Pediatric and Adolescent Health Care*, 46, 36–51.
- Josefsson, E. (2010) Immigrant background and orthodontic treatment need. Quantitative and qualitative studies in Swedish adolescents. *Swedish Dental Journal*, suppl 207, 1–92.
- Simões, R.C., Goettems, M.L., Schuch, H.S., Torriani, D.D. and Demarco, F.F. (2017) Impact of malocclusion on oral health-related quality of life of 8-12 years old schoolchildren in southern Brazil. *Brazilian Dental Journal*, 28, 105–112.
- Agou, S., Locker, D., Muirhead, V., Tompson, B. and Streiner, D.L. (2011) Does psychological well-being influence oral-health-related quality of life reports in children receiving orthodontic treatment? *American Journal of Orthodontics and Dentofacial Orthopedics*, 139, 369–377.

III

Randomized Controlled Trial

An RCT on clinical effectiveness and cost analysis of correction of unilateral posterior crossbite with functional shift in specialist and general dentistry

Ola Sollenius¹, Sofia Petré² and Lars Bondemark²

Department of Orthodontics, ¹County Council, Halland, Halmstad and ²Malmö University, Malmö, Sweden

Corresponding to: Ola Sollenius, Department of Orthodontics, County Council, Halland, S-30246 Halmstad, Sweden. E-mail: ola.sollenius@regionhalland.se

Summary

Objectives: To evaluate the costs of quad-helix (QH) and removable expansion plate (EP) treatments performed either in specialist or general dentistry for the correction of unilateral posterior crossbite with functional shift in the mixed dentition.

Trial design: Four-arm parallel group multicentre randomized controlled trial.

Materials and methods: One hundred and ten patients with unilateral posterior crossbite with functional shift were recruited. The patients were randomized by an independent person not involved in the trial. The randomization used blocks of 20 and into the following four groups: QH treatments in specialist orthodontic clinics (QHS), QH treatments in general dentistry (QHG), EP treatments in specialist orthodontic clinics (EPS), and EP treatments in general dentistry (EPG). Blinding was accomplished of the outcome assessor and data analyst. A cost analysis was performed with reference to intention-to-treat (ITT), regarding direct costs, indirect costs, and societal costs (the sum of direct and indirect costs) for calculations of successful treatments alone and for retreatments when required. To determine which alternative has the lower cost, a cost-minimization analysis was undertaken, based on that the outcome of the treatment alternatives were broadly equivalent, so the difference between them reduces to a comparison of costs.

Results: In the QHS group, 28 of 28 patients were successfully corrected compared to 23 of 27 in the QHG group. Treatment with expansion plate was less successful: 18 of 27 patients in the EPS group and 18 of 28 in the EPG group. QH treatment performed in specialist orthodontic clinics had significantly lower costs than QH or EP treatment accomplished in general dentistry as well as EP treatments in specialist orthodontic clinics.

Limitations: Costs depend on local factors and should not be directly extrapolated to other locations.

Conclusion: Treatment of unilateral posterior crossbite in the mixed dentition is recommended to be performed by a specialist orthodontist using the QH appliance.

Trial registration: The trial was not registered.

Introduction

Unilateral posterior crossbite with functional shift is one of the most common malocclusions in the mixed dentition, with a reported prevalence ranging from 8.5 to 17 per cent (1,2). Left untreated, it may cause long-term effects on the growth and development of the jaws and teeth (3). Treatment of crossbite may consume a substantial part of the total resources in orthodontic care and, usually, is performed by rapid maxillary expansion, quad-helix (QH) or removable expansion plates (EP) (4).

In recent years, economic evaluations have become an integral component of health services (5). The main reason is that resources within the health sector (personnel, time, facilities, equipment, and knowledge) are limited (6). Thus, failure to analyse economic aspects of dental health service evaluations may result in unsustainable over-expenditure or withdrawal or reduction of services or resources in other areas of healthcare (7). In allocating resources, including dental care and orthodontics, health service purchasers need to take into account not only evidence of clinical effectiveness of treatment procedures but also relative costs, i.e. 'value for money' (8). Two features characterize health economic evaluation: the relationship between costs (input) and consequences (output) and preferential selection of diagnostic or therapeutic options, implying alternative ways of allocating resources.

Four types of evaluations can be used to gather evidence and compare the expected costs and consequences of different procedures. 'A cost-effectiveness analysis' is characterized by analysis of both costs and outcomes, where the outcomes of alternative methods might differ in magnitude. In a 'cost-minimization analysis', which is a form of cost-effectiveness analysis, typically this is used to describe the situation where the consequences of two or more treatments are broadly equivalent, so the difference between them reduces to a comparison of costs. In a 'cost-utility analysis', the quality of life following treatment is also evaluated, while in a 'cost-benefit analysis', the consequences are expressed in monetary units, for example, when distribution of resources to diverse areas of healthcare is evaluated (6).

A systematic review found insufficient evidence available about the health economics of orthodontic interventions (9); however, it is important to note that studies of costs may not always be possible to compare between countries or economic systems. An example of this may partly be mirrored in a Finnish study (10) where the factors affecting the costs of children's free orthodontic treatment were studied and significant differences in treatment patterns and costs between different health centres were found. One attempt to find ways of drawing comparisons between studies of costs in orthodontic care has been described in a study of exploratory nature (11). Large differences in costs were found between included practitioners from seven countries in Europe. Nevertheless, the differences were largely due to differences in costs not in effects.

In Finland and also in Sweden, the role of orthodontists in general dental care is essential in diagnostics and treatment planning, whereas general dentists may often provide a part of the treatment after consultation with, or under the supervision of, an orthodontist (12, 13). Nevertheless, studies comparing the costs of orthodontic treatments performed in general versus specialist dentistry are virtually non-existent. Consequently, we conducted a multicentre randomized controlled trial with the aim of comparing the costs of QH and EP for treatment of unilateral posterior crossbite with functional shift in the mixed dentition as well as to evaluate the costs of such crossbite treatments performed either in specialist or general

dentistry. It was hypothesized that both treatment methods are equal regarding costs as well as for the costs of treatment in general and specialist dentistry.

Material and methods

Between 2013 and 2018, we consecutively recruited 110 patients, i.e. all patients who attended and met the inclusion criteria, from 10 general dental health clinics at the Public Dental Health Service, Halland County Council, Sweden. The inclusion criteria were: mixed dentition (all incisors and first molars erupted), unilateral posterior crossbite with functional shift including the maxillary first molar, and no sucking habits or sucking habit ceased at least one year before the trial. The exclusion criteria were: craniofacial syndromes and children who had previously undergone orthodontic treatment.

Before the treatment started, two experienced orthodontists provided the patients and their parents with oral and written information about the trial and the responsible general dentist got the same information at this visit. Subjects were randomized into one of four groups: treatment in general dentistry with QH (QHG) or EP (EPG) or in specialist orthodontic clinics with QH (QHS) or EP (EPS). The randomization procedures were prepared and carried out by an independent person not involved in the study and the randomization used blocks of 20. Thus, the orthodontists carried boxes of already prepared envelopes. Six opaque envelopes were prepared with 20 sealed notes in each (5 notes for each group) and for every new patient in the study, a note was extracted from the first envelope. When the envelope was empty, the second envelope was opened, and the 20 new notes were extracted as the patients were recruited to the trial. This procedure was then repeated 4 more times. After enrolment and randomization, all patients were treated according to a present standard concept, detailed descriptions of the design and handling of the appliances (see The Quad-helix and The removable expansion plate), and also a form to fill out, for example, time for ordinary and emergency visits, parents travelling time and absence from work.

The ethics committee of Lund University, Lund, Sweden (Dnr: 2012/650), which follows the guidelines of the Declaration of Helsinki, approved the informed consent form and protocol. Thus, the patients and parents received oral and written information about the trial and then signed the consent form.

All clinics, three specialist orthodontic and 10 general clinics, used the same consent form and protocol for the treatment interventions QH and EP. The two orthodontists who provided the information to the patients had 15 years of experience and treated all the patients in the specialist orthodontic clinics and did not treat any patients at the general dental clinics. The 17 general dentists had between 1 and 30 years of experience of interceptive orthodontic treatments.

In this trial we used the intention-to-treat principle (ITT), i.e. all cases, whether successful or not as well as dropouts, were included in the final analysis.

Patients who declined to participate in the trial were offered orthodontic treatment with either QH or EP at their general practitioner's clinic.

The quad-helix

The QH consisted of a standard stainless steel arch (MIA system, 3M Unitek, Monrovia, California, USA) with stainless steel bands attached with glass ionomer cement on the maxillary first molars

Figure 1. Occlusal view of the quad-helix.

(Figure 1). The QH was activated 10 mm before placement and then reactivated every 6 weeks until a normal transverse relationship was achieved; no overcorrection was produced. Steel or elastic ligatures were used to secure the QH to the bands. The treatment result was retained for 3 months.

The removable expansion plate

The EP consisted of acrylic, with an expansion screw and stainless steel clasps on the primary and permanent first molars (Figure 2). The EP was activated 0.2 mm once a week by the patient until a normal transverse relationship was achieved; no overcorrection was produced. The dentist thoroughly instructed the patient to use the EP day and night except for meals and tooth brushing. Treatment progression was evaluated every 6 weeks, and the result was retained for 3 months. Adjustment of steel clasps was performed when needed.

Outcome measures

The outcome measures to be assessed in the trial were: 1. success rate of crossbite correction (yes or no), i.e. normal transverse relationships achieved for all teeth within 1 year; 2. if midline correction was achieved or not; 3. treatment duration—time in months to correct the crossbite to normal occlusion (if normal occlusion was not achieved, the treatment time was stated as 1 year, i.e. the total observation period); 4. total number and time of each visits including emergency visits and missed appointments. The primary outcomes were the cost outcomes; 5. direct costs such as costs for chair time were calculated based on the costs per hour for the general practitioner (2235 SEK/€214 and for the specialist in orthodontics (2626 SEK/€251) for the year 2017 and the material costs for the QH or the EP were based on the dental technician costs including bands, separation ligatures, and Mobile Intraoral Arch system; 6. indirect costs, i.e. parent's time absent from work when accompanying the patient to the dental clinic.

It was assumed that the parents of 8- and 10-year olds were between 35 and 45 years old. In 2017, the mean income per year for 35- to 45-year olds living in Sweden was 364 743 SEK/year (€35 071/year) or 278 SEK/hour (€26.82/hour). The mean income per hour was calculated by dividing mean income per year with 1700 working hours, then adding the social costs by factor 1.3142. The data were sourced from the Swedish National Bureau of Statistics (www.scb.se). All costs were based on 2017 prices and were expressed in Euros (€), SEK 100 = €10.4 at a mean currency value (www.xe.com).

Figure 2. Occlusal view of the removable expansion plate.

Blinding

Patients and trial staff could not be blinded because of the character of treatment but blinding was accomplished of the outcome assessor; thus, the assessor was unaware of the group to which the patient belonged or whether the data were for before or after treatment.

Direct costs

Direct costs comprised material costs and treatment time costs. Material costs were compiled and calculated according to average commercial prices. Treatment time costs, including the costs of the premises, dental equipment, maintenance, and cleaning, were calculated according to average commercial prices in Sweden; these figures were used to establish estimated costs for each unit in the study. Similarly, staff salaries, including payroll tax, were calculated for dental assistants, general dental practitioners, and orthodontists. Treatment time (in hours) for both scheduled and unscheduled appointments, along with the number of appointments, were registered on an individual basis, on a form included in the patient records.

Indirect costs

Indirect costs were defined as loss of income (wages plus social security costs) incurred by the patients' parents' absence from work to accompany the patient to the orthodontic appointment, i.e. total time (including waiting time) at the clinic and the travelling duration.

Cost-minimization analysis

A cost-minimization analysis was used to describe the situation since the consequences of the two treatments were broadly equivalent, so the difference between them reduces to a comparison of costs.

The sum of direct and indirect costs was defined as 'societal costs' and the cost minimization was calculated as follows:

1. The mean societal costs of successful cases only on completion of treatment within 1 year in all groups:
 - (a) the mean societal costs for all successful QH treatments at the specialist orthodontic clinic divided by the number of successful patients, i.e. societal costs for 28 treatments divided by 28 patients.
 - (b) the mean societal costs for all successful QH treatments at the general clinic divided by the number of successful patients, i.e. societal costs for 23 treatments divided by 23 patients.

- (c) the mean societal costs for all successful EP treatments at the specialist orthodontic clinic divided by the number of successful patients, i.e. societal costs for 17 treatments divided by 17 patients.
 - (d) the mean societal costs for all successful EP treatments at the general clinic divided by the number of successful patients, i.e. societal costs for 18 treatments divided by 18 patients.
2. The mean societal costs of the successful and retreated cases (cases that had to be retreated since no correction was achieved during 1 year) on completion of treatment in all groups:
- (a) the mean societal costs for all successful treatments and retreatments of QH at the specialist orthodontic clinic divided by the number of successful patients, i.e. 28 treatments divided by 28 patients, thus no retreatments and unaltered costs.
 - (b) the mean societal costs for all successful treatments and retreatments of QH at the general clinic divided by the number of successful patients, i.e. 27 treatments divided by 23 patients, thus 4 retreatments and an increase in costs.
 - (c) the mean societal costs for all successful treatments and retreatments of EP at the specialist orthodontic clinic divided by the number of successful patients, i.e. 27 treatments divided by 17 patients, thus 10 retreatments and an increase in costs.
 - (d) the mean societal costs for all successful treatments and retreatments of EP at the general clinic divided by the number of successful patients, i.e. 28 treatments divided by 18 patients, thus 10 retreatments and an increase in costs.

Statistical analysis

The sample size for each group to detect a relevant cost difference of 1000 SEK/€96 [standard deviation (SD): 1000/960] and a power of 90 per cent was calculated for 21 patients in each group. To compensate for possible dropouts of approximately 20 per cent per group, the number of patients per group was increased to at least 25.

The data were statistically analysed using SPSS software (version 24, SPSS, Chicago, Illinois). The chi-square and Fisher's exact test were used to calculate the success rate. Numerical variables, arithmetic means, and standard deviations were calculated. Analysis of variance (ANOVA) with the Tukey *post hoc* test was used to compare the costs and treatment time within and between groups. Differences with probabilities of less than 5 per cent ($P < 0.05$) were considered statistically significant.

Results

A total of 110 patients, 59 girls and 51 boys (mean age: 9.2 years; SD: 1.02), entered the study and were randomized into four groups. Accordingly, QHS included 11 boys and 17 girls (mean age: 9.3; SD: 1.14), QHG involved 13 boys and 14 girls (mean age: 9.6; SD: 1.05), EPS comprised 14 boys and 13 girls (mean age: 8.8; SD: 0.85), while EPG consisted of 13 boys and 15 girls (mean age: 9.0; SD: 0.87). Figure 3 illustrates the flow of patients throughout the trial. No significant differences in age and gender distribution were found between the groups at baseline.

Successful treatments

In the QHS group, 28 out of 28 patients were successfully corrected compared to 23 out of 27 in the QHG group. Treatment with expansion plate was less successful: 18 out of 27 in the EPS group and 18 out of 28 in the EPG group (Table 1). Thus, QHS were significantly

more successful compared with EPS and EPG, $P = 0.001$ and $P = 0.000$, respectively. No other differences in crossbite corrections between groups were seen.

Considering mandibular midline correction, significant differences were also found between the QHS and QHG groups ($P = 0.02$) and the QHS and EPG groups ($P = 0.007$) (Table 1).

Number of appointments

The mean number of appointments to the clinic of the successful treatments including the retention period of 3 months was 7.1 (SD: 2.63) appointments in the QHS group, 8.4 (SD: 2.21) in the QHG group, 8.2 (SD: 2.52) in the EPS group, and 9.2 (SD: 1.62) appointments in the EPG group. Significantly fewer numbers of appointments were found in the QHS compared with the EPG group ($P = 0.013$). The mean numbers of appointments for both the successful treatments and retreatments were 8.1 (SD: 3.97) for the QHS group, 10.9 (SD: 3.54) for the QHG group, 12.7 (SD: 6.13) for the EPS group, and 14.7 (SD: 6.32) appointments for the EPG group. Consequently, the QHS group had significantly fewer appointments compared with the EPS ($P = 0.007$) and EPG groups ($P = 0.000$) as well as infrequent number of appointments in the QHG compared with the EPG group ($P = 0.032$).

Complications

No appliances were lost in the two QH groups. However, 33 new EP had to be produced due to lost or broken plates, i.e. 13 in the EPS and 20 in the EPG group, which increased the material costs for the patients in these groups and especially in the EPG group. (Table 2).

Lost separation ligatures and ligatures to secure the QH arch wire into the molar bands were the most frequent reasons for emergency visits in the QH groups (7 in the QHS group and 14 in the QHG group).

No significant differences in emergency visits were detected between the groups; hence, the mean number of emergency visits was 1.0 (SD: 1.56) in the QHS group, 1.9 (SD: 1.24) in the QHG group, 1.29 (SD: 1.50) in the EPS, and 1.54 (SD: 1.26) in EPG group.

Cost-minimization analysis

Societal costs—successful treatments

The mean societal, direct and indirect costs per patient are shown in Table 2. For the patients with successful treatments, the societal costs were €765 for the QHS patients, €1025 for the QHG patients, €844 for the EPS patients, and €878 for the EPG patients. The cost per patient was significantly lower for the QHS compared with the QHG ($P = 0.000$). Moreover, costs for expansion plates in specialist orthodontic clinics were significantly lower compared with the QHG ($P = 0.015$) (Table 2).

Material costs—successful treatments

The mean costs were €120 for the QHS and QHG patients, €185 for the EPS patients, and €229 for the EPG patients. The mean material cost per patient was significantly lower for QH compared to expansion plate treatments in special as well as in general dentistry ($P = 0.000$). In addition, the material costs of expansion plate treatments in specialist orthodontic clinics were lower than plate treatments in general dentistry ($P = 0.013$).

Treatment time costs—successful treatments

The mean costs were €601 for the QHS patients, €788 for the QHG patients, €575 for the EPS patients, and €569 for the EPG

Figure 3. Flow diagram of patients in this trial.

patients. Thus, the QHG costs were significantly higher compared to both QH and expansion plate treatments in specialist orthodontic clinics ($P = 0.000$). Furthermore, in general dentistry the QH treatment time costs were significantly higher than plate treatments ($P = 0.000$).

Indirect costs—successful treatments

The costs were €44 for the QHS patients, €117 for the QHG patients, €83 for the EPS patients, and €80 for the EPG patients (Table 2). Overall, the indirect costs per patient were 5 to 10 per cent of the societal costs. The indirect cost per patient was significantly lower for the QHS compared with QHG ($P = 0.008$).

Costs including retreatments

In general, when the retreatments were included, the costs on all levels were increased for expansion plate treatments performed in general and special dentistry as well as for QHG (Table 2). The lower success rate for the expansion plate treatments in general and special dentistry and, additionally, for QHG was the main reason for the increased or added costs when retreatments were also included.

Finally, according to cost-minimization and since QH treatments accomplished in specialist orthodontic clinics revealed a 100 per cent

success rate (no retreatments), it was obvious that QHS were the most cost effective choice (Table 2). In the QHG group, 4 retreatments have to be performed and 10 retreatments in each of the EPS and EPG group (Table 2).

Discussion

Main findings

Economic analyses of orthodontic care are seldom presented but preserve essential data for the outlining and administration of orthodontic services. This study is the first to assess the costs of crossbite treatments performed in either specialist or general dentistry. It was clearly found that QH treatment performed in specialist orthodontic clinics had lower costs than QH or EP treatments accomplished in general dentistry as well as EPS. Consequently, the results rejected our hypotheses that the crossbite treatment methods are equal regarding costs and that treatment in general and specialist dentistry would be equally effective. In this evaluation we used the ITT (14), i.e. all cases successful or not were included, which means that in the cost-minimization analysis we also calculated the costs for retreatments of unsuccessful cases. Hence, the costs on all levels were increased for expansion plate treatments performed in general and special dentistry as well as for QHG (Table 2). Thus, the low cost of

Table 1. Success rate of crossbite and midline correction for patients treated with quad-helix in specialist orthodontist clinics (QHS) and general dentistry (QHG) and for patients treated with removable expansion plate in specialist orthodontist clinics (EPS) and in general dentistry (EPG)

Outcome	QHS <i>n</i> = 28	QHG <i>n</i> = 27	EPS <i>n</i> = 27	EPG <i>n</i> = 28	Group difference ^a
	<i>n</i>	<i>N</i>	<i>n</i>	<i>n</i>	<i>P</i>
Successful crossbite correction	28	23	17	18	QHS/QHG <i>P</i> = 0.051
Unsuccessful crossbite correction	0	4	10	10	QHS/EPS <i>P</i> = 0.001
					QHS/EPG <i>P</i> = 0.000
					QHG/EPS <i>P</i> = 0.111
					QHG/EPG <i>P</i> = 0.121
					EPS/EPG <i>P</i> = 1.000
Successful midline correction	24	10	16	10	QHS/QHG <i>P</i> = 0.020
Unsuccessful midline correction	4	10	8	10	QHS/EPS <i>P</i> = 0.104
					QHS/EPG <i>P</i> = 0.007
					QHG/EPS <i>P</i> = 0.474
					QHG/EPG <i>P</i> = 0.669
					EPS/EPG <i>P</i> = 0.263

^aChi-square test

treatments of unilateral posterior crossbite in specialist orthodontic clinics was underpinned.

One reason for the lower costs of QH treatment was that this treatment is less dependent on patient cooperation compared with expansion plate treatment. The lower costs of treatments in specialist orthodontic clinics can be explained by trained specialists in orthodontics being employed in specialist clinics and that these orthodontists have greater experience working with fixed appliances compared with general dentists who may be more familiar with using removable appliances. Furthermore, there is some variety among the general dentists in clinical orthodontic skills.

It is notable that the differences in costs were primarily related to treatment time and material costs (direct costs), and no statistically significant differences between the groups were found for indirect costs. Additionally, indirect costs represented 5 to 10 per cent of the societal costs regardless of whether treatment was performed in general or specialist dentistry. Comparable cost studies on treatment of unilateral posterior (15) and anterior crossbite (16) in the mixed dentition have shown the proportion of indirect costs amounting to between 35 and 45 per cent. Nonetheless, higher indirect costs regarding treatments performed at the specialist orthodontic clinics could be expected since those clinics are not so frequently distributed over the country as the general clinics and, therefore, the travelling route would be longer and more time consuming for many patients and parents. However, it seems that the indirect costs from this aspect described above were balanced by the lower number of appointments/visits for the patients at the specialist orthodontic clinics. In fact, this trial was conducted in a mixed rural and urban area where the overall distances for patients and parents to the general and specialist clinics were relatively short and thereby less time consuming and resulting in lower indirect costs.

Interpretations

The results of this study are in accordance with some previous studies in this field. Pietilä *et al.* (17) found that general dentists with little orthodontic experience were associated with the highest costs for orthodontic treatments. Furthermore, Godoy *et al.* (18) showed an equal success rate for QH and EP but concluded that treatment with QH revealed significantly shorter treatment time and lower

costs compared with expansion plate. In addition, Petréň *et al.* (15) presented in general dentistry a higher success rate and lower costs with QH compared with expansion plate treatments.

Drummond *et al.* (6) have claimed that it is important to calculate the opportunity cost for different treatments. From our trial, it can be shown that 10 patients treated with QH in general dentistry (successfully only) represent on average the same level of costs or resources (€10 250) as 13 patients (€10 250 divided by €765) treated in specialist orthodontic clinics. Moreover, if 10 patients are treated with EP in specialist orthodontic clinics (successfully only), it will result in a mean cost of €8440 and this cost will cover 11 patients treated with QH in specialist orthodontic clinics as well. This means, if QH treatment in general dentistry is used for standard cases this will reduce the number of conducted treatments by 3 out of 13 cases (23 per cent).

The mean cost differences in the present study were clearly significant, and extrapolation of the data to a national level further highlights the differences between the economics of the two treatment modalities. In the year 2017, there were 120 977 9-year-old children living in Sweden and the prevalence of posterior crossbite is estimated to be around 10 per cent. Treating all these children for posterior crossbite, the differences in societal costs between QHS and expansion plate treatments performed in general dentistry would be approximately €3.7 million.

Generalizability

When this multicentre RCT was undertaken, attempts were made to reduce the risk of bias and confounding factors. The prospective design and protocol ensured that the baseline characteristics and treatment progress were easily observed. Furthermore, we used a treatment protocol as close as possible to the normal treatment procedure to ensure that the results could be implemented in everyday work.

Moreover, the multicentre design resulted in treatment outcomes from a large number of operators, and this made the patient-related outcome closer to reality compared with if only a few operators were involved in the study. Additionally, the ITT approach was used to evaluate every patient record, regardless of whether the treatment was successful or not. Using ITT instead of a per-protocol approach will avoid overoptimistic estimates of efficacy of an intervention resulting from removal of not successfully treated subjects.

Table 2. Mean societal, direct and indirect costs (Euro) for the quad-helix treatment in specialist orthodontic clinics (QHS), quad-helix in general dentistry (QH), removable expansion plate treatment in specialist orthodontic clinics (EPS), and removable expansion plate treatment in general dentistry (EPG). The mean total costs per patient for successful treatments only and including retreatment. There were no retreatments in the QHS group, while there were 4 in the QHG and 10 each for the EPS and EPG groups, thus explaining the increased costs in the QHG, EPS, and EPG groups for including retreatments. CI, confidence interval; SD, standard deviation

	QHS			QH			EPS			EPG			Difference between groups ^a P
	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI	Mean	SD	95% CI	
Successful treatments	n = 28			n = 23			n = 17			n = 18			
Societal costs	765	182	695–836	1025	291	910–1140	844	215	759–929	878	163	814–940	QHG > QHS; P = 0.000 QHG > EPS; P = 0.015
Direct costs													
Material costs	120	0	120–120	120	0	120–120	185	68	159–212	229	78	199–259	EPS > QHS; EPG > QHS; EPS > QHG; EPG > QHG; P = 0.000
Treatment time costs													EPG > EPS; P = 0.013
Indirect costs	44	51	24–64	117	93	80–154	83	90	48–119	569	78	539–600	QHG > QHS; QHG > EPS; QHG > EPG; P = 0.000
Including retreatment	n = 28			n = 27			n = 27			n = 28			QHG > QHS; P = 0.008
Societal costs	765	182	695–836	1139	402	980–1298	1099	426	931–1268	1151	407	993–1309	QHG > QHS; P = 0.002 EPS > QHS; P = 0.006 EPG > QHS; P = 0.001
Direct costs													
Material costs	120	0	120–120	138	44	121–155	226	89	190–261	272	98	234–310	EPS > QHS; EPG > QHS; EPS > QHG; EPG > QHG; P = 0.000
Treatment time costs													
Indirect costs	601	148	544–659	877	341	742–1012	775	318	650–901	784	304	666–902	QHG > QHS; P = 0.003
Material costs	44	51	24–64	124	94	86–161	98	93	61–135	95	92	60–131	QHG > QHS; P = 0.004

^aOne-way analysis of variance with Tukey's *post hoc* test.

Limitations

It is important to point out that monetary variables are influenced by local factors such as insurance systems, staff salaries, rental costs, taxes, urban versus rural areas, etc., and consequently, the figures shown in this trial could not be directly extrapolated to other locations. Thus, the generalizability of the findings to other settings may be limited as the study was carried out on a regional scale, and furthermore, it may not be legally appropriate for general dentists in all countries to treat orthodontic patients. Nonetheless, it must be admitted that the success rate was very high (100 per cent) when unilateral crossbite was treated with QH in specialist orthodontic clinics. Even if diverse countries and health providers use different health economic systems of oral healthcare, the cost differences between the specialist and general dentistry found in this trial may still be of interest. Consequently, for any health provider or clinician, the results of this study could be valuable when making decisions on whether general or specialist dentistry instances should be utilized or which appliance is used.

Since this trial had a relatively short observation period (within 1 year), it must be acknowledged that the trial runs the risk of the Hawthorne effect (positive bias) whereby patients tend to perform better as participants in the trial (19). Of course, the Hawthorne effect induced all the four groups in this trial, but probably patients in the two groups treated with expansion plates could have been more compliant than the average patients in ordinary orthodontic practice. This implies that the success rate found in the EPS and EPG groups will conceivably be lower in everyday practice.

Conclusion

Based on our findings, we recommend that treatment of unilateral posterior crossbite in the mixed dentition shall be performed by a specialist in orthodontics using the QH appliance.

Funding

European Orthodontic Society Research Grant (2015); Region Halland, Sweden.

Acknowledgements

The authors would like to thank all patients and their parents for contributing to this study. We also want to thank the staff at the general and specialist dentistry clinics for their help in completing this study.

Conflict of interest

None to declare.

References

1. Thilander, B. and Myrberg, N. (1973) The prevalence of malocclusion in Swedish schoolchildren. *Scandinavian Journal of Dental Research*, 81, 12–21.
2. Heikinheimo, K. (1978) Need of orthodontic treatment in 7-year-old Finnish children. *Community Dentistry and Oral Epidemiology*, 6, 129–134.
3. Andrade, A.S., Gameiro, G.H., Derossi, M. and Gavião, M.B. (2009) Posterior crossbite and functional changes. A systematic review. *The Angle Orthodontist*, 79, 380–386.
4. Petré, S., Bondemark, L. and Söderfeldt, B. (2003) A systematic review concerning early orthodontic treatment of unilateral posterior crossbite. *The Angle Orthodontist*, 73, 588–596.
5. Kumar, S., Williams, A.C. and Sandy, J.R. (2006) How do we evaluate the economics of health care? *European Journal of Orthodontics*, 28, 513–519.
6. Drummond, M.F., Sculpher, M.J., Claxton, K., Stoddart, G.L. and Torrance, G.W. (2015) *Methods for the Economic Evaluation of Healthcare Programmes*. Oxford University Press, Oxford, UK, 4th edn.
7. Tickle, M. (1997) Clinical effectiveness and primary dental care. 2. The influence of health economics. *Primary Dental Care*, 4, 85–87.
8. Buck, D. (2000) Economic evaluation and dentistry. *Dental Update*, 27, 66–73.
9. Sollenius, O., Petré, S., Björnsön, L., Norlund, A. and Bondemark, L. (2016) Health economic evaluations in orthodontics: a systematic review. *European Journal of Orthodontics*, 38, 259–265.
10. Järvinen, S. and Widström, E. (2002) Determinants of costs of orthodontic treatment in the Finnish public health service. *Swedish Dental Journal*, 26, 41–49.
11. Deans, J., Playle, R., Durning, P. and Richmond, S. (2009) An exploratory study of the cost-effectiveness of orthodontic care in seven European countries. *European Journal of Orthodontics*, 31, 90–94.
12. Pietilä, T., Pietilä, I., Widström, E., Varrela, J. and Alanen, P. (1997) Extent and provision of orthodontic services for children and adolescents in Finland. *Community Dentistry and Oral Epidemiology*, 25, 150–155.
13. Petré, S., Bjerklin, K., Hedén, P. and Ecorcheville, A. (2014) Orthodontic treatment by general practitioners in consultation with orthodontists—a survey of appliances recommended by Swedish orthodontists. *Swedish Dental Journal*, 38, 121–132.
14. Bondemark, L. and Abdullaheem, S. (2018) Intention to treat (ITT) analysis as reported in orthodontic randomized controlled trials—evaluations of methodology and recommendations for the accurate use of ITT analysis and handling dropouts. *European Journal of Orthodontics*, 40, 409–413.
15. Petré, S., Bjerklin, K., Marké, L.Å. and Bondemark, L. (2013) Early correction of posterior crossbite—a cost-minimization analysis. *European Journal of Orthodontics*, 35, 14–21.
16. Wiedel, A.P., Norlund, A., Petré, S. and Bondemark, L. (2016) A cost minimization analysis of early correction of anterior crossbite—a randomized controlled trial. *European Journal of Orthodontics*, 38, 140–145.
17. Pietilä, T., Sintonen, H., Pietilä, I., Widström, E., Varrela, J. and Alanen, P. (1998) Cost and productivity analysis of orthodontic care in Finland. *Community Dentistry and Oral Epidemiology*, 26, 283–288.
18. Godoy, F., Godoy-Bezerra, J. and Rosenblatt, A. (2011) Treatment of posterior crossbite comparing 2 appliances: a community-based trial. *American Journal of Orthodontics and Dentofacial Orthopedics*, 139, e45–e52.
19. Abdullaheem, S. and Bondemark, L. (2018) Hawthorne effect reporting in orthodontic randomized controlled trials: truth or myth? Blessing or curse? *European Journal of Orthodontics*, 40, 475–479.

IV

Randomized controlled trial

Three-dimensional evaluation of forced unilateral posterior crossbite correction in the mixed dentition: a randomized controlled trial

Ola Sollenius¹, Aljaž Golež², Jasmina Primožič², Maja Ovsenik²,
Lars Bondemark³ and Sofia Petrén³

¹Department of Orthodontics, County Council, Halland, Halmstad, Sweden, ²Department of Orthodontics and Dentofacial Orthopaedics, Faculty of Medicine, University of Ljubljana, Slovenia, and ³Department of Orthodontics, Malmö University, Malmö, Sweden

Correspondence to: Ola Sollenius, Department of Orthodontics, Halland County Council, S-30246 Halmstad, Sweden. E-mail: ola.sollenius@regionhalland.se

Summary

Objectives: The objectives of this study were to assess the three-dimensional (3D) treatment changes (palatal surface area and volume) of forced unilateral posterior crossbite correction using either quad-helix or removable expansion plate appliances in the mixed dentition, and to compare the treatment changes with the three-dimensional changes occurring in age-matched untreated unilateral posterior crossbite patients as well as in subjects with normal occlusion and with no or mild orthodontic treatment need.

Trial design: Six-arm parallel group multicentre randomized controlled trial.

Materials and methods: One-hundred and thirty-five patients with unilateral posterior crossbite with functional shift were recruited. The patients were randomized by an independent person not involved in the trial. The randomization used blocks of 25, and the patients were randomized into the following five groups: quad-helix treatments in specialist orthodontic clinics (QHS), quad-helix treatments in general dentistry (QHG), removable expansion plate treatments in specialist orthodontic clinics (EPS), removable expansion plate treatments in general dentistry (EPG), and untreated crossbite (UC). Twenty-five patients with normal occlusion who served as normal controls were also included in the trial. Blinding of the outcome assessor and data analyst was accomplished. Data on all children were evaluated on an intention-to-treat basis, regarding 3D palatal surface area, palatal projection area, and palatal shell volume; two-dimensional linear measurements were registered at the same time.

Results: After treatment, the surface and projection area and shell volume increased in the four treatment groups (QHS, QHG, EPS, and EPG). QHS increased significantly more than EPG for the surface and projection area. The QHS and EPS had significantly higher mean difference for shell volume.

Limitations: The trial considers a short-term evaluation.

Conclusion: After treatment, there were no significant differences between the four treatment groups and the normal group, which implies that the surface and projection area together with the shell volume for the four treatment groups and the normal group were equivalent.

Trial registration: The trial was registered with <https://www.researchweb.org/is/sverige>, registration number: 220751.

Introduction

Unilateral posterior crossbite (UPC) with functional shift is one of the most common malocclusions in the mixed dentition, with a reported prevalence ranging from 8.5 to 17% (1, 2). Left untreated, it may cause long-term effects on the growth and development of the jaws and teeth (3, 4). Several studies on treatment outcomes of crossbite correction have been published (5–8). Correction of UPC may be carried out in different dental stages; for example, Primožič *et al.* (7) advocate treatment in the primary dentition whereas Petřén and Bondemark (8) in the mixed dentition.

Maxillary expansion is usually assessed regarding correction of posterior crossbite. However, in growing subjects, the assessment should also involve the reestablishment of normal growth and occlusal development. Hence, as treatment of UPC is performed in growing children, it is also advisable to relate the treatment changes to growth changes associated in untreated subjects with UPC as well as in subjects with normal occlusion and with no or mild orthodontic treatment need.

Treatment of posterior crossbite is usually performed by a specialist in orthodontics but in some countries, general dentists under supervision of a specialist in orthodontics carry out these treatments. A recent study considering cost-effectiveness (9) has presented that posterior crossbite in the mixed dentition should be treated with quad-helix (QH) and by specialists in orthodontics. In this study, a further evaluation regarding the three-dimensional (3D) and two-dimensional (2D) morphological changes in treated crossbite patients were compared to untreated posterior crossbite children and children with normal occlusion.

When the treatment effects of unilateral crossbite correction have been assessed, most studies have used 2D evaluations such as linear measurements, for example, the amount of expansion in millimetres between the permanent first molars and between the primary canines (8, 10). However, orthodontic treatment effects including crossbite correction may, if possible, be described in all three planes. In 2018, Leonardi *et al.* (11) published a study on 3D evaluation of crossbite patients. They found a bilateral symmetrical contraction of the palatal vault and an asymmetric contraction of the alveolar process in patients with UPC. Nevertheless, 3D evaluations of the maxilla using 3D technique have been performed before (6, 7). In the study by Primožič *et al.* (7), they showed that correction of UPC by an expansion plate in the primary dentition resulted in an increase of palatal surface area and palatal volume.

Several studies (12–15) have shown that intraoral scanning presents good reliability of the scanned structures when compared with conventional analogic measures on study casts. Also, the 3D images from intraoral scanning or from scanning on study casts have been proven to be valid and reliable (16), and therefore, may be used to measure, for instance, palatal surface area and palatal volume (7).

To the best of our knowledge, there are no studies in the literature that explicitly have investigated the 3D treatment effects of unilateral crossbite correction between different appliances and, in this context, related the treatment changes to growth changes associated with untreated subjects with UPC as well as in subjects with normal occlusion and with no or mild orthodontic treatment need. Therefore, the aims of this investigation were 1. to assess the 3D treatment changes (specifically palatal surface area and volume) of forced UPC correction using either QH or removable expansion plate appliances in the mixed dentition, and 2. to compare the treatment changes with the 3D changes occurring in age-matched untreated UPC patients as well as in subjects with normal occlusion and with no or mild orthodontic treatment need.

It was hypothesized that there are no significant differences regarding palatal surface area and volume between patients treated with QH and expansion plate. Furthermore, it was hypothesized that there were no significant morphological changes in the untreated control group. It was also hypothesized that the palatal surface area and volume in the treated groups would not reach the dimensions of the normal group.

Material and methods

The study sample comprised data from an randomized controlled trial (RCT) regarding correction of UPC children in the mixed dentition. Between 2013 and 2018, we consecutively recruited 135 children, all children who attended and met the inclusion criteria, from 10 general dental health clinics at the Public Dental Health Service, Halland County Council, Sweden. Data collection took place from December 2013 to November 2018.

The trial was registered in 'FoU i Sverige' (<https://www.researchweb.org/is/sverige>) registration number: 220751. The inclusion criteria were the following:

- Consecutive children, boys and girls
- Mixed dentition (between 8 and 10 years of age)
- UPC including the first permanent molar and with a functional shift of more than 1 mm
- Class I occlusion

The exclusion criteria were the following:

- Children with sucking habits or ceased sucking habits during the year before the trial was started
- Children who had previously undergone orthodontic treatment
- Children who had severe crowding of teeth (extraction of teeth necessary)
- Children with craniofacial syndromes

Before treatment started, two experienced orthodontists provided the patients and their parents with oral and written information about the trial, and the responsible general dentist got the same information at this visit.

Subjects were randomized into one of five groups: treatment in general dentistry clinics with quad-helix (QH) or removable expansion plate (EPG), or in specialist orthodontic clinics with quad-helix (QHS), or removable expansion plate (EPS) or in to an untreated control group (UC) where the treatment of posterior crossbite was postponed for 1 year.

Normal control group

Parallel to the recruitment of crossbite patients, 25 patients with normal occlusion were collected as normal controls. The reason for using a normal control group was to also evaluate the maxillary dimensions in subjects with normal occlusion, i.e. if the treated crossbite patients reached the same dimension values as the normal children. The inclusion criteria for the normal control group (NC) were children, boys and girls, in the mixed dentition (between 8 and 10 years of age), with normal occlusion, and with no or mild orthodontic treatment need, that is, Index of Orthodontic Treatment Need—Dental Health Component (17) of 1 or 2. The exclusion criteria were craniofacial syndromes, children who had previously undergone orthodontic treatment, or severe crowding of teeth (extraction of teeth necessary).

Ethical approval

The Regional Ethical Review Board in Lund, Sweden, approved the trial protocol and informed consent form (Dno: 2012/650).

Randomization procedure

The randomization was prepared and carried out by an independent person not involved in the study and the randomization used blocks of 25 (5 + 5 + 5 + 5 + 5). Thus, the orthodontists carried boxes of already prepared envelopes. Six opaque envelopes were prepared with 25 sealed notes in each (5 notes for each group), and for every new patient in the study, a note was extracted from the first envelope. When the envelope was empty, the second envelope was opened, and the 25 new notes were extracted as the patients were recruited to the trial. This procedure was then repeated five more times.

Sample size calculation

As Primožič *et al.* (18) are the only authors who have evaluated 3D palatal vault changes in assessing treatment of constricted maxilla in growing subjects, we used their sample size calculation as a base for clinical relevant changes.

With the assumption of means in variances for different groups taken from Primožič *et al.* (18), and using multiple testing adjustment suggested by Dunnett and Tamhane (19), with standard parameters of 80% power and α 0.05, a number of 23 children per group was needed, and as we expected some dropouts to occur, the required sample size was estimated to be 25 in each of the groups.

Treatment procedures

After recruitment and randomization, all children were treated at 3 orthodontic specialist clinics and 10 general dentistry clinics by two orthodontic specialists and 17 general dentists. The two orthodontists who provided the information to the patients had 15 years of experience and treated all the patients in the specialist orthodontic clinics and did not treat any patients at the general dental clinics. The 17 general dentists had between 1 and 30 years of experience of interceptive orthodontic treatments.

No specific calibration was performed, but the treatments followed a preset standard concept with specific descriptions of the design and management of the appliances; see The quad-helix and The removable expansion plate sections later.

The quad-helix

The QH consisted of a standard stainless-steel arch (MIA system; 3M Unitek, Monrovia, California, USA) with stainless-steel bands attached with glass ionomer cement on the maxillary first molars. The QH was activated 10 mm before placement and then reactivated every 6 weeks until a normal transverse

relationship was achieved (maximum treatment time 1 year) and no overcorrection was produced. Steel- or elastic ligatures were used to secure the QH to the bands. The treatment result was retained for 3 months.

The removable expansion plate

The removable expansion plate (EP) consisted of acrylic with an expansion screw, lined up with the midpalate suture, and stainless-steel clasps on the primary and permanent first molars. There was no acrylic covering the occlusal surfaces. The EP was activated 0.2 mm once a week by the patient until a normal transverse relationship was achieved (maximum treatment time 1 year) and no overcorrection was produced. The dentist thoroughly instructed the patient to use the EP day and night except for meals and tooth brushing. Treatment progression was evaluated every 6 weeks, and the result was retained for 3 months. Adjustment of steel clasps was performed when needed.

Outcome measures

The primary outcome measures to be assessed in the trial before and after active treatment or at baseline and after 1 year of expectance in the untreated or normal control groups were 1. palatal surfaces area changes, 2. palatal projection area changes, and 3. palatal shell volume (Figure 1). Further outcomes were the linear maxillary and mandibular intercanine and intermolar distances as well as the success rate of crossbite correction, which means that normal transverse relations would be achieved within 1 year.

Assessment of palatal surface area, projection area, and shell volume

Plaster casts before and after treatment were scanned with Trios 3 Intraoral Scanner (3Shape, Copenhagen, Denmark), the 3D data were imported to a reverse modelling software package Rapidform™ 2006 (INUS Technology, Seoul, South Korea), and the analysis was made as described by Primožič *et al.* (7, 20).

To measure the palatal surface area and to calculate the palatal volume, the boundaries/lines of the palate had to be defined. The gingival plane and a distal plane were used as boundaries for the palate. The gingival plane was created by connecting the midpoints of the dento-gingival junction of all primary teeth, permanent incisors, and first permanent molars (Figure 1). The distal plane was created through two points at the distal of the first permanent molar perpendicular to the gingival plane (Figure 1). The palatal surface area and the volume were then calculated. Also, intercanine and intermolar maxillary and mandibular transverse widths at the cusp and gingival levels were measured.

Figure 1. Illustrations of surface area (a), projection area (b), and surface volume (c) on three-dimensional scanmings.

Figure 2. Flowchart.

Intention-to-treat

Data on all children were evaluated on an intention-to-treat (ITT) basis; consequently, if the crossbite was not corrected during the trial period of 1 year, the outcome was judged as unsuccessful, and the possible change in palatal area and volume as well as linear expansion effect was quoted. Accordingly, all children, successful or not, were incorporated in the final analysis. Moreover, any withdrawals in the course of the trial were regarded as unsuccessful and with no expansion or changes. If data were lacking at the follow-up or after treatment, the last observation carried forward principle was used (21). This implies that the last measured values or data were imputed at the follow-up. After treatment and 3 months of retention, a second impression was taken 1 month after the removal of appliance. This was performed to enable blinding. For UC and NC, the second evaluation was performed after 1 year of observation.

Blinding

Children and trial personnel could not be blinded due to the character of treatment, but blinding was performed of the outcome evaluator and the person who analysed the data. Hence, the evaluator was unaware of the group to which the child had been allocated or whether the data were from baseline, follow-up, or after treatment.

Statistical analysis

The data were statistically analysed using SPSS software (version 24, SPSS, Chicago, Illinois, USA). Numerical variables, arithmetic means, and standard deviations (SDs) were calculated. Analysis of variance with the Tukey *post hoc* test was used to compare the palatal surfaces area, palatal projection area, and palatal shell volume changes as well as the linear maxillary and mandibular intercanine

and intermolar distance changes within and between groups. The chi-square and Fisher's exact test was used to calculate the success rate of crossbite correction. Differences with probabilities of less than 5% ($P < 0.05$) were considered statistically significant.

Method error analysis

Twenty randomly selected scanned models were measured at two separate times. No significant mean differences between the records were found using paired *t*-test, i.e. no systematic errors could be detected. The ICC and error of the measurements (in percentages) were calculated and were for palatal surface area less than 3.5 and 4.5 for palatal volume. For the linear measurements (in millimetres), the error according to Dahlberg's formula (22) was less than 0.17 mm or between 0.1 and 0.2 mm.

Results

A total of 160 children entered the study, and of these, 135 children with crossbite were randomized into 5 groups, and 25 children with normal occlusion constituted the normal group (Figure 2). In Table 1, the age and gender distribution in the groups are displayed. No significant differences were found between the groups, except that the age in the EPS and UC groups at baseline was significantly lower compared to the other groups ($P < 0.05$).

Palatal surface area, projection area, and shell volume

At baseline or before treatment, all five crossbite groups had significantly less palatal surface and projection area as well as palatal shell volume compared to the normal group (Table 2). No significant differences between the five crossbite groups were found regarding surface or projection area or shell volume (Table 2).

After treatment, the surface and projection area and shell volume increased in the four treatment groups (QHS, QHG, EPS, and EPG) (Table 3). However, in the untreated crossbite group and the normal occlusion group, the surface and projection area as well as the shell volume were more or less unaltered in size after 1 year. Nevertheless, no significant differences were found between the QHS, QHG, and EPS groups, but these groups revealed significantly larger surface and projection area compared to the untreated crossbite group and the normal group (Table 3). Moreover, the surface and projection area increased significantly more in the QHS than in the EPG group. The QHS and EPS groups also showed significantly higher mean differences for the

palatal shell volume (Table 3). Also, the surface areas were significantly larger in the QHS and EPS groups compared to the untreated control group whereas the projection area and shell volume were significantly larger in the EPS than in the untreated control group (Table 3).

After treatment, the four treatment groups reached the normal group values considering the surface and projection area together with the shell volume (Table 2).

No within-group changes were found for the untreated and normal groups during the 1-year follow-up.

In the mandible, small or negligible and insignificant changes within and between the groups were noted considering surface and projection area as well as shell volume.

Linear measurements

At baseline, there were significantly smaller maxillary intermolar and intercanine distances except for distances between the intercanine cusp tips when the five crossbite groups were compared to the normal group; however, no significant differences were found between the crossbite groups (Table 4).

After treatment, the intermolar and intercanine distances in the maxilla significantly increase within all four treatment groups (Table 4). The increase was mostly pronounced in the QHS and QHG groups, which exceeded some of the values of the normal group (Table 4).

Between the groups, the maxillary intermolar and intercanine expansion was significantly larger in the QHS than in the QHG, EPS, and EPG groups whereas all four treatment groups showed greater expansion than in the untreated control and normal group especially on the molars (Table 5).

Small or negligible and insignificant changes were found in the mandible regarding the intermolar and intercanine distances.

Treatment time

The mean treatment time to correct the crossbite, including retention of 3 months, was 7.5 months (SD = 1.45, range 6.0–10.5) for QHS, 8.2 months (SD = 2.23, range 6.0–12.0) for QHG, 11.4 months (SD = 3.40, range 7.5–20.0) for EPS, and 12.0 months (SD = 3.63, range 6.0–21.0) for EPG.

Success rate of crossbite correction

In the QHS group, 28 of 28 children were successfully corrected whereas in the QHG group 23 of 27 children ($P = 0.051$) were successfully corrected. The expansion plate treatments were less successful, that is, 18 of 27 in the EPS group and 18 of 28 in the EPG group. Consequently, QH treatment in orthodontic specialist

Table 1. Baseline/pre-treatment demographic characteristics. QHS, quad-helix treatments in specialist orthodontist clinics; QHG, quad-helix treatments in general dentist clinics; EPS, removable expansion plate treatments in specialist orthodontist clinics; EPG, removable expansion plate treatments in general dental clinics; UC, untreated posterior crossbite control group; NC, normal control group with normal occlusion and no or mild orthodontic treatment need; SD, standard deviation

Baseline/pre-treatment						
Group	Number			Age (year)		
	Total	Boys	Girls	Mean	Range	SD
QHS	28	11	17	9.3	7.4–11.5	1.09
QHG	27	14	13	9.5	8.1–12.4	1.04
EPS	27	15	12	8.7	7.6–10.7	0.78
EPG	28	11	17	9.2	6.5–10.6	1.11
UC	25	16	9	8.5	7.7–11.1	0.57
NC	25	17	8	9.3	6.5–12.4	1.06

Table 2. Surface area, projection, area, and shell volume at baseline/pre-treatment and after treatment. QHS, quad-helix treatments in specialist orthodontist clinics; QHG, quad-helix treatments in general dentist clinics; EPS, removable expansion plate treatments in specialist orthodontist clinics; EPF, removable expansion plate treatments in general dentist clinics; UC, untreated posterior crossbite control group; NC, normal control group with normal occlusion and no or mild orthodontic treatment need. One-way analysis of variance with Tukey's *post hoc* test. SD, standard deviation; CI, confidence interval

	Group QHS			Group QHG			Group EPS			Group EPF			Group UC			Group NC			Difference
	Mean	SD	CI (%)	Mean	SD	CI (%)	Mean	SD	CI (%)	Mean	SD	CI (%)	Mean	SD	CI (%)	Mean	SD	CI (%)	
Baseline																			
Maxilla Surface area (mm ²)	1154	87.69	1120–1188	1205	122.60	1156–1253	1159	85.26	1126–1193	1154	68.73	1128–1181	1170	89.06	1133–1207	1292	123.19	1241–1343	QHS, QHG, EPS, EPF, UC < NC <i>P</i> < 0.02
Projection area (mm ²)	810	67.22	784–836	781	160.34	718–844	836	69.85	809–864	841	56.99	819–863	835	64.17	808–861	939	77.15	907–970	QHS, QHG, EPS, EPF, UC < NC <i>P</i> < 0.002
Shell volume (mm ³)	4823	639.53	4375–5071	5079	845.92	4744–5414	4887	638.19	4635–5140	4843	486.59	4655–5032	4815	711.57	4521–5108	5676	1013.52	5258–6095	QHS, QHG, EPS, EPF, UC < NC <i>P</i> < 0.05
Post-treatment																			
Maxilla Surface area (mm ²)	1261	95.51	1224–1298	1276	130.31	1224–1328	1275	95.14	1238–1313	1220	73.88	1191–1248	1184	91.07	1146–1221	1300	129.06	1247–1354	QHG > UC <i>P</i> = 0.02 EPS > UC <i>P</i> = 0.022 NC > UC <i>P</i> = 0.002
Projection area (mm ²)	922	74.52	894–951	870	182.76	797–942	933	90.52	897–969	894	59.87	871–917	843	63.13	817–869	940	79.16	907–972	EPS > UC <i>P</i> = 0.02 NC > UC <i>P</i> = 0.012
Shell volume (mm ³)	5423	649.70	5171–5675	5618	943.60	5245–5992	5758	785.97	5447–6069	5355	538.61	5146–5563	4976	833.90	4632–5320	5823	1119.68	5361–6286	EPS > UC <i>P</i> = 0.011 NC > UC <i>P</i> = 0.005

Table 3. Mean differences in surface area, projection area, and shell volume within and between the groups. QHS, quad-helix treatments in specialist orthodontist clinics; QHG, quad-helix treatments in general dentist clinics; EPS, removable expansion plate treatments in specialist orthodontist clinics; EPG, removable expansion plate treatments in general dentist clinics; UC, untreated posterior crossbite control group; NC, normal control group with normal occlusion and no or mild orthodontic treatment need. One-way analysis of variance with Tukey's *post hoc* test. SD, standard deviation; CI, confidence interval

	Group QHS			Group QHG			Group EPS			Group EPG			Group UC			Group NC			Difference
	Mean	SD	CI (%)	Mean	SD	CI (%)	Mean	SD	CI (%)	Mean	SD	CI (%)	Mean	SD	CI (%)	Mean	SD	CI (%)	
Maxilla																			
Surface area (mm ²)	107.4	52.94	86.9–127.9	69.7	61.67	45.3–94.1	91.4	71.01	63.3–119.5	52.6	57.30	30.4–74.8	13.6	54.96	–9.1 to 36.3	8.1	66.19	–19.2 to 35.5	QHS, QHG, EPS > UC and NC $P < 0.015$ QHS > EPG $P = 0.012$
Projection area (mm ²)	112.7	42.23	96.3–129.0	88.2	85.71	54.3–122.1	76.5	60.45	52.6–100.4	44.4	47.43	26.0–62.8	8.4	31.66	–4.6 to 21.5	0.7	45.65	–18.1 to 19.6	QHS, QHG, EPS > UC, and NC $P < 0.000$ QHS > EPG $P = 0.000$ QHG > EPG $P = 0.042$ EPS > EPG $P = 0.045$
Shell volume (mm ³)	599.9	344.76	466.2–733.5	440.1	384.79	287.9–592.3	697.3	547.25	480.8–913.8	404.7	425.92	239.6–569.9	161.2	402.86	–5.1 to 327.5	147.0	529.29	–71.5 to 365.3	QHS, EPS > UC, and NC $P < 0.06$

clinics was significantly more successful compared with expansion plate treatments in orthodontic specialist or general dentist clinics, ($P = 0.001$ and $P = 0.000$).

Harm

No harm was observed in any subjects of the included groups. However, a few parents of the patients in the untreated control group were concerned that their child has to wait for one year before the treatment started. However, all 25 untreated control children were later successfully treated with QH, and the patients and parents were satisfied.

Discussion

Main findings

The palatal surface and projection area increased the most in children treated with QH in orthodontic specialist clinics (QHS), but a pronounced increase was also found in general dentistry (QHG). Also, children treated with removable expansion plates in specialist orthodontic clinics (EPS) revealed adequate treatment effects. Considering the maxillary shell volume, this variable increased mostly in the QHS and EPS groups. Another important finding was that, after treatment, all four treated groups reached the same palatal surface area and volume as the normal group. Thus, our initial hypothesis that there were no significant differences regarding palatal surface area and volume between patients treated with QH or EP was partially rejected, but the hypothesis that there were no significant morphological changes in the untreated control group was supported. Finally, the hypothesis that the palatal surface area and volume after treatment in the treated groups would not reach the dimensions of the normal group was rejected.

It can also be pointed out that the lower increase in expansion found in the expansion plate group compared to the QH group can be explained by a higher success rate of QH group treatments which reached 100% in the QHS group.

Interpretations

To our knowledge, this is the first RCT that has used and analysed the 3D changes after posterior crossbite correction in the mixed dentition and simultaneously compared the treatment results with those changes that occurred in two control groups; one untreated crossbite control group and another control group with normal occlusion. Hence, a search in the literature has failed to identify any comparable studies in the mixed dentition, but similar studies have been performed in the primary dentition by Primožič *et al.* (23, 24) that stated that after treatment there were no differences between treated crossbites and noncrossbite patients.

Both the 3D and 2D evaluations in this study revealed that the maxillary variables of the crossbite children had smaller values compared to children with normal occlusion at baseline. After treatment with QH or EP, the maxillary expansion of the crossbite patients significantly improved, and the 3D as well as the 2D evaluations showed that the maxillary parameters reached the size of the parameters found in the normal group with normal occlusion. These post-treatment results contrasted with the treatment outcomes presented by Petrén and Bondemark (8). In that 2D study, the expanded maxilla never reached the same level of width as the normal controls.

Concerning the growth changes during the one-year trial, it could be concluded that small 3D and 2D changes took place. For

Table 4. Intermolar and intercanine distances at baseline/pre-treatment and after treatment. QHS, quad-helix treatments in general dentist clinics; EPS, removable expansion plate treatments in specialist orthodontist clinics; QHG, quad-helix treatments in general dentist clinics; EPB, removable expansion plate treatments in general dental clinics; UC, untreated posterior crossbite control group; NC, normal control group with normal occlusion and no or mild orthodontic treatment need. One-way analysis of variance with Tukey's post hoc test. SD, standard deviation; CI, confidence interval

[illegible]

Table 5. Mean differences in intermolar and intercanine distances within and between the groups. QHS, quad-helix treatments in specialist orthodontist clinics; QHG, quad-helix treatments in general dentist clinics; EPS, removable expansion plate treatments in specialist orthodontist clinics; EPG, removable expansion plate treatments in general dental clinics; UC, untreated posterior crossbite control group; NC, normal control group with normal occlusion and no or mild orthodontic treatment need. One-way analysis of variance with Tukey's *post hoc* test. SD, standard deviation; CI, confidence interval

	Group QHS			Group QHG			Group EPS			Group EPG			Group UC			Group NC			Difference
	Mean	SD	CI%	Mean	SD	CI%	Mean	SD	CI%	Mean	SD	CI%	Mean	SD	CI%	Mean	SD	CI%	
Maxilla																			
Intermolar distance, mesio-buccal cusp tips	5.2	1.61	4.62–5.86	3.8	2.20	2.90–4.65	3.3	1.63	2.67–3.96	2.6	2.21	1.59–3.50	0.5	0.36	0.34–0.65	0.2	0.67	–0.07 to 0.49	QHS > QHG <i>P</i> = 0.013, QHS > EPS, EPG, UC and NC <i>P</i> = 0.000. QHG > UC and NC <i>P</i> = 0.000 EPS > UC and NC <i>P</i> = 0.000. EPG > UC and NC <i>P</i> = 0.000
Intermolar distance, gingival margin	4.9	1.47	4.32–5.46	3.3	1.90	2.59–4.09	2.9	1.46	2.36–3.51	2.1	1.80	1.35–2.91	0.3	0.37	0.16–0.48	0.2	0.57	–0.04 to 0.43	QHS > QHG <i>P</i> = 0.01 QHS > EPS, EPG, UC, NC <i>P</i> = 0.000 QHG > EPG <i>P</i> = 0.034 QHG > UC and NC <i>P</i> = 0.000 EPGUC and NC <i>P</i> = 0.000 EPGUC and NG <i>P</i> = 0.000
Intercanine distance, cusp tips	4.4	1.38	3.87–5.01	2.3	1.73	1.52–3.06	3.3	1.46	2.71–3.92	1.8	1.72	0.93–2.60	0.3	0.77	–0.03 to 0.68	0.6	0.90	0.06 to 1.20	QHS > QHG, EPG, UC and NC <i>P</i> = 0.000 QHG > UC and QHG > NC <i>P</i> = 0.017 EPS > EPG <i>P</i> = 0.006 EPS > UC and NC <i>P</i> = 0.000 EPG > UC <i>P</i> = 0.019
Intercanine distance, gingival margin	3.3	1.23	2.74–3.76	1.7	1.21	1.16–2.23	2.5	1.29	2.01–3.07	1.3	1.42	0.65–2.02	0.4	0.56	0.13–0.64	0.5	0.81	–0.02 to 1.02	QHS > QHG, EPG, UC, NC <i>P</i> = 0.000 QHG > UC <i>P</i> = 0.004 EPS > EPG <i>P</i> = 0.011 EPS > UC and NC <i>P</i> = 0.000

instance, between 0.3 and 0.6 mm of the increased palatal width corresponded to natural growth in the untreated and normal control groups (Table 5)

Strengths

One of the strengths of this trial was that there were few dropouts and we used a multicenter RCT with an ITT approach. Thus, all children, successfully treated or not, were included, which minimized the risk of false positive treatment results. Moreover, measurement errors were judged to be small and clinically irrelevant.

Blinding of the trial staff and the treated children could not be performed because of the character of the treatments but blinding of the outcome evaluator was performed. Consequently, the evaluator did not know to which group the child belonged or whether the data were at baseline/before or after intervention or at the one-year follow-up.

Limitations

The strength of this trial might have been further enhanced if we also had a follow-up period after treatment to evaluate the stability of the treatments performed. Nonetheless, the primary objectives of this trial were to assess the short-term 3D treatment changes of forced UPC correction using either QH or removable expansion plate in the mixed dentition and then to compare the treatment changes with the changes occurring in age-matched untreated UPC patients as well as in subjects with normal occlusion. Since it is important also to evaluate the post-treatment 3D changes from a long-term perspective, the treated children will be followed for at least a further two-year period.

Generalizability

A treatment protocol as near as possible to the established treatment procedures was applied and all clinics and therapists were instructed to follow this pre-set protocol as close as possible, and by that, the results could be used in the regular clinical work for crossbite treatment in the mixed dentition. In addition, since many therapists were engaged in this trial, this made the patient-related results nearer to reality compared to if infrequent therapists were included in the trial. In addition, both the prospective trial design and protocol assured that the baseline characteristics and treatment progress were readily noted.

Conclusions

- Children with UPC with functional shift in the mixed dentition revealed before treatment significantly smaller palatal surface and projection area as well as smaller palatal shell volume than normal control children.
- UPC correction resulted in a normalization of the palatal structures in terms of palatal area and volume.
- QH treatments in orthodontic specialist clinics but also in general dentistry as well as removable expansion plate treatments in specialist orthodontic clinics revealed a significant increase of palatal area and volume.
- The natural growth changes during the one-year trial were small.

Funding

This study was supported by the European Orthodontic Society Research grant (2015), Region Halland, Sweden.

Conflict of interest

None declared.

References

1. Thilander, B. and Myrberg, N. (1973) The prevalence of malocclusion in Swedish schoolchildren. *Scandinavian Journal of Dental Research*, 81, 12–21.
2. Heikinheimo, K. (1978) Need of orthodontic treatment in 7-year-old Finnish children. *Community Dentistry and Oral Epidemiology*, 6, 129–134.
3. Andrade, A.D.A.S., Gameiro, G.H., Derossi, M. and Gavião, M.B. (2009) Posterior crossbite and functional changes. A systematic review. *The Angle Orthodontist*, 79, 380–386.
4. Iodice, G., Danzi, G., Cimino, R., Paduano, S. and Michelotti, A. (2016) Association between posterior crossbite, skeletal, and muscle asymmetry: a systematic review. *European Journal of Orthodontics*, 38, 638–651.
5. Boysen, B., La Cour, K., Athanasiou, A.E. and Gjessing, P.E. (1992) Three-dimensional evaluation of dentoalveolar changes after posterior crossbite correction by quad-helix or removable appliances. *British Journal of Orthodontics*, 19, 97–107.
6. Oliveira De Felipe, N.L., Da Silva, A.C., Viana, G., Kusnoto, B., Smith, B. and Evans, C.A. (2008) Relationship between rapid maxillary expansion and nasal cavity size and airway resistance: short- and long-term effects. *American Journal of Orthodontics and Dentofacial Orthopedics*, 134, 370–382.
7. Primožič, J., Richmond, S., Kau, C.H., Zhurov, A. and Ovsenik, M. (2013) Three-dimensional evaluation of early crossbite correction: a longitudinal study. *European Journal of Orthodontics*, 35, 7–13.
8. Petré, S. and Bondemark, L. (2008) Correction of unilateral posterior crossbite in the mixed dentition: a randomized controlled trial. *American Journal of Orthodontics and Dentofacial Orthopedics*, 133, 790.e7–790.13.
9. Sollenius, O., Petré, S. and Bondemark, L. (2019) An RCT on clinical effectiveness and cost analysis of correction of unilateral posterior crossbite with functional shift in specialist and general dentistry. *European Journal of Orthodontics*, Apr 11 pii, ciz014. doi:10.1093/ejoc/cjz014. [Epub ahead of print].
10. Thilander, B. and Lennartsson, B. (2002) A study of children with unilateral posterior crossbite, treated and untreated, in the deciduous dentition-occlusal and skeletal characteristics of significance in predicting the long-term outcome. *Journal of Orofacial Orthopedics*, 63, 371–383.
11. Leonardi, R., Lo Giudice, A., Rugeri, M., Muraglia, S., Cordasco, G. and Barbato, E. (2018) Three-dimensional evaluation on digital casts of maxillary palatal size and morphology in patients with functional posterior crossbite. *European Journal of Orthodontics*, 28, 556–562. doi:10.1093/ejoc/cjx103
12. Sfondrini, M.F., Gandini, P., Malfatto, M., Di Corato, F., Trovati, F. and Scribante, A. (2018) Computerized casts for orthodontic purpose using powder-free intraoral scanners: accuracy, execution time, and patient feedback. *BioMed Research International*, 2018, 4103232.
13. Nawi, N., Mohamed, A.M., Marizan Nor, M. and Ashar, N.A. (2018) Correlation and agreement of a digital and conventional method to measure arch parameters. *Journal of Orofacial Orthopedics*, 79, 19–27.
14. Gül Amuk, N., Karsli, E. and Kurt, G. (2019) Comparison of dental measurements between conventional plaster models, digital models obtained by impression scanning and plaster model scanning. *International Orthodontics*, 17, 151–158.
15. Wiranto, M.G., Engelbrecht, W.P., Tutein Nolthenius, H.E., van der Meer, W.J. and Ren, Y. (2013) Validity, reliability, and reproducibility of linear measurements on digital models obtained from intraoral and cone-beam computed tomography scans of alginate impressions. *American Journal of Orthodontics and Dentofacial Orthopedics*, 143, 140–147.
16. Czarnota, J., Hey, J. and Fuhrmann, H. (2016) Measurements using orthodontic analysis software on digital models obtained by 3D scans of plaster casts. *Journal of Orofacial Orthopedics*, 77, 22–23.
17. Brook, P.H. and Shaw, W.C. (1989) The development of an index of orthodontic treatment priority. *European Journal of Orthodontics*, 11, 309–320.

18. Primožič, J., Perinetti, G., Contardo, L. and Ovsenik, M. (2013) Diagnostic performance of 3-dimensional evaluation of palatal vault changes in assessing successful treatment of constricted maxilla in growing subjects. *American Journal of Orthodontics and Dentofacial Orthopedics*, 143, 42–49.
19. Dunnett, C.W. and Tamhane, A.C. (1992) A step-up multiple test procedure. *Journal of the American Statistical Association*, 87, 162–170.
20. Generali, C., Primožič, J., Richmond, S., Bizzarro, M., Flores-Mir, C., Ovsenik, M. and Perillo, L. (2017) Three-dimensional evaluation of the maxillary arch and palate in unilateral cleft lip and palate subjects using digital dental casts. *European Journal of Orthodontics*, 39, 641–645.
21. Bondemark, L. and Abdullaheem, S. (2018) Intention to treat (ITT) analysis as reported in orthodontic randomized controlled trials-evaluations of methodology and recommendations for the accurate use of ITT analysis and handling dropouts. *European Journal of Orthodontics*, 40, 409–413.
22. Dahlberg, G. (1940) *Statistical Methods for Medical and Biological Students*. Allen and Unwin, London, UK, pp. 122–132.
23. Primožič, J., Baccetti, T., Franchi, L., Richmond, S., Farčnik, F. and Ovsenik, M. (2013) Three-dimensional assessment of palatal change in a controlled study of unilateral posterior crossbite correction in the primary dentition. *European Journal of Orthodontics*, 35, 199–204.
24. Primožič, J., Ovsenik, M., Richmond, S., Kau, C.H. and Zhurov, A. (2009) Early crossbite correction: a three-dimensional evaluation. *European Journal of Orthodontics*, 31, 352–356.

ISBN 978-91 7877-018-2 (print)

ISBN 978-91-7877-019-9 (pdf)

MALMÖ UNIVERSITY
205 06 MALMÖ, SWEDEN
WWW.MAU.SE