SUSANNA GILLBORG

OROFACIAL PAIN AND TOOTH WEAR IN SWEDISH ADULTS

OROFACIAL PAIN AND TOOTH WEAR IN SWEDISH ADULTS

© Copyright Susanna Gillborg 2019

Foto framsida Per Alstergren, foto baksida Linda Forslund, lay-out Martin Gillborg

ISBN 978-91-7877-048-9 (print)

ISBN 978-91-7877-049-6 (pdf)

ISSN 1650-6065

Holmbergs, Malmö 2019

**SUSANNA
GILLBORG**

**OROFACIAL PAIN AND
TOOTH WEAR
IN SWEDISH ADULTS**

Cross-sectional studies in southern Sweden

Malmö University, 2019
Faculty of Odontology
Orofacial Pain and Jaw Function

Publikationen finns även elektroniskt,
se muep.mau.se

To my parents, Carina and Lars-Göran Martinsson

CONTENTS

- PREFACE 9
- ABSTRACT 11
- POPULÄRVETENSKAPLIG SAMMANFATTNING 13
- INTRODUCTION 15
 - Paper I 15
 - Paper II 17
- AIMS 19
- HYPOTHESES 20
- MATERIAL AND METHODS 21
 - Paper I 21
 - Data analysis paper I 21
 - Paper II 22
 - Data analysis paper II 22
 - Statistical analysis 24
 - Flowchart of the selection process 25
- RESULTS 26
 - Paper I 26
 - Paper II 26
 - BEWE group 27
 - BEWE-adjusted group 28

DISCUSSION.....	29
Paper I	29
Response rate	29
Screening for TMD pain	29
Prevalence of TMD pain	30
TMD pain and Oral health-related quality of life	32
Screening for TMD pain in general practice	32
Paper II	33
Methods of measuring tooth wear	33
Prevalence of tooth wear	33
The etiology to tooth wear	33
No information about erosion.....	34
In the future	34
IMPLICATIONS	36
Paper I	36
Paper II	36
CONCLUSIONS	38
ACKNOWLEDGEMENTS	39
REFERENCES	41
PAPERS I-II.....	49

PREFACE

This thesis is based on the following papers, which will be referred to in the text by their numerals:

- Paper I: Gillborg S, Akerman S, Lundegren N, Ekberg EC.
Temporomandibular Disorder Pain and Related Factors
in an Adult Population: A Cross-Sectional Study in
Southern Sweden. J Oral Facial Pain Headache 2017
Winter;31(1):37-45.
- Paper II: Gillborg S, Akerman S, Ekberg EC.
Tooth wear in swedish adults – a cross-sectional study.
Journal of Oral Rehabilitation 2019 Sep 14. doi:
10.1111/joor.12887. (epub ahead of print).

Papers are reprinted with the kind permission from the publishers.

ABSTRACT

Aim. The present licentiate thesis investigated the prevalence of TMD-pain and related factors, the prevalence and severity of tooth wear, and the etiology and factors related to tooth wear in adults in southern Sweden.

Methods. The methods used included a questionnaire, history, clinical examination, intraoral photographs, and saliva sample. In Paper I, two screening questions for TMD pain were used to query a study sample comprising 6123 questionnaire participants about their pain experience. In Paper II, a clinical examination and intraoral photographs helped determine the presence and severity of tooth wear. Information from a questionnaire, patient histories, and participant saliva samples were analyzed regarding tooth wear-related factors. The study sample comprised 831 individuals.

Results. Paper I found a prevalence of TMD pain once a week or more often in 11% of the study sample. Related factors were female gender, subjects under 50 years of age, weekly headache, self-reports of poor general health, impaired oral health-related quality of life, and tooth wear. Paper II showed tooth wear in all individuals. Attrition, the most common tooth wear, was found in over 90% of the study sample. Signs of erosion were found in almost 80% of the individuals. Men had more tooth wear than women, but none of the factors that were investigated as related factors differed between the genders. Only some of the individuals, including the group with severe tooth wear reported having received information

about tooth wear from their clinician. Participants reported receiving information about tooth wear due to extensive tooth brushing more than about erosion.

Conclusions. Paper I found a prevalence of TMD pain in 11% of the study sample. In Paper II, attrition was found in over 90% of the study sample. Almost 80% of the individuals exhibited signs of erosion. Only a few reported having received information about tooth wear due to erosion from their clinician.

POPULÄRVETENSKAPLIG SAMMANFATTNING

Syfte. Denna licentiatuppsats undersökte prevalensen av Temporomandibulär Dysfunktions-smärta (TMD-smärta) och relaterade faktorer, prevalensen av tandslitage av olika svårighetsgrader och etiologin och relaterade faktorer till tandslitage i den vuxna befolkningen i södra Sverige.

Material och metod. Underlaget för studierna baseras på ett frågeformulär, anamnes, klinisk undersökning, intraorala foto och salivprov. I delarbete I användes två specifika frågor för att identifiera individer med TMD-smärta bland 6123 individer som deltog i studien. I delarbete II bestämdes prevalensen och graden av tandslitage genom kliniska undersökningar och granskning av kliniska foto. Information från frågeformuläret, anamnes och salivprov analyserades med avseende på möjliga relaterade faktorer till tandslitage. Studien innefattade 831 individer.

Resultat. Prevalensen av TMD-smärta en gång i veckan eller oftare i delarbete I var 11%. Relaterade faktorer var kvinnligt kön, åldrarna 20-50 år, huvudvärk en gång i veckan eller oftare, självrapporterad dålig allmänhälsa, försämrad livskvalité till följd av den orala hälsan och tandslitage.

I delarbete II uppvisade alla individer tecken på tandslitage. Attrition var den vanligast förekommande tandslitage och registrerades hos över 90% av individerna. Tecken på erosion registrerades hos nästan 80% av individerna. Män hade mer tandslitage än kvinnor men ingen av de relaterade faktorer som identifierades i studien skilde sig mellan könen. Endast en

del av individerna rapporterade att de hade fått information om tandslitage från sin tandläkare eller tandhygienist, även i gruppen med omfattande tandslitage. Fler uppgav ha fått information om tandslitage på grund av för hård tandborstning än om erosion.

Konklusion. Delarbete 1 fann en prevalens av TMD-smärta på 11% i den vuxna befolkningen. I delarbete II hade över 90% av de vuxna studiedeltagarna tecken på attrition. Tecken på erosion fanns hos nästan 80%. Endast en mindre del av individerna med tandslitage uppgav sig ha fått information av sin tandläkare eller tandhygienist om tandslitage orsakad av erosion, även bland de med omfattande tandslitage.

INTRODUCTION

Paper I

Life expectancy has long been a common measure for evaluating general health in a population; using that measure, health in Sweden has never been better (1). The mortality in some of the more common public diseases as heart and vascular diseases has decreased (2). The World Health Organization (WHO) defines health as the “state of complete physical, mental and social well-being and not merely the absence of disease or infirmity” (3). To live with a diagnose today does not necessarily mean that you experience a bad health. With modern medicine it is often possible to offer the patient a treatment, that will allow them to have a long life and experience a good general health.

Today, the most common reasons for adults to report a low level of general health are chronic pain and psychiatric disorders and are also the two most common reasons of all reported sick leave in Sweden (4). The Swedish Council on Health Technology Assessment (SBU) has estimated that, chronic pain alone has an overall societal cost in Sweden that exceeds SEK 80 billion per year, which corresponds to one-tenth of the annual budget expenses for the Swedish government.

Chronic pain is an important co-morbidity associated with several diseases or injuries, but is also acknowledged as a condition in its own (5,6). In the adult population in Europe, one of five report chronic pain of moderate to severe intensity (7). The most common chronic pain is back pain. Knee pain is next most common, with pain in the head area, including orofacial pain third most common. Temporomandibular disorder pain (TMD pain) is the most common non-dental-related orofacial pain condition (8). TMD is a collective term embracing a number of disorders affecting the temporomandibular joint, the masticatory musculature, and associated structures (9). The main symptoms are pain and functional limitation. The prevalence of TMD pain is reported to be 10–15%, it is more

common among women, and young and middle-aged adults are more often affected than children or the elderly(10,11).

The World Dental Federation defines oral health as being “Oral health is multi-faceted and includes the ability to speak, smile, smell, taste, touch, chew, swallow and convey a range of emotions through facial expressions with confidence and without pain, discomfort and disease of the cranio-facial complex.”(12). TMD patients report an impaired oral health-related quality of life (OHRQoL), and the more symptoms and signs that patients exhibit, the greater the impact of TMD on OHRQoL seems to be (13,14).

Psychiatric disorders are known risk factors both for developing chronic pain and for worsening the prognosis for recovery. In addition, long-term pain often has an impact on psychosocial well-being in the form of stress, anxiety, and depression. Epidemiological studies have found that many psychiatric disorders are more commonly found among individuals with chronic pain (15). In Europe, 21% of the individuals (24% in Sweden) with chronic pain report to have been diagnosed with depression because of their pain (7). Culture, genetics, and life style are factors that affect both psychiatric disorders and pain. For this reason prevalence will differ depending on the population sample. Moreover, the method used to investigate prevalence can strongly effect the results.

Pain is a complex condition, and there is no gold standard for the method of investigation (10,16). Most studies use questionnaires to measure self-reported pain and determine prevalence and intensity of TMD pain (11). In 2000, a Swedish study on TMD pain in adolescents queries 60 adolescents with a self-report of TMD pain and 60 controls (17). The adolescents were considered to have TMD pain if they answered yes to one or both of the following questions: (1) Do you have pain in your temples, face, jaw joint, or jaws once a week or more? (2) Do you have pain when you open your mouth wide or chew once a week or more?. Later, a clinician – blinded to the outcome of the questions – clinically examined the adolescents using standardized methods for diagnosing TMD pain: the RDC/TMD (Research Diagnostic Criteria for TMD) (18,19). In the group with self-reported TMD pain, 80% had myofascial pain compared to 3.3% in the control group. The two questions were thus considered to have very

good reliability and strong validity for screening TMD pain in adolescents.

In 2016, these screening questions were tested for validity in adults with TMD; together with a third question, on dysfunction. The procedure was again to compare the screening responses with the results of a standardized clinical examination, this time the Diagnostic Criteria for TMD (DC/TMD), which is a revision of the RDC/TMD. In this study, all patients coming for their routine dental check-up at the Public Dental Health Service in Västerbotten (northern Sweden) in 2014 were screened for TMD. Of 7,831 individuals, 152 screened positive for TMD. Another 148 gender- and age-matched individuals consented to join the study as controls. In the group with a self-report of TMD, 74% met the criteria for DC/TMD pain or dysfunction compared to 16% in the control group. The conclusion was that the screening questions also showed substantial validity to TMD in adults and especially for the TMD pain diagnoses (20).

Paper II

As with general health, oral health over time has changed for the better. The discovery and use of fluoride toothpaste was revolutionary for preventing caries (21). Today, although we are living longer, tooth loss and edentulism have become less common (22); other conditions, however, may come to play a larger role. In the younger population, tooth wear caused by dental erosion (23-26) seems to be increasing. There is much support in the literature since most studies on tooth wear focus on erosion in children and adolescents (27).

If the tooth wear seen in children and adolescents continues into adulthood, need for future dental care may increase. Rehabilitation of a worn dentition is often complex and time consuming. Because the vertical dimension has decreased and the teeth already have lost substance, conventional techniques that use preparations to create space for the restorative material are not possible and full-arch restorations are often needed to succeed (28,29).

Studies on tooth wear in adults have been sparse but interest is beginning to increase; still, studies on current levels of tooth wear in adults are

needed. Whether erosion is the most common cause of tooth wear in adults, as in children and adolescents, is unknown. If erosion is a common finding among adults, the question will be if the erosion is caused by acidic drinks, as reported in the literature for children and adolescents, or if tooth wear in adults has another etiology.

A review published in 2009 listed only 13 studies from various countries that had investigated the prevalence of tooth wear in adults (30). The review estimated the prevalence of severe tooth wear to be 3% among 20-year olds and 17% among 70-years olds. New studies on tooth wear in adults have since been published, but the variations in study design are substantial (31-34).

Due to differences in scoring systems, samples, and examiners, it is difficult to compare the results of the studies on tooth wear. Reviews of available literature report a wide range of prevalences. (23,30). There is, however, a consensus on the definition and diagnostic criteria of various forms of tooth wear (27). Since 1970, tooth wear has been classified according to three etiological factors: attrition, abrasion or erosion (35). Since then, abfraction has been suggested as a fourth wear-related process.

Efforts have been made to reach a consensus on an index that can be adaptable to other indexes, both in the collection and the presentation of data. For this reason, the Basic Erosive Wear Examination (BEWE) was developed at a workshop in 2007 (36,37). The scoring system quantifies the size of a given lesion in terms of the percentage of affected tooth surface. BEWE has been shown to have a sensitivity of 48.6% and a specificity of 96.1% for moderate tooth wear and 90.9% and 91.5% for severe tooth wear. (38,39)

AIMS

The aims were to:

- measure the prevalence of TMD pain in adults in southern Sweden (Paper I)
- evaluate the association of TMD pain with gender and other factors (Paper I)
- investigate the prevalence and severity of tooth wear in adults in southern Sweden (Paper II)
- study the etiology of the tooth wear (Paper II)
- assess factors that might be related to tooth wear (Paper II)

HYPOTHESES

The hypotheses of the studies were that:

- The prevalence of TMD pain would be in line with that reported in reviews, approximately 10-15%. (Paper I)
- Women between the ages of 20 and 50 would present a higher prevalence of TMD pain. (Paper I)
- Immigrants from outside the Nordic countries, individuals unemployed or on sick leave would have a higher prevalence of TMD pain. (Paper I)
- Self-reported bad general health and weekly headache would be related to a higher prevalence of TMD pain. (Paper I)
- Tooth wear would be found in almost all individuals. (Paper II)
- Only a small part of the population would have severe tooth wear. (Paper II)
- Attrition would be the most prevalent etiology in registered tooth wear. (Paper II)
- More than 50% of the individuals would have signs of erosion.
- The relation between tooth wear and consumption of soft drinks would be weak. (Paper II)
- The relation between tooth wear and fruit would be stronger than between tooth wear and soft drinks. (Paper II)
- Males would show more tooth wear than females. (Paper II)

MATERIALS AND METHODS

Paper I

Paper I used data from a large epidemiological study on oral health that had been carried out in southern Sweden in 2006-2007. The study on oral health was based on a questionnaire that was sent to 10,000 randomly selected individuals aged 20–89 years, which had been drawn from the Swedish Population Address Register (SPAR). To ensure a high response rate, four reminders was sent out in the next 12 months. The final response rate was 63%.

The questionnaire contained 58 questions, of which 2 were specific for identifying TMD pain (40):

Do you have pain in your face, jaw, temple, in front of the ear, or in your ear once a week or more often?

Do you have pain when you open your mouth or when you chew once a week or more often?

These questions are identical to questions used in the Research Diagnostic Criteria for Temporomandibular Disorders (RDC/TMD) and similar to the questions used by Nilsson 2000 and Lövgren 2016.

Data Analysis

Paper I

Individuals were regarded as having TMD pain if they answered yes to one or both of the above listed questions. Sixteen of the remaining 56 questions in the questionnaire were regarded having a possible relation to TMD pain and were selected for further analysis. After the first statistical analysis, 3 of the 16 questions were excluded: in one, there was no statistical relation to TMD pain in terms of frequency; in the other two, no

scientific hypothesis about the causative effect was possible. The remaining thirteen questions were studied in more detail using a logistic regression analysis (LRA) to determine their role in the etiology of TMD pain. The questions involved the following: demographics, experience of headache, self-reported general and oral health-related quality of life, satisfaction with life, number of missing teeth, information about tooth wear from their regular dentist/hygienist, and smoking habits.

Paper II

The material for paper II was collected from 2 larger epidemiological studies performed in 2 regions in southern Sweden. The first study was conducted in Skåne county in 2007–08, and the second in Kalmar County in 2011–12. Participants were randomly selected from SPAR. Each study comprised a questionnaire and a clinical examination. The Skåne study questionnaire contained 58 questions and was the same questionnaire used in Paper I. The Kalmar study questionnaire contained 56 questions; most were the same or similar to the questions in the Skåne study questionnaire. The clinical examinations in both studies included: a saliva sample; intraoral photographs; and a history, in which information about medication and dietary habits were obtained. All patients were examined using standard examination procedures. The dentists were coordinated regarding examination techniques through comprehensive written instructions, practice, and discussion of clinical cases.

In the Skåne study, the questionnaire to 1,000 individuals aged 20–89. All but 1 of the 451 individuals participating in the clinical examination completed the questionnaire. In the Kalmar study, the questionnaire was sent to 2,000 individuals aged 20–90; 900 of these were offered a dental examination, and 380 individuals were examined.

Data Analysis

Paper II

The thesis author (SG) compiled and analyzed the data on tooth wear for the 831 participants. Tooth wear level was recorded using the Basic Erosive Wear Examination (BEWE). The calibration of SG with a specialist in orofacial pain and jaw function had moderate inter-rater agreement (41). To minimize the risk of overlooking clinical signs of tooth wear, a

calibration for recording clinical signs of tooth wear was done twice: before the clinical examinations and another one for evaluating tooth wear on photographs. The level of tooth wear was then estimated by supplementing the results of the clinical examination with information from the intraoral photographs. Any tooth wear found in the photographs that had been undetected clinically was added to the record. SG then used BEWE to estimate the level of tooth wear for all participants.

BEWE – No risk

BEWE – low risk

BEWE – medium risk

BEWE – high risk

The clinical appearance of tooth wear was evaluated according to the descriptions of the different types of tooth wear: attrition, erosion, abrasion and abfraction (27,42). Signs of attrition were registered if glossy plane facets with sharp margins were found on occluding surfaces. Reversed V-sign incisal on maxillary central incisors, thin enamel on the palatal surface of the maxillary incisors, loss of the distinct shape on the tooth surface, thinning out of the enamel without signs of attrition, loss of surface lustre, raised restorations, grooves on the cusps or incisal edges, and/or restorations rising above the level of the adjacent morphology, were all considered signs of erosion. Signs of attrition and erosion were the most easily identifiable. Abrasion was difficult to distinguish from erosion without more detailed histories of each individual and was not recorded. Abfraction, was seen as a possible part of the etiology of noncarious cervical lesions, and we decided to report the prevalence of NCCLs instead.

Information from the questionnaires, histories, and examinations allowed us to study factors related to tooth wear. Information from questions and clinical examinations of interest in relation to tooth wear underwent further analysis: demographics, knowledge about tooth wear, dietary habits, medications, tobacco use, alcohol consumption, saliva (quantity and quality), and experience of TMD pain.

Statistical Analysis

Paper I

Descriptive statistics with means and standard deviations characterized individuals with and without pain. The χ^2 -test compared the distribution of categorical variables. Possible related factors related to TMD pain were analyzed using logistic regression models with a likelihood-ratio test. In the first model, TMD pain was tested by correcting for the related factors. In the second and third models, gender was tested for TMD pain by correcting for the related factors.

Paper II

Participants were stratified on the basis of risk levels of tooth wear and age group. The χ^2 -test compared the distribution of categorical variables. Possible factors related to levels of tooth wear were analyzed using logistic regression models with a likelihood-ratio test.

Figure 1. Flowchart of the selection process for the Paper I and II study cohorts.

RESULTS

Paper I

The prevalence of self-reported TMD pain (once a week or more often) in Swedish adults was 11.0% in 2006–2007. TMD pain prevalence in women and individuals under age 50 was significantly higher than the mean value for the entire study group (aged 20–89). Prevalence was also higher in individuals who were unemployed, on sick leave, or taking ill-health pension. Individuals living with a partner reported less TMD pain than individuals living alone or in other family configurations.

Prevalence also differed depending on birth country. TMD pain prevalence in individuals born in Sweden or other Nordic countries was similar to the prevalence in individuals born in Western European countries and in North and South America but lower than in other nationalities.

Individuals with TMD pain reported more headache and more often experienced a poorer general and oral health than others of the same age. A higher score on the Oral Health Impact Profile scale (OHIP-14) was found among individuals with TMD pain, which means that individuals with TMD pain felt that their oral health had a bad influence on their everyday life to a larger extent than individuals without TMD pain; they also felt less satisfied with life more often. Men and women showed the same related factors for TMD pain.

Paper II

Data from 831 individuals were collected. Of these, 54 were edentulous or had restorations covering most or all tooth substance, and had to be excluded. Another 106 individuals with partial dentition restoration or who were partially edentulous could not be assigned a BEWE score since the index was designed for use on individuals with a full dentition; thus, tooth wear in 81% (671) of the study group could be evaluated with the BEWE (BEWE group). The 106 individuals who could not be assigned a

BEWE score were analyzed separately (BEWE-adjusted group) by multiplying their results to obtain a comparable score.

Table 1. Participants in the Skåne questionnaire study (2006–2007; Paper I) and the questionnaire + clinical exam study (Kalmar, 2011–2012; Skåne, 2007–2008; Paper II). The studies used in Paper II comprised a Basic Erosive Wear Examination (BEWE) group (a) and a BEWE-adjusted group (b).						
	No.	Mean age yrs (SD)	Women		Men	
			(n)	(%)	(n)	(%)
Paper I	6123	51 (17.5)	3480	57	2643	43
Paper II						
(a)	671	46 (15.5)	347	52	324	48
(b)	106	69 (9.8)	58	55	48	45

Only 4 individuals in the BEWE-adjusted group were younger than 50 years, whereas all ages up to 70 were similarly represented in the BEWE group.

The BEWE group

In the BEWE group, nearly 80% of the group exhibited signs of erosion and over 90%, signs of attrition. Retraction of the gingival margin was registered in 78%, and NCCLs were registered on at least one tooth in 29% of the individuals and became more frequent with higher levels of tooth wear. All individuals with a high-risk level of tooth wear had signs of both erosion and attrition.

Significant gender-related differences were found: the no-risk group comprised 68% women and 32% men, whereas the high-risk group comprised 39% women and 61% men.

In total, half (52%) of the individuals in the high-risk group reported that their dentist or hygienist had informed them about tooth wear due to tooth

grinding, and 10% reported being informed about erosion. Of the individuals with NCCL, 40% reported being informed about tooth wear due to extensive tooth brushing.

Individuals with severe tooth wear had more fractured teeth and/or fillings. Tooth wear was more related to daily consumption of acidic fruit than to consumption of acidic drinks. No differences in related factors (signs of erosion, signs of attrition, NCCLs, gingival retractions, fractured teeth and/or fillings, information about tooth grinding and daily consumption of fruit) were found between the genders.

Table 2. Frequency of the different severities of tooth wear in the BEWE group and BEWE-adjusted group.								
Group	No risk		Low risk		Medium risk		High risk	
	(n)	(%)	(n)	(%)	(n)	(%)	(n)	(%)
BEWE	22	3.3	333	49.6	285	42.5	31	4.6
BEWE-adjusted	0	0	27	25.5	56	52.8	23	21.7

The BEWE-adjusted group

More than 60% of the individuals exhibited signs of erosion and almost all, signs of attrition. For NCCL, the figure was almost the same as in the BEWE group, with signs in at least one tooth in 29% of the individuals. All individuals in the high-risk group had signs of attrition, as in the BEWE group, but only 70% showed signs of erosion.

DISCUSSION

Paper I

Response rate

Paper I is based on a questionnaire that was mailed to 10,000 randomly selected individuals in 2006-2007. After three reminders, the final response rate was 63% ($n = 6,123$). A systematic review of response rates in epidemiologic and public health studies compared response rates to web-based surveys with more traditional data collection methods such as mail or phone and concluded that traditional data collection resulted in a higher response rate (43). The mean response rate reported by the studies in the review, which used traditional data collection methods and were carried out during 2002–2009 was 40% with a 95% CI of 23.7–56.3 (43). In comparison, the response rate of 63% in the present study can be considered high.

Screening for TMD pain

The two screening questions used in this study had been widely used since 1992, when they were incorporated into the RDC/TMD (18). One of these screening questions is also identical to one of the screening questions used and validated by Nilsson et al and Lövgren et al to screen for TMD pain (17,20)(20). The other question is similar to the screening question used by Nilsson et al and Lövgren, but refers to the area of the jaw joint as “in front of the ear or in the ear” whereas Nilsson et al and Lövgren et al write “the jaw joint”. The Nilsson et al study was published after the study in Paper I had begun collecting data and the study from Lövgren et al, some years later, in 2016.

If the study in Paper I had been performed today, the screening questions would have been identical to the questions used by Lövgren et al. as these questions have been validated for screening of TMD pain in adults. It can only be speculated whether referring to the jaw joint as the area in front of the ear or in the ear strengthens validity by decreasing false negatives

of individuals who lack knowledge of the anatomy of the mandible, or weakens validity by increasing false positive responses of individuals with recurrent ear pain of other origin.

Prevalence of TMD pain

In the present study, the prevalence of TMD pain among adults in southern Sweden was 11%. Between May 2010 and October 2012 in northern Sweden (Västerbotten County), 55% of the county population ($n = 137,718$ individuals) was screened for TMD pain with the same questions that were later validated by Lövgren for screening TMD pain in adults. This northern Sweden study found that 5.3% screened positive for TMD pain in the 20–69-year age group (44). Between November 2006 and September 2008, clinicians in another part of Sweden (Södermanland County) screened 2,837 adults for TMD pain at a Public Dental Health Service (PDHS) clinic during routine dental examinations, using the same screening questions. They reported a prevalence of TMD pain of 3.7%. Between 2011 and 2013 all adults (144,000), who attended the PDHS in the same area as previous study were screened for TMD pain, again with the same screening questions. Registrations were made for 51% of the individuals and the prevalence was 1.5% (45).

Why these three Swedish studies found different prevalences of TMD pain in adults, despite using the same screening question, appears puzzling. The study in Paper I was based on volunteers, with the attendant risk of selection bias. The second study at the PDHS clinics reported a dropout of nearly half in the responses to the screener. An uncertain risk is whether clinicians who forgot to ask the screening questions simply registered negative responses in order to complete the form, thus lowering prevalence.

Possibly, prevalence does vary between different areas of Sweden. The regions differ in other ways. The population density in southern Sweden is 2.5 times higher than in Södermanland County and 25 times higher than in Västerbotten County (46). At the time of the investigations, 8% of the population in Västerbotten County had been born outside Sweden compared to 18% in southern Sweden (47). The prevalence of TMD pain

among immigrants varied substantially, depending on the individual's reported birth country. These results should be interpreted with caution since some nationalities are represented only by a small group of individuals. A study performed in Saudi Arabia reported a much higher frequency of TMD pain than European and American studies have suggested, indicating that prevalence could be much higher in other parts of the world (48). The large differences in prevalence of TMD pain among immigrants is probably also due to socioeconomic differences and the sometimes difficult process of acculturation which can cause stress (49). Such stress could heightened the pain sensitivity and affect the ability to cope with pain.

Table 2. Prevalence of self-reported TMD pain according to birth country (n=6,123) in adult residents of southern Sweden (Paper I).

Birth country	TMD pain (n=648) (%)	95% CI	<i>P</i> -value
Sweden (n=5117)	10.1	9.3–11.0	.000
Other Nordic country (n=161)	11.2	6.3–16.1	
Former Yugoslavia (n=141)	22.0	15.1–28.9	
Former Soviet (n=16)	37.5	10.9–64.1	
Western Europe (n=94)	10.6	4.3–17.0	
Eastern Europe (n=104)	15.4	8.3–22.4	
Iran/Iraq (n=70)	21.4	11.6–31.3	
Asia (n=84)	20.2	11.5–29.0	
Africa (n=26)	23.1	5.7–40.4	
North/South America (n=38)	10.5	0.3–20.7	

TMD = temporomandibular disorders; CI = confidence interval.

TMD pain and Oral Health-related Quality of Life

We found an association between TMD pain and high scores on the OHIP. This is in line with other reports on TMD pain and its negative effect on oral health-related quality of life (14,50). Pain affects individuals to varying degrees. If the individual has bad expectations and experience that the pain has a negative affect on the quality of life, this have been shown to be a risk factors for future pain and disability and worsens the prognosis for recovery (51). Several easy-to-administer, well-validated screening instruments for exploring the thoughts, feelings, and expectations of the individual are available and could be used in dental care(52,53). The DC/TMD includes screening and comprehensive self-report instrument sets, which assess psychological distress (52-54). This additional information about the patient could improve patient management and through communication between dental and general health care, also help patients access professional help at an earlier stage.

Screening for TMD pain in general dentist practice

TMD pain is underdiagnosed and undertreated in Sweden (55-57). In an effort to improve identification of individuals with TMD pain, TMD pain screening using the validated questions was introduced in several regions through the PDHS. Lövgren et al have evaluated the impact of the introduced screening questions on clinical decision-making. Additional to the TMD pain questions, a question about dysfunction was added. Significantly, more treatment was done or recommended for the individuals with one or more positive answers to the screening questions, but according to the dental records, the majority of individuals screening positive for TMD received no assessment or treatment (58). A 5-year follow-up reviewed TMD-related treatment decisions as reported in the dental records by selecting a random sample of 200 individuals with at least one positive response to the screening questions and 200 controls. In 45.5% of the cases, a clinical decision related to TMD was absent, indicating undertreatment of TMD in general dental practice despite identification of individuals with likely TMD (59).

Paper II

Methods of measuring tooth wear

Tooth wear studies have over 100 indices from which to choose to evaluate the severity of tooth wear (42). Although the index used in Paper II has not been the most frequently used historically, it was designed to be able to allow comparison between studies using different types of index (36). The weakness is that although it is relatively insensitive, inter-examiner agreement is moderate and acceptable. It is also designed for use in individuals with a full or almost full dentition, which results in the exclusion of some parts of the population, especially in the older age groups. As with all indices designed to measure visible damage, there is a risk of the individual developing unnecessary, irreversible damage before their dentist or dental hygienist become aware of it.

Prevalence of tooth wear

Being able to study tooth wear in clinical photos made it possible to systematically and in detail search for even the smallest signs of tooth wear. Some signs of tooth wear was found in all individuals. Severity varied from a single bruxofascett to loss of more than half of the tooth substance. The severity of tooth wear increased with age. The risk of tooth wear was high in 3% of the youngest group (20–29 years). No individual in the oldest group (80–89 years) had enough remaining tooth substance to determine a score on the BEWE. In the 70–79-year age group, 11% had a high risk of tooth wear. If the BEWE-adjusted group had been included, this risk would be even higher in the older age groups. As in all epidemiological studies, the study populations need to be comparable in order to compare results with other studies. Many studies on tooth wear in adults only include individuals up to age 35, and for lack of a standard, a wide variety of scoring systems is used. Thus, comparisons between studies are challenging (23,30).

The etiology of tooth wear

The etiology in each case of tooth wear was determined solely on clinical appearance. Many well-written descriptions of the clinical signs for the four types of tooth wear can be found in the literature (27,42). Anamnestic information was an additional source of information, but it was not used as a discriminator; known risk factors such as acidic diet, when sourced

from patient histories, would be able to show a relation to tooth wear at the population level, but not at the individual level. Commonly, the history noted current risk factors and did not query, for instance, diet habits earlier in life. It was impossible to determine whether tooth wear was new or old since the measurement was only made once. The true causative factor behind the tooth wear was not always shown in the history.

No information on erosion

A major finding in our study was that individuals with severe tooth wear due to erosion did not report having received information about erosion from their dentist or dental hygienist. This result was unexpected since it is well known that damage due to erosion has been increasing in children and adolescents for many years(60) and should be checked at the annual visits, also for adults.

Between 2009 and 2016, prescriptions for new oral appliances in Sweden increased by 30% (61). During this time, the population increased only 7%. One could speculate whether dentists underdiagnose erosion and incorrectly treat patients with an oral appliance.

In the future

To avoid underdiagnosis of TMD pain, screening questions should be added to the routine anamnesis for the patients when they see their dentist or dental hygienist for their yearly check-up. In Sweden, the two screening questions validated by Nilsson et al in adolescents and thereafter by Löfgren et al in adults has already been implemented in several regions in Sweden. Other versions of screening questions have been presented in the literature (62), and perhaps, different countries/populations will need to develop and validate their own country-specific screener so that it will be compatible with their population. The next question is how to support general practitioners in their treatment of patients with identified TMD pain.

Tooth wear seems to be increasing not only in children and adolescents but also among adults. Dentists and dental hygienists need more knowledge about tooth wear and, especially, dental erosion. A standardized method for identifying pathological progression rates of tooth wear

should be available. For this, research needs to establish normative values for tooth wear and a screening method that is easy to use. With the technology available today, with intraoral scanners, perhaps we can detect abnormal progression rates before detectable for the eye. Preventative treatment could then be provided early and avoid unnecessary and irreversible damage to the dentition.

The patient history taken in the Paper II studies could not explain the cause of tooth wear at an individual level. Thus, there is a need for improved questionnaires that also query old habits in order to better understand the etiology of the tooth wear.

IMPLICATIONS

Paper I

The consequences of chronic pain are not only physical but also mental and social. By interfering with the ability to participate in the normal activities of life, chronic pain is responsible for substantial costs to not only the individual but also society. To minimize the risk of developing chronic pain, individuals with higher risk must be identified as soon as possible, in an early stage.

Knowledge of TMD pain prevalence is important for determining the scope and severity of the condition in the population. To identify TMD pain-related factors, is necessary for a better understanding of the etiology. Information on prevalence and related factors are valuable for strategic planning in the dental health service, to be able to identify these individuals and provide appropriate care.

Paper II

Tooth wear has been recognized as a growing oral health problem among children and adolescents, and it seems that this is also true among adults. Studies have found that erosion is most often involved in the etiology of severe tooth wear; erosion weakens and speeds up the loss of tooth substance in the presence of abrasion or attrition. In Paper II, all individuals with severe tooth wear had signs of erosion. Despite this, only a few individuals reported having received information about erosion from their clinician. This indicates a need for more knowledge of tooth wear, and for how to detect and interpret clinical findings.

Today, no normative values have yet been set for tooth wear at different ages. The results of this study provide information on the current prevalence of tooth wear in the adult population and on the variances in severity at different ages. The results of this study could serve as a basis for future studies that wish to track how tooth wear in adults changes over time or

wish to identify normative values for the various age groups. Furthermore, the results may help clinicians in general dental practice change their approach regarding tooth wear and be more observant to changes in tooth wear caused by erosion. Knowledge of current prevalences of tooth wear in different ages can help the clinician better identify those individuals in need of prophylactic actions to prevent further wear. The anamnestic information collected in our study on possible causative factors did not always explain the recorded tooth wear. This indicates that a standard anamnesis is not always able to identify the etiological factor behind tooth wear. The clinician will sometimes need to expand the anamnesis to include past habits and more unusual causative factors. The clinician will need to determine etiology by the clinical appearance of tooth wear. Clinical photos are an easy and cost-effective way to document tooth wear and monitor changes over time.

CONCLUSIONS

The prevalence of TMD pain was 11% in the adult population in southern Sweden, which as expected, is in line with figures that reviews have reported. Prevalence was higher among women than men and highest prevalence was found in individuals aged 20–49 years. Immigrants from Asia, Africa, Eastern Europe, Former Yugoslavia, Former Soviet and Iran/Iraq and individuals who were unemployed or on sick leave had a higher prevalence of TMD pain. Individuals with TMD pain more often reported bad general health and high frequency of headache (once a week or more often). They also reported more impaired oral health-related quality of life than individuals without TMD pain.

Tooth wear was registered in the 831 individuals participating in this study. Severe tooth wear was found in 4.6% of the individuals. Attrition was the most prevalent etiology, but erosion occurred to a larger extent than was expected: almost 80% had signs of erosion. Males had more severe tooth wear. Tooth wear was more related to daily consumption of acidic fruit than to consumption of acidic drinks.

The results from this licentiate thesis can be used both by general dentists and by specialists in identifying patients with TMD pain and tooth wear of different etiology and provide appropriate care.

ACKNOWLEDGEMENTS

I wish to thank my primary supervisor and trusted mentor, Professor EwaCarin Ekberg. You noticed my special interest in orofacial pain and jaw function early on, when I was an undergraduate dental student at Malmö University. You have guided me in the clinic, in research, and sometimes also in life, with empathy and a unique ability to sense what is needed: a push or patience.

I would also like to say thank you to:

Professor Sigvard Åkerman, my secondary supervisor, for letting me take part of all the data collected in the Skåne and Kalmar studies. For valuable criticism in Paper I and Paper II.

Nina Lundegren, Ph.D., for your generosity in sharing your work, and for your help with formatting Materials and Methods in Paper I.

To the Faculty of Odontology at Malmö University, Sweden, who made it possible for me to do this licentiate thesis and with financially support, to participate in international courses and congresses.

Per-Erik Isberg, for being service minded, patient, and pedagogical when performing the statistical analyses together with me, EwaCarin, and Sigvard.

All staff members of the Faculty of Odontology, who were my supervisors when I was an undergraduate and who now are my inspiring colleagues. To be able to work in this environment is inspirational. Thank you.

The dental assistants I have worked with who supported me when my mind perhaps was scattered, thinking of too many things at the same time.

The Language Editing Group at Malmö University and Gail Conrod for linguistic revision.

To the Public Dental Health Service in Skåne, and especially my old clinic in Genarp, for all the help and support when I was absent for 50% of the week for several years.

My “extended family” of orofacial pain specialists in Sweden and around the world for including me, supporting me, and being people I could look up to.

My beloved friends, old and new. My childhood friend, Christina Brunner, for all the fun we had through the years, you know a part of me no one else knows. My good friend and therapist, Michelle Andersson, it is a blessing to have you. Sandhya Nair for loyal friendship and the gift of seeing things differently. My old gang Jenny Flygare, Annika Rask, and Sofia Wallinder for loving friendship, hard work, and cooking.

My parents-in-law, for support in everyday life. I am truly blessed to have you.

My parents, Carina and Lars-Göran Martinsson, for giving me courage and self-esteem. For endless support even as a grown-up.

My sisters for unconditional love. One for all, all for one. I love you.

My husband and big love, Martin. For everything you do for me and our children everyday.

This work has been supported by the Regional Board of Dental Public Health in Skåne and Kalmar, Sweden.

REFERENCES

- (1) Statistiska Centralbyrån. Life expectancy 17-51-2018. 2018; Available at: <https://www.scb.se/en/finding-statistics/statistics-by-subject-area/population/population-composition/population-statistics/pong/tables-and-graphs/yearly-statistics--the-whole-country/life-expectancy/>. Accessed 9/19, 2019.
- (2) Socialstyrelsen. Öppna jämförelser 2014, Folkhälsa. 2015; Available at: <https://www.socialstyrelsen.se/globalassets/sharepoint-dokument/artikelkatalog/oppna-jamforelser/2014-12-3.pdf>. Accessed 10/21, 2019.
- (3) WHO. Constitution. 2019; Available at: <https://www.who.int/about/who-we-are/constitution>. Accessed 10/20, 2019.
- (4) Försäkringskassan. Rapport - uppföljning av sjukfrånvarons utveckling 2018. 2019; Available at: <https://www.forsakringskassan.se/wps/wcm/connect/d3d2d056-0ae7-46d9-b350-ac87e4696f1c/uppfoljning-av-sjukfranvarons-utveckling-2018.pdf?MOD=AJPERES&CVID=>. Accessed 10-24, 2019.
- (5) Tracey I, Bushnell MC. How neuroimaging studies have challenged us to rethink: is chronic pain a disease? J Pain 2009 Nov;10(11):1113-1120.
- (6) Nicholas M, Vlaeyen JWS, Rief W, Barke A, Aziz Q, Benoliel R, et al. The IASP classification of chronic pain for ICD-11: chronic primary pain. Pain 2019 Jan;160(1):28-37.

- (7) Breivik H, Collett B, Ventafridda V, Cohen R, Gallacher D. Survey of chronic pain in Europe: prevalence, impact on daily life, and treatment. *Eur J Pain* 2006 May;10(4):287-333.
- (8) McNeill C. Management of temporomandibular disorders: concepts and controversies. *J Prosthet Dent* 1997 May;77(5):510-522.
- (9) American Academy of Orofacial Pain, De Leeuw R, American Academy of Orofacial Pain, American Academy of Orofacial Pain. Orofacial pain: guidelines for assessment, diagnosis, and management. 5th ed. Chicago: Quintessence; 2013.
- (10) Macfarlane TV, Glenny AM, Worthington HV. Systematic review of population-based epidemiological studies of oro-facial pain. *J Dent* 2001 Sep;29(7):451-467.
- (11) LeResche L. Epidemiology of temporomandibular disorders: implications for the investigation of etiologic factors. *Crit Rev Oral Biol Med* 1997;8(3):291-305.
- (12) The World Dental Federation. FDI's definition of oral health. 2019; Available at: <https://www.fdiworlddental.org/oral-health/fdi-definition-of-oral-health>. Accessed 10/20, 2019.
- (13) Oghli I, List T, John M, Larsson P. Prevalence and oral health-related quality of life of self-reported orofacial conditions in Sweden. *Oral Dis* 2017 Mar;23(2):233-240.
- (14) Dahlstrom L, Carlsson GE. Temporomandibular disorders and oral health-related quality of life. A systematic review. *Acta Odontol Scand* 2010 Mar;68(2):80-85.
- (15) Velly AM, Mohit S. Epidemiology of pain and relation to psychiatric disorders. *Prog Neuropsychopharmacol Biol Psychiatry* 2018 Dec 20;87(Pt B):159-167.

- (16) Crombie, I.K. Croft, P.R. Linton, S.J. LeResche, L. Von Korff, M. editor. *Epidemiology of Pain*. Seattle, USA: IASP Press; 1999.
- (17) Nilsson IM, List T, Drangsholt M. The reliability and validity of self-reported temporomandibular disorder pain in adolescents. *J Orofac Pain* 2006 Spring;20(2):138-144.
- (18) Dworkin SF, LeResche L. Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. *J Craniomandib Disord* 1992 Fall;6(4):301-355.
- (19) List T, Dworkin SF. Comparing TMD diagnoses and clinical findings at Swedish and US TMD centers using research diagnostic criteria for temporomandibular disorders. *J Orofac Pain* 1996 Summer;10(3):240-253.
- (20) Lovgren A, Visscher CM, Haggman-Henrikson B, Lobbezoo F, Marklund S, Wanman A. Validity of three screening questions (3Q/TMD) in relation to the DC/TMD. *J Oral Rehabil* 2016 Oct;43(10):729-736.
- (21) O'Mullane DM, Baez RJ, Jones S, Lennon MA, Petersen PE, Rugg-Gunn AJ, et al. Fluoride and Oral Health. *Community Dent Health* 2016 Jun;33(2):69-99.
- (22) Hugoson A, Koch G, Gothberg C, Helkimo AN, Lundin SA, Norderyd O, et al. Oral health of individuals aged 3-80 years in Jonkoping, Sweden during 30 years (1973-2003). II. Review of clinical and radiographic findings. *Swed Dent J* 2005;29(4):139-155.
- (23) Jaeggi T, Lussi A. Prevalence, incidence and distribution of erosion. *Monogr Oral Sci* 2014;25:55-73.
- (24) Bartlett DW. The role of erosion in tooth wear: aetiology, prevention and management. *Int Dent J* 2005;55(4 Suppl 1):277-284.
- (25) Johansson AK. On dental erosion and associated factors. *Swed Dent J Suppl* 2002;(156)(156):1-77.

- (26) Carlsson GE, Johansson A. Prevalens av dental erosion. Dental erosion: Förlagshuset Gothia; 2006.
- (27) Ganss C, Young A, Lussi A. Tooth wear and erosion: methodological issues in epidemiological and public health research and the future research agenda. *Community Dent Health* 2011 Sep;28(3):191-195.
- (28) Lussi A, Carvalho TS. Erosive tooth wear: a multifactorial condition of growing concern and increasing knowledge. *Monogr Oral Sci* 2014;25:1-15.
- (29) Johansson A, Johansson AK, Omar R, Carlsson GE. Rehabilitation of the worn dentition. *J Oral Rehabil* 2008 Jul;35(7):548-566.
- (30) Van't Spijker A, Rodriguez JM, Kreulen CM, Bronkhorst EM, Bartlett DW, Creugers NH. Prevalence of tooth wear in adults. *Int J Prosthodont* 2009 Jan-Feb;22(1):35-42.
- (31) Wetselaar P, Vermaire JH, Visscher CM, Lobbezoo F, Schuller AA. The prevalence of tooth wear in the Dutch adult population. *Ned Tijdschr Tandheelkd* 2018 Apr;125(4):205-213.
- (32) Awad MA, El Kassas D, Al Harthi L, Abraham SB, Al-Khalifa KS, Khalaf ME, et al. Prevalence, severity and explanatory factors of tooth wear in Arab populations. *J Dent* 2019 Jan;80:69-74.
- (33) Vered Y, Lussi A, Zini A, Gleitman J, Sgan-Cohen HD. Dental erosive wear assessment among adolescents and adults utilizing the basic erosive wear examination (BEWE) scoring system. *Clin Oral Investig* 2014 Nov;18(8):1985-1990.
- (34) Bartlett DW, Lussi A, West NX, Bouchard P, Sanz M, Bourgeois D. Prevalence of tooth wear on buccal and lingual surfaces and possible risk factors in young European adults. *J Dent* 2013 Nov;41(11):1007-1013.
- (35) Pindborg J editor. Pathology of the dental hard tissues. Copenhagen: Munksgaard; 1970.

- (36) Bartlett D, Ganss C, Lussi A. Basic Erosive Wear Examination (BEWE): a new scoring system for scientific and clinical needs. *Clin Oral Investig* 2008 Mar;12 Suppl 1:S65-68.
- (37) Young A, Amaechi BT, Dugmore C, Holbrook P, Nunn J, Schiffner U, et al. Current erosion indices--flawed or valid? Summary. *Clin Oral Investig* 2008 Mar;12 Suppl 1:S59-63.
- (38) Bernhardt O, Gesch D, Splieth C, Schwahn C, Mack F, Kocher T, et al. Risk factors for high occlusal wear scores in a population-based sample: results of the Study of Health in Pomerania (SHIP). *Int J Prosthodont* 2004 May-Jun;17(3):333-339.
- (39) Fares J, Shirodaria S, Chiu K, Ahmad N, Sherriff M, Bartlett D. A new index of tooth wear. Reproducibility and application to a sample of 18- to 30-year-old university students. *Caries Res* 2009;43(2):119-125.
- (40) Nilsson IM. Reliability, validity, incidence and impact of temporomandibular pain disorders in adolescents. *Swed Dent J Suppl* 2007(183):7-86.
- (41) Viera AJ, Garrett JM. Understanding interobserver agreement: the kappa statistic. *Fam Med* 2005 May;37(5):360-363.
- (42) Wetselaar P. A PhD completed 6. Tooth Wear Evaluation System: development and applications. *Ned Tijdschr Tandheelkd* 2016 Oct;123(10):492-494.
- (43) Blumenberg C, Barros AJD. Response rate differences between web and alternative data collection methods for public health research: a systematic review of the literature. *Int J Public Health* 2018 Jul;63(6):765-773.
- (44) Lovgren A, Haggman-Henrikson B, Visscher CM, Lobbezoo F, Marklund S, Wanman A. Temporomandibular pain and jaw dysfunction at different ages covering the lifespan--A population based study. *Eur J Pain* 2016 Apr;20(4):532-540.

- (45) Adern B, Minston A, Nohlert E, Tegelberg A. Self-reportance of temporomandibular disorders in adult patients attending general dental practice in Sweden from 2011 to 2013. *Acta Odontol Scand* 2018 Oct;76(7):530-534.
- (46) Statistiska Cantral Byrån. Befolkningstäthet (invånare per kvadrat-kilometer), folkmängd och landareal efter region och kön. År 1991 - 2018. 2019; Available at: http://www.statistikdatabasen.scb.se/pxweb/sv/ssd/START_BE_BE0101_BE0101C/BefAreal-TathetKon/?loadedQueryId=65972&timeType=from&timeValue=2000. Accessed 17/10, 2019.
- (47) Statistiska Central Byrån. Inrikes och utrikes födda efter region, ålder och kön. År 2000 - 2018. 2019; Available at: http://www.statistikdatabasen.scb.se/pxweb/sv/ssd/START_BE_BE0101_BE0101E/InrUtr-FoddaRegAlKon/?loadedQueryId=65089&timeType=from&timeValue=2000. Accessed 10/20, 2019.
- (48) Al-Harthy M, Al-Bishri A, Ekberg E, Nilner M. Temporomandibular disorder pain in adult Saudi Arabians referred for specialised dental treatment. *Swed Dent J* 2010;34(3):149-158.
- (49) Edwards CL, Fillingim RB, Keefe F. Race, ethnicity and pain. *Pain* 2001 Nov;94(2):133-137.
- (50) John MT, Reissmann DR, Schierz O, Wassell RW. Oral health-related quality of life in patients with temporomandibular disorders. *J Orofac Pain* 2007 Winter;21(1):46-54.
- (51) Boersma K, Linton SJ. Expectancy, fear and pain in the prediction of chronic pain and disability: a prospective analysis. *Eur J Pain* 2006 Aug;10(6):551-557.
- (52) Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001 Sep;16(9):606-613.

(53) Lowe B, Decker O, Muller S, Brahler E, Schellberg D, Herzog W, et al. Validation and standardization of the Generalized Anxiety Disorder Screener (GAD-7) in the general population. *Med Care* 2008 Mar;46(3):266-274.

(54) Schiffman E, Ohrbach R, Truelove E, Look J, Anderson G, Goulet JP, et al. Diagnostic Criteria for Temporomandibular Disorders (DC/TMD) for Clinical and Research Applications: recommendations of the International RDC/TMD Consortium Network* and Orofacial Pain Special Interest Groupdagger. *J Oral Facial Pain Headache* 2014 Winter;28(1):6-27.

(55) Al-Jundi MA, John MT, Setz JM, Szentpetery A, Kuss O. Meta-analysis of treatment need for temporomandibular disorders in adult nonpatients. *J Orofac Pain* 2008 Spring;22(2):97-107.

(56) Nilsson IM, List T, Drangsholt M. Prevalence of temporomandibular pain and subsequent dental treatment in Swedish adolescents. *J Orofac Pain* 2005 Spring;19(2):144-150.

(57) Wanman A, Wigren L. Need and demand for dental treatment. A comparison between an evaluation based on an epidemiologic study of 35-, 50-, and 65-year-olds and performed dental treatment of matched age groups. *Acta Odontol Scand* 1995 Oct;53(5):318-324.

(58) Lovgren A, Marklund S, Visscher CM, Lobbezoo F, Haggman-Henrikson B, Wanman A. Outcome of three screening questions for temporomandibular disorders (3Q/TMD) on clinical decision-making. *J Oral Rehabil* 2017 Aug;44(8):573-579.

(59) Lovgren A, Karlsson Wirebring L, Haggman-Henrikson B, Wanman A. Decision-making in dentistry related to temporomandibular disorders: a 5-yr follow-up study. *Eur J Oral Sci* 2018 Dec;126(6):493-499.

(60) Johansson AK, Omar R, Carlsson GE, Johansson A. Dental erosion and its growing importance in clinical practice: from past to present. *Int J Dent* 2012;2012:632907.

(61) Forsakringskassan. 2016; Available at: <https://www.forsakringskassan.se/statistik/ovrigaersatt/tandvard>. Accessed 02/01, 2018.

(62) Gonzalez YM, Schiffman E, Gordon SM, Seago B, Truelove EL, Slade G, et al. Development of a brief and effective temporomandibular disorder pain screening questionnaire: reliability and validity. J Am Dent Assoc 2011 Oct;142(10):1183-1191.

II

Tooth wear in Swedish adults—A cross-sectional study

Susanna Gillborg | Sigvard Åkerman | EwaCarin Ekberg

Department of Orofacial Pain and Jaw
Function, Faculty of Odontology, Malmö
University, Sweden

Correspondence

Susanna Gillborg, Department of Orofacial
Pain and Jaw Function, Faculty of
Odontology, Malmö University, Malmö
SE-205 06, Sweden.
Email: susanna.gillborg@mau.se

Funding information

The regional board of Skane and Kalmar,
Sweden, Grant/Award Number: n/a

Abstract

Background: Tooth wear has been recognised as a growing oral health problem in children and adolescents, with erosion often cited as the main cause of the tooth wear. Most studies on tooth wear have been conducted on children and adolescents, and only few studies focus on adults. Our aim was to study the prevalence of different types of tooth wear in an adult population and investigate related factors to tooth wear.

Methods: A total of 831 adults in Sweden participated in the study by completing a questionnaire about oral health, a clinical examination, saliva sample and intraoral photographs. Tooth wear was estimated according to the Basic Erosive Wear Examination index, and the aetiology was determined based on the clinical appearance.

Results: Almost 80% of the individuals had signs of erosion, and over 90% had signs of attrition. A high level of tooth wear was found in 4.6% of the individuals, few of who reported having received information about both attrition and erosion. Significantly, more men had tooth wear. Daily consumption of fruit had a stronger correlation to tooth wear than acidic drinks.

Discussion & Conclusion: A high level of tooth wear was found in 4.6% of the individuals, and it was more common in men than women. Aside from attrition, tooth wear due to erosion was a frequent finding in adults. Only a few of the individuals with a high level of tooth wear reported to have received information about tooth wear from their dentist or dental hygienist.

KEYWORDS

adult, cross-sectional studies, epidemiologic factors, prevalence, tooth attrition, tooth erosion, tooth wear

1 | BACKGROUND

In dentistry, tooth wear is a major problem and a serious risk in maintaining the long-term health of the dentition.¹ Today, we live longer and tooth loss and edentulism are conditions that have become less common.² However, in the younger population, tooth wear that is likely due to dental erosion^{3–6} seems to be increasing. Most studies on tooth wear and erosion focus on children and adolescents.⁷

Tooth wear in the primary dentition has been described as a possible indicator for developing tooth wear in the permanent dentition.^{8,9} The data on tooth wear in adults are scattered, and there is scant research available showing what the current levels of tooth wear are for adults.¹⁰

From a review published in 2009, only 13 studies could present the prevalence of tooth wear in adults: seven of these studies were randomized population-based samples and the remaining six were convenient data. The conclusion was that, by the age of 20, about 3% of patients show signs of severe tooth wear, and by

The peer review history for this article is available at <https://publons.com/publon/10.1111/joor.12887>.

age 70, the number increases to 17%. However, due to different scoring systems, samples and examiners, it is difficult to compare studies.^{10,11} Since then new studies on tooth wear in adults have been published. Still the variations in study design are substantial. A study of 3773 adults, 20 years or older, in United States showed a prevalence of tooth wear of 79.8%. Severe tooth wear was registered in 0% of the younger individuals and in 11% of the older individuals.¹² In 2013, a study was performed in the Netherlands of 1125 individuals aged 25-74. The prevalence of tooth wear was 99%. Severe tooth wear was found in 2% of the younger individuals and in 12% of the older individuals. Their results could also be compared with data from a survey performed in the Netherlands in 2007 and they concluded that the prevalence of tooth wear had increased since then.¹³ A multicenter, multinational cross-sectional study was conducted among 2924 participants between the ages of 18 and 35 years in six Arab countries. The prevalence of tooth wear was reported as relatively high with the highest severity of tooth wear in Oman with a prevalence of severe tooth wear of 60%. They conclude that tooth wear is a significant oral health problem in the Arab population.¹⁴

There is a consensus regarding the definition and diagnostic criteria of various forms of tooth wear.⁷ Tooth *attrition* is the physical wear caused by the action of antagonistic teeth with no foreign substances intervening. This form of tooth wear is characterised by antagonistic glossy plane facets with sharp margins, that only occur on occluding surfaces. Tooth *abrasion* is the physical wear caused by mechanical processes, which also involve foreign substances or objects. The morphological changes can be diffuse or localised depending on the predominant impact. Tooth *erosion* is the chemical wear due to extrinsic or intrinsic acids. Tooth erosion appears first as loss of the physiological surface lustre. In more advanced stages, changes in the original tooth morphology occur. On smooth surfaces, the convex areas flatten and concavities can develop, in which the width clearly exceeds the depth. Lesions are located coronal to the enamel-cementum junction, with an intact border of enamel along the gingival margin. On occlusal surfaces, the cusps become rounded and grooves develop on the cusps and incisal edges. Restorations can rise above the level of the adjacent tooth surfaces.⁷ In 1970, Pindborg had already claimed that tooth wear is the result of attrition, abrasion or erosion.¹⁵ Since then, abfraction has been suggested as a fourth wear-related process. Tooth *abfraction* is described as physical wear resulting from tensile and compressive forces in the cervical region due to the flexing of teeth under occlusal loads, provoking microfractures in the enamel and dentine.⁷ However, the true aetiology of abfraction lesions has been controversial, as other causative factors such as abrasion and erosion have also been considered in the development of these lesions.^{1,16} Many researchers agree that the aetiology is multifactorial and the term, *non-carious cervical lesion* (NCCL), is preferred to describe the loss of tooth substance at the cementum-enamel junction without bacterial involvement. In the clinical setting, it can be difficult to distinguish the various forms of tooth wear from one another. A clinical examination is not adequate to fully understand the aetiology behind the

tooth wear of each individual tooth, but a thorough history should be added.^{17,18}

Although the extent of tooth wear increases with age,¹⁹⁻²² it is not possible to determine age by studying the extent of tooth wear. Historically, the amount of tooth wear caused by the mastication of a coarse diet created a linear curve in time. Today, our food no longer creates that amount of tooth wear. It is suggested that, during one's lifetime, there are periods of more or less tooth wear caused by a variety of factors such as functional activity, parafunctional habits, patterns of mandibular movement, diet, diseases, salivary factors, occupational environment, oral hygiene and various aspects of the modern lifestyle which differ between people and stages of life.²³⁻²⁵ Few studies can present data on progression rates of tooth wear. It is time consuming to evaluate the progression rate of tooth wear for every patient in the clinic. There have also been a lack of clear thresholds for determining whether the tooth wear for each patient can be considered physiological or pathological. A definition of severe tooth wear and pathological tooth wear was published in 2017.²⁶ Severe tooth wear is defined as a substantial loss of tooth structure, with dentin exposure and significant loss ($\geq 1/3$) of the clinical crown. Pathological tooth wear is defined as atypical for the age of the patient, causing pain or discomfort, functional problems or deterioration of aesthetic appearance, which, if it progresses, may give rise to undesirable complications of increasing complexity.

Over 100 different grading systems for the quantification of tooth wear exist.²⁷ A validated, standardised and internationally accepted tooth wear index has been requested. With this intention, a new index—the Basic Erosive Wear Examination (BEWE)—was developed at a workshop in 2007, held on current epidemiological approaches in the field of dental erosion entitled *Current Erosion Indices - Flawed or Valid*.^{28,29} The BEWE is a partial scoring system recording the most severely affected surface in a sextant and quantifies the size of the given lesion in terms of the percentage of the tooth surface affected. The vestibular, occlusal and palatal surfaces of all teeth (except third molars) are graded. The index is not only directed to epidemiological studies but also intended to help clinicians in the registration and evaluation of tooth wear. Attempts to validate the BEWE have been published³⁰; BEWE had a sensitivity of 48.6% and a specificity of 96.1% for moderate tooth wear and 90.9% vs 91.5% for severe tooth wear. Inter-examiner reliability and intra-examiner reliability were found to be moderate. A number of studies on tooth wear using BEWE as an index have been published in recent years.^{14,31}

Evidence suggests that men have more tooth wear than women, but no explanation for the reason behind this difference between genders is given.¹⁰ Bruxism has been discussed to be the main factor behind tooth wear, but this is not supported by the literature.³²⁻³⁵ Acids are potential factors in causing tooth wear. Extrinsic acids like those found in soft drinks, citrus fruit, ketchup, sour candy and juice can cause extensive damage if overused. Foods and drinks with a low pH have increased in modern times. Exposing children to sour-tasting foods at an early age increases their preference for acidic food and drinks later in life.³⁶ In addition, individuals exposed to

FIGURE 1 Flowchart of the individuals who participated in this study

intrinsic acids from vomiting and reflux have shown to have more tooth wear than controls.³⁷

The large increase in the numbers of children and adolescents with tooth wear due to erosion is known, but it is important to increase knowledge about tooth wear in the adult population. Accordingly, if tooth wear is a condition that follows into adulthood, this would result in a greater need for future dental care. We need more knowledge about whether erosion is the most common cause of tooth wear in adults and related factors in the different forms of tooth wear. One hypothesis was that tooth wear would be registered in almost all adults. Another hypothesis was that high levels of tooth wear would be registered in only a small percentage of the individuals. Furthermore, we expected attrition to be the most common reason for tooth wear in adults, and that signs of erosion would be seen in more than 50% of the individuals examined, as reported in a study from 2015, using BEWE to evaluate tooth wear among 500 individuals in Israel.³¹ A relation between tooth wear and soft drinks has been reported in many studies, but we expected this to be weak among Swedish adult individuals. Rather, we expected a

stronger relation between tooth wear and frequent intake of fruit. We also expected to find more tooth wear among males, as most of the studies that have investigated gender differences in tooth wear have shown this relation.^{10,11}

We aim to study the prevalence of different types of tooth wear in an adult population and investigate related factors to the different types of tooth wear. Another aim is that the results will help guide the clinician in diagnosing tooth wear and to better plan for the preventive care and the rehabilitation of the dentition.

2 | METHODS

In 2007–08, a study concerning general oral health was performed in the county of Skåne in southern Sweden. A questionnaire was sent to a randomly selected sample of 1000 individuals. The individuals were 20–89 years old and registered as living in the county during 2007. The sample was obtained from SPAR (Swedish Population Address Register). After excluding individuals who had moved from

TABLE 1 Description of 13 variables selected from questionnaire and history that we expected to have a relation to tooth wear

Age	0 = 20-29 y 1 = 30-39 y 2 = 40-49 y 3 = 50-59 y 4 = 60-69 y 5 = 70-79 y 6 = 80-89 y
Gender	0 = man 1 = woman
Information about tooth grinding	Has your dentist or hygienist told you that you have tooth wear due to grinding? 0 = no 1 = yes
Information about tooth erosion	Has your dentist or hygienist told you that you have tooth wear due to tooth erosion from drinks (Coca-Cola, juice, sport drinks), food (citrus fruit) or reflux? 0 = no 1 = yes
Information about tooth abrasion	Has your dentist or hygienist told you that you have tooth wear due to tooth brushing? 0 = no 1 = yes
Fruit	How often do you eat apples or citrus fruit? 0 = almost never 1 = once a month 2 = once a week 3 = daily 4 = more than two times a day 5 = more than five times a day
Drinks	How often do you drink Coca-Cola, other kind of soft drinks, energy drinks or juice? 0 = almost never 1 = once a month 2 = once a week 3 = daily 4 = more than two times a day 5 = more than five times a day
Smoking	Do you smoke? 0 = no 1 = yes
Snuff	Do you use snuff? 0 = no 1 = yes
Alcohol	How often do you drink wine, beer or spirit? 0 = never 1 = once a month 2 = twice a month 3 = once per week 4 = twice per week 5 = daily

(Continues)

TABLE 1 (Continued)

Occupation	0 = working/studying 1 = housewife/senior citizen/on parental leave 2 = unemployed/sick listed/early retired
TMD pain	"Yes" to one or both of the following two questions: Do you have pain in your face, jaw, temple, in front of the ear or in your ear once a week or more often? Do you have pain when you open your mouth or when you chew once a week or more often?
Medications	What medications do you take?

the region, had an unknown address or were deceased, 966 individuals remained in the final sample (Figure 1).

The questionnaire contained 58 questions, of which, all had multiple choice answers and several questions had been validated and used previously in other questionnaires.³⁸ In this study, 13 questions relating to tooth wear were selected among the 58 questions for analysis (Table 1).

The individuals in the study were offered a clinical examination. A total of 451 individuals participated in a clinical examination, and of these individuals, all but one completed the questionnaire. The examination included a saliva sample, intraoral photographs and a history which included medication and dietary habits. The clinical examination was carried out at the Faculty of Odontology at Malmö University in Sweden and at three clinics of the Public Dental Health Service in Skåne during the period of March 2007 and November 2008. Of the clinical examinations, 90.5% were performed by four out of eight dentists, all employed at the Department of Oral Diagnostics, Faculty of Odontology, Malmö University in Sweden. All patients were examined using standard examination procedures. The dentists were co-ordinated regarding the diagnostic criteria through comprehensive written instructions, practice and by discussing clinical cases.³⁸

A similar investigation was carried out in another county in southern Sweden—Kalmar. This sample was also randomized from SPAR and initially consisted of 2000 individuals between the ages of 20 and 90 (Figure 1). The questionnaire consisted of 56 questions, of which, most were the same or very similar questions as those of the study in Skåne. Of the 2000 individuals, 900 were offered a dental examination and 380 individuals were examined. Two dentists performed the clinical examinations. The examinations took place at four different public dental clinics in the county of Kalmar during the period of April 2011 and July 2012 and consisted of the same variables as the examination performed in Skåne.

The data from the investigations in Skåne and Kalmar were compiled in this study (Figure 1) and evaluated by one dentist (SG) concerning tooth wear. A calibration was performed between the dentist (SG) and a specialist in orofacial pain and jaw function (SÅ). The inter agreement after examining clinical photographs from 10 individuals was moderate.³⁹ The clinical photographs from 831 individuals were compared with the records from the clinical examination concerning tooth wear. If tooth wear was found in the

photographs that had not been noted at the clinical examination, this was added to the record and the level of tooth wear was then estimated by using BEWE for all individuals. The scores range from 0 to 3:0 represents "No erosive tooth wear"; 1 represents "Initial loss of surface texture"; 2 represents "Distinct defect, hard tissue loss < 50% of the surface area"; and 3 represents "Hard tissue loss ≥ 50% of the surface area." Once all the sextants have been assessed, the sum of the scores is calculated and converted to a total risk level of tooth wear. If the sum is between 0 and 2, it is considered as having no risk. A sum between 3 and 8 indicates a low risk, 9 and 13 a medium risk and 14 and 18 is a high risk. Individuals who did not fulfil the criteria for BEWE, by not having teeth in all 6 sextants with enough tooth substance to evaluate the level of tooth wear, were calculated by using their score and multiplying it by the factor resulting in a comparable score. Individuals with assessable tooth substance in two sextants were evaluated, and the score was then multiplied by 3. For 3 sextants, we multiplied the number with 2, for 4 sextants with 1.5 and for 5 sextants with 1.2. The group with teeth in all 6 sextants comprised the BEWE group and the group that were calculated with the adjusted version of BEWE comprised the BEWE-adjusted group.

When evaluating different types of tooth wear, signs of attrition were registered if glossy plane facets with sharp margins were found on occluding surfaces in the photographs or if they were registered in the journal. In the majority of individuals, these were registered on incisal surfaces and on the cusps of the canines, and in some individuals, on teeth more posteriorly. Several signs of erosion were recorded and used to determine the aetiology: reversed V-sign incisal on maxillary central incisors, thin enamel on the palatal surface of the maxillary incisors, loss of the distinct shape on the tooth surface, thinning out of the enamel without signs of attrition, loss of surface lustre, raised restorations, grooves on the cusps or incisal edges and/or restorations rising above the level of the adjacent morphology. Signs of abrasion and abfraction were not recorded but seen as possible parts of the aetiology if the individual had NCCLs. Other damage due to abrasion was not possible to distinguish from erosion.

These studies were approved by the Ethics Board at the University of Lund in Lund, Sweden. All participants signed written informed consent prior to enrolment and were free to omit any item in the questionnaire or withdraw from the study at any time without providing an explanation.

2.1 | Statistical methods

The individuals were stratified in four groups on the basis of risk levels of tooth wear (no risk, low risk, medium risk and high risk) and in seven age groups (20-29, 30-39, 40-49, 50-59, 60-69, 70-79, 80-89 years). The χ^2 -test compared the distribution of categorical variables. Possible related factors related to levels of tooth wear were analysed using logistic regression models with a likelihood-ratio test. The related factors were as follows: age, gender, information about tooth wear, information about tooth erosion, information about tooth abrasion, fruit consumption, consumption of acidic drinks, smoking habits, use of snuff, alcohol consumption, occupation, TMD pain and medications. Statistical calculations were performed using the Statistical Package for the Social Sciences (SPSS 22 for Windows) with a significance level of 5% for all tests.

3 | RESULTS

Clinical photographs and records from 831 individuals were evaluated (Figure 1). A total of 81% of the individuals had teeth in all 6 sextants that were possible to evaluate regarding the level of tooth wear (BEWE group). Of the remaining individuals, 54 had prostheses, full arch bridges, fillings/crowns covering most part of the teeth, or were edentulous, leaving 106 individuals (13% of the original group) with teeth that could be evaluated for tooth wear in 2-5 sextants (BEWE-adjusted group). Table 2 shows the number of sextants with teeth with enough visible enamel/dentine to be evaluated for risk levels of tooth wear and the number of individuals counted for each number of sextant. Of the 671 individuals in the BEWE group, a score between 0 and 2 (no risk) was registered in 3.3% of the individuals, a score between 3 and 8 (low risk) was registered in 49.6% and a score between 9 and 13 (medium risk) in 42.5%. The BEWE score of 14 or over (high risk) was registered in 4.6% of the individuals in the BEWE group. Of the 106 individuals in the BEWE-adjusted group, all showed signs of tooth wear: a low risk level of tooth wear was registered in 25.5%, medium risk in 52.8%, and high risk in 21.7% (Figure 2).

TABLE 2 Number of sextants with teeth with visible enamel/dentine to be evaluated for risk levels of tooth wear and the number of individuals counted for each number of sextants

Sextants	Individuals
0	47
1	7
2	16
3	24
4	23
5	43
6	671
Total	831

When analysing age in the BEWE group and the BEWE-adjusted group, significant differences were registered for the different age groups (Tables 3A,B).

In the BEWE-adjusted group, only four individuals are younger than 50 years, whereas in the BEWE group, all ages are represented equal up to the age of 70.

In addition, significant differences were found between genders in the BEWE group (Table 4). In the group with no risk level of tooth wear, 68% were women and 32% were men, whereas in the group of high risk level of tooth wear, 39% were women and 61% were men. When performing a regression analysis between genders on the related factors identified as having a significant association with tooth wear in our study, no differences were found.

In the BEWE group, almost 80% of the individuals had signs of erosion and over 90% had signs of attrition. A shiny facet on either an incisor or a canine was the most common finding of tooth wear due to attrition. NCCLs were registered on at least one tooth in 28% of the individuals. All individuals with a high risk level of tooth wear had signs of both erosion and attrition (Table 4). In the BEWE-adjusted group, 63% showed signs of erosion and 98% showed signs of attrition. For NCCL, the figure was almost the same as for the BEWE group, with signs in at least one tooth in 29% of the individuals. Of the individuals with high risk level of tooth wear, all had signs of attrition, as in the BEWE group, but only 70% showed signs of erosion.

Retraction of the gingival margin was registered in 78% of the individuals. NCCL was registered in 29% of the individuals and became more frequent with higher levels of tooth wear (Table 4).

There was a statistically significant relation between the level of tooth wear and the frequency of fractured teeth or fillings. Three times as many individuals with fractures in the high risk group were found in comparison with the no risk group (Table 4).

In total, 52% of the individuals in the high risk group reported that they had been informed about tooth wear due to tooth grinding from their dentist or hygienist, and 10% reported being informed about erosion (Table 4). Of the individuals with NCCL, 40% reported being informed about tooth wear due to tooth brushing.

Daily consumption of acidic fruit was significantly associated with a higher grade of tooth wear (Table 4). Twice as many individuals reported drinking acidic drinks every day or more often in the high risk group compared with the no risk group, but the relation was not statistically significant. The difference in consumption of acidic drinks disappears if the risk groups, "no risk" and "low risk," are made into one group and "medium risk" and "high risk" are made into another group.

Neither the dentist clinical registration of mouth dryness nor the registration of low saliva sample of stimulated saliva showed a statistical relation to tooth wear. In addition, smoking, use of snuff, high alcohol consumption, occupation or signs of TMD pain showed no statistical relation to tooth wear. Taking prescribed medications and specifically antidepressants or sleeping medications were not related to risk level of tooth wear.

FIGURE 2 Frequency of the different risk levels of tooth wear in the BEWE group (n = 671) and the BEWE-adjusted group (n = 106)

4 | DISCUSSION

As expected, tooth wear was found in all individuals. Not all tooth wear was registered at the clinical examination, but when evaluating the clinical photographs, defects on minor surfaces or more diffuse defects could be detected and it amounted to all the individuals having at least 1 sextant with signs of tooth wear. A high risk level of tooth wear was found in all age groups, and the risk increased with age. The youngest individual with a high risk level of tooth wear was 26 years old and had extensive tooth wear on all molars due to erosion. According to these data, a dentist treating 1000 patients a year will meet 25 individuals aged 20-29 years with a high risk level of tooth wear who need a thorough investigation concerning the aetiology and preventive care and/or restorations.

Most of the individuals had visible signs of attrition. This result does not automatically translate to prevalence of bruxism since an individual could have periods with bruxism and periods without bruxism during a lifetime.⁴⁰

In 1988, Hugoson et al reported that tooth wear in Swedish adults mostly affected canines and incisors. This finding showed that the distribution of wear in adults at that time differed in distribution from the tooth wear seen in prehistorical skulls, where molars were affected to a higher extent. Historically, tooth abrasion was caused by the mastication of food and affected the whole dentition. Today, our food is not as abrasive as it was in the past and the most common cause of abrasion is tooth cleaning.⁴¹ NCCL is often suggested to be tooth wear caused by excessive tooth brushing.¹⁶ Studies have shown that the duration, force and frequency of tooth brushing are associated with gingival recession and NCCL, but the available evidence does not confirm these to be the most important aetiological factors.⁴² Today's toothpaste gives 1 mm of tooth wear if used daily for a hundred years and cannot explain the aetiology to NCCL alone.⁴³ Larger impacts on the tooth surface have been reported when exposed to acid combined with abrasion such as tooth brushing. In historical or aboriginal populations in which the diet was abrasive but not erosive, NCCL is rare or absent.⁴⁴

Gingival retraction and NCCL were common findings among the individuals in this study. There are other suggested explanations to the aetiology behind NCCL than tooth cleaning. Tooth abfraction is

described as loss of tooth substance in the cervical region due to stress caused by mastication, malocclusion or bruxism. The loading of the tooth results in cyclic tension and compression concentrated in the cervical area and eventually causes cracking of the tooth enamel and dentine. The theory is mainly supported by laboratory studies and currently lacks support from clinical studies.¹⁶ NCCL is likely the result of erosion, abrasion and attrition working together. More research is needed to fully understand the aetiology to NCCL.¹⁶ In our results, we have not specified the prevalence of abrasion. Without a complete history from each individual about diet, working environment, et cetera, both today and earlier in life, it is difficult to separate abrasion from damage caused by erosion.

When identifying tooth erosion, cuppings were a frequent finding. Cuppings are strongly correlated with tooth erosion, and it has been stated that they can be regarded as an indicator of the onset of erosion.^{45,46} A total of 48% of the individuals had cuppings, and the surface with most cuppings was the cusps on the first molars in the lower jaw. In the study of tooth wear in Swedish adults in 1988, tooth wear in the molar region was uncommon indicating that the tooth wear in Swedish adults has changed its distribution in the dentition since 1988. That cuppings is not found on all tooth surfaces equally could be explained by the differences in salivary protection. The further the site is from the secretion of the major salivary glands, the longer time it takes for the pH to be stabilized again after exposure to acid.⁴⁷ The mucin and other macro molecules in the saliva create a lining, called the *pellicle*, over the tooth surface and protect it from acids and decreases the friction. The thickness of the pellicle differs between different sites in the mouth.

There is also variation concerning the buffering capacity in the saliva between different individuals. Demineralized enamel/dentine can regain strength by absorbing new minerals from the saliva. We studied both the amount of saliva in relation to tooth wear and the acid neutralisation ability in the saliva. We also chose to study whether having many prescribed medications was related to tooth wear, as many medications often result in mouth dryness. However, none of these three factors showed any relation to tooth wear in this population.

As more than 100 indices are available to rate the level of tooth wear in adults, it was not obvious which one to choose in this study.

Age group		20-29	30-39	40-49	50-59	60-69	70-79	80-89	Total
(A)									
No risk									
n	15	5	1	1	0	0	0	0	22
%	68.2	22.7	4.5	4.5	0.0	0.0	0.0	0.0	100.0
Low risk									
n	74	81	73	48	42	10	5	333	
%	22.2	24.3	21.9	14.4	12.6	3.0	1.5	100.0	
Medium risk									
n	27	47	53	73	61	21	3	285	
%	9.5	16.5	18.6	25.6	21.4	7.4	1.1	100.0	
High risk									
n	3	3	5	9	7	4	0	31	
%	9.7	9.7	16.1	29.0	22.6	12.9	0.0	100.0	
Total									
n	119	136	132	131	110	35	8	671	
%	17.7	20.3	19.7	19.5	16.4	5.2	1.2	100.0	
(B)									
No risk									
n	-	-	-	-	-	-	-	-	-
%	0	0	0	0	0	0	0	0	0
Low risk									
n	-	-	2	1	12	10	2	27	
%	0	0	7.4	3.7	44.4	37.0	7.4	100.0	
Medium risk									
n	1	0	1	7	23	20	4	56	
%	1.8	0.0	1.8	12.5	41.1	35.7	7.1	100.0	
High risk									
n	-	-	-	1	7	8	7	23	
%	0	0	0	4.3	30.4	34.8	30.4	100.0	
Total									
n	1	-	3	9	42	38	13	106	
%	0.9	0	2.8	8.5	39.5	35.8	12.3	100.0	

TABLE 3 Levels of tooth wear in the different age groups in the (A) BEWE group (n = 671); (B) BEWE-adjusted group (n = 106)

BEWE was developed after a consensus procedure between several researchers in the field of dental erosion, from several countries.²⁸ It was decided that this study would use BEWE because it was designed to provide a simple scoring system that can be used with the diagnostic criteria of all existing indices that aim to transfer their results into one unit (the BEWE score sum). The validation studies performed using BEWE have been carried out on the younger population.³⁰ Our results show that the index successfully captures individuals up to the age of 70. After this age, the number of individuals markedly drops due to not fulfilling the criteria of having teeth in all six sextants. According to BEWE, we should have to exclude 20% of the individuals due to tooth loss, but we chose to include these individuals in the study by adjusting the BEWE. Although, we separated the results from the group fulfilling the criteria for BEWE. The results

from the adjusted BEWE group have to be interpreted with that in mind that the tooth wear is perhaps not comparable to the tooth wear registered in the individuals with teeth in all six sextants. The tooth wear in the sextants missing in the data could have been both less and more than the tooth wear in the remaining sextants, and having fewer teeth could be the cause of a higher level of tooth wear.

It is known that gender plays a role in the aetiology of tooth wear.¹⁰ In most of the studies that report a difference, men have significantly more tooth wear than women, which was in line with our results. No information is available that explains why men have more tooth wear than women. A more detailed history about recent and present risk factors needs to be obtained.

In 2013, there was a collaboration between seven European countries to assess the prevalence of tooth wear in 3187 patients

TABLE 4 Distribution of factors that could be related to tooth wear divided between the risk levels of tooth wear for the BEWE group (n = 671)

	No risk (n = 22) %	Low risk (n = 333) %	Medium risk (n = 285) %	High risk (n = 31) %	P-value	Total n = 671%
Women	68.2	57.4	45.3	38.7	.004**	51.7
Signs of erosion	22.7	74.7	83.2	100.0	.000***	77.8
Signs of attrition	86.4	90.1	97.2	100.0	.001**	90.1
NCCL (Non-Carious Cervical Lesions)	0.0	23.4	35.8	41.9	.000***	28.8
Gingival retraction	54.5	72.7	84.6	83.9	.000***	77.6
Fractured teeth	4.5	7.2	15.4	6.5	.006**	10.6
Fractured fillings	9.1	13.8	23.5	29.0	.004**	18.5
Information about tooth grinding	0.0	9.9	14.5	51.6	.000***	13.4
Information about tooth erosion	0.0	4.2	7.1	9.7	.194	5.5
Information about wear due to toothbrushing	18.2	21.0	24.7	12.9	.379	22.1
Daily intake of fruit	27.3	37.8	47.5	54.8	.019*	42.3
Daily intake of acidic drinks	9.1	26.4	25.4	16.1	.199	24.9

P = * ≤ 0.05 , ** ≤ 0.01 , *** ≤ 0.001

aged 18–35 years and to identify related risk factors.⁴⁸ Our results concerning the relation between soft drinks and fruit and tooth wear is in line with their results, and confirm the hypothesis—a stronger association between fruit and high levels of tooth wear than soft drinks and high levels of tooth wear. This result, as mentioned earlier, does not take in consider habits that may have been before. Soft drinks that are reported in many studies as a risk factor for tooth wear are perhaps something that is excluded from the diet as the tooth wear starts being a problem, functionally or aesthetically.

One strength of our study is the number of individuals evaluated both clinically and with intraoral photographs. Another strength is having a clinical evaluation of the degree of tooth wear chairside, while another dentist evaluated the degree of tooth wear in the clinical photographs guided by the clinical evaluation.

The weakness in this study lies in how we asked about dietary habits and other possible related factors to tooth wear. This leaves open the possibility that the individuals have changed their habits during their lifetime, and thus, the true factor(s) behind their tooth wear will not be identified. Tooth wear can be caused by a variety of factors, and it would be difficult to create a questionnaire with the ability to screen for all the possibilities. The questionnaire was intended to evaluate the probable aetiology behind the tooth wear, but there were individuals with obvious clinical signs of dental erosion that did not report any of the known risk factors in the questionnaire. Therefore, the aetiology behind the tooth wear in our results is based solely on the clinical signs. For future studies, it would be a significant advantage if related factors could be identified by better methods.

The results of this study could serve as a basis for future studies to be able to compare how the tooth wear of adults changes over

time. Furthermore, the results may help dentists and hygienists in general dental practice to change their approach regarding tooth wear, and be more observant to changes in tooth wear caused by erosion. Many individuals with extensive tooth wear were not informed or did not remember being informed about tooth wear. More often, they reported receiving information about bruxism than erosion, and many individuals reported being informed about tooth wear due to tooth brushing. This result was not expected, as the literature often mentions the growing problem of dental erosion.⁴⁶ The softening effect erosion has on dental hard tissue enhances the effect of tooth attrition and tooth abrasion, speeding up the progression rate.

There are few studies that attempt to measure the progression rate of tooth wear. Data of 1239 permanent maxillary and mandibular central incisors from 346 persons covering the ages between 10 and 73 years showed a decrease by an average of 1.01 mm of the height of the maxillary incisors over the 60-year span. For mandibular incisors, the decrease was an average of 1.46 mm.⁴⁹ Another study followed the progression of wear in enamel due to tooth attrition during 4 years. Areas with occlusal contact in the premolar region had an enamel wear of 88 μm and in the molar region 153 μm .⁵⁰ The existing methods, to measure tooth wear in the everyday clinic, are not sensitive enough to measure small changes in progression. The consequence is that the clinician registers tooth wear first when the eye can detect tooth wear atypical for the age of the patient. Unless the patient complains from discomfort or other undesirable complications, alerting the clinician to be more observant, unnecessary damage to the dentition is likely to occur. The definition of severe tooth wear and pathological tooth wear, published in 2017, clarifies that severe tooth

wear does not have to be considered pathological and vice versa, pathological tooth wear does not have to be severe.²⁶ The absence of normative values for tooth wear is a question for future studies. In addition, a further intention is to develop a more sensitive instrument that not only calculates for present tooth wear but also the speed of the progression of tooth wear over time.

5 | CONCLUSION

Tooth wear was registered in almost all adults, with high levels of tooth wear being found in 4.6% of the individuals. Attrition was the most common reason for the tooth wear, but signs of erosion were seen in almost 80% of the individuals. The erosion was related more to the frequent intake of fruit than to soft drinks. Men had more tooth wear than women, but no explanation could be found among the related factors studied here. We found a moderate level of inter-examiner agreement with BEWE and it was possible to evaluate tooth wear in individuals between 20 and 69 years.

ACKNOWLEDGMENTS

This study was performed in collaboration with the Regional Board of Dental Public Health in Skåne and Kalmar, Sweden.

ORCID

Susanna Gillborg <https://orcid.org/0000-0002-8387-8710>

REFERENCES

- Lee A, He LH, Lyons K, Swain MV. Tooth wear and wear investigations in dentistry. *J Oral Rehabil*. 2012;39(3):217-225.
- Lussi A, Carvalho TS. Erosive tooth wear: a multifactorial condition of growing concern and increasing knowledge. *Monogr Oral Sci*. 2014;25:1-15.
- Bartlett DW. The role of erosion in tooth wear: aetiology, prevention and management. *Int Dent J*. 2005;55(4 Suppl 1):277-284.
- Jaeggi T, Lussi A. Prevalence, incidence and distribution of erosion. *Monogr Oral Sci*. 2006;20:44-65.
- Carlsson GE, Johansson A. *Dental Erosion: Prevalens av Dental Erosion*. Stockholm, Sweden: Förlagshuset Gothia; 2006.
- Johansson AK On dental erosion and associated factors. *Swed Dent J Suppl*. 2002;(156):1-77.
- Ganss C, Young A, Lussi A. Tooth wear and erosion: methodological issues in epidemiological and public health research and the future research agenda. *Community Dent Health*. 2011;28(3):191-195.
- Harding MA, Whelton HP, Shirodaria SC, O'Mullane DM, Cronin MS. Is tooth wear in the primary dentition predictive of tooth wear in the permanent dentition? Report from a longitudinal study. *Community Dent Health*. 2010;27(1):41-45.
- Ganss C, Klimek J, Giese K. Dental erosion in children and adolescents—a cross-sectional and longitudinal investigation using study models. *Community Dent Oral Epidemiol*. 2001;29(4):264-271.
- Van't Spijker A, Rodriguez JM, Kreulen CM, Bronkhorst EM, Bartlett DW, Creugers NH. Prevalence of tooth wear in adults. *Int J Prosthodont*. 2009;22(1):35-42.
- Jaeggi T, Lussi A. Prevalence, incidence and distribution of erosion. *Monogr Oral Sci*. 2014;25:55-73.
- Okunseri C, Wong MC, Yau DT, McGrath C, Szabo A. The relationship between consumption of beverages and tooth wear among adults in the United States. *J Public Health Dent*. 2015;75(4):274-281.
- Wetselaar P, Vermaire JH, Visscher CM, Lobbezoo F, Schuller AA. The prevalence of tooth wear in the Dutch adult population. *Caries Res*. 2016;50(6):543-550.
- Awad MA, El Kassas D, Al Harthi L, et al. Prevalence, severity and explanatory factors of tooth wear in Arab populations. *J Dent*. 2019;80:69-74.
- Pindborg J, ed. *Pathology of the Dental Hard Tissues*. Copenhagen, Denmark: Munksgaard; 1970.
- Bartlett DW, Shah P. A critical review of non-carious cervical (wear) lesions and the role of abfraction, erosion, and abrasion. *J Dent Res*. 2006;85(4):306-312.
- Johansson A, Johansson AK, Omar R, Carlsson GE. Rehabilitation of the worn dentition. *J Oral Rehabil*. 2008;35(7):548-566.
- Fares J, Shirodaria S, Chiu K, Ahmad N, Sherriff M, Bartlett D. A new index of tooth wear. Reproducibility and application to a sample of 18- to 30-year-old university students. *Caries Res*. 2009;43(2):119-125.
- Bernhardt O, Gesch D, Splieth C, et al. Risk factors for high occlusal wear scores in a population-based sample: results of the Study of Health in Pomerania (SHIP). *Int J Prosthodont*. 2004;17(3):333-339.
- Hugoson A, Bergendal T, Ekfeldt A, Helkimo M. Prevalence and severity of incisal and occlusal tooth wear in an adult Swedish population. *Acta Odontol Scand*. 1988;46(5):255-265.
- Johansson A, Haraldson T, Omar R, Kiliaridis S, Carlsson GE. An investigation of some factors associated with occlusal tooth wear in a selected high-wear sample. *Scand J Dent Res*. 1993;101(6):407-415.
- Nunn J, Morris J, Pine C, Pitts NB, Bradnock G, Steele J. The condition of teeth in the UK in 1998 and implications for the future. *Br Dent J*. 2000;189(12):639-644.
- Johansson A. A cross-cultural study of occlusal tooth wear. *Swed Dent J Suppl*. 1992;86:1-59.
- Barbosa Tde S, Miyakoda LS, Poczaruk Rde L, Rocha CP, Gavião MB. Temporomandibular disorders and bruxism in childhood and adolescence: review of the literature. *Int J Pediatr Otorhinolaryngol*. 2008;72(3):299-314.
- Johansson AK, Johansson A, Birkhed D, et al. Dental erosion associated with soft-drink consumption in young Saudi men. *Acta Odontol Scand*. 1997;55(6):390-397.
- Loomans B, Opdam N, Attin T, et al. Severe tooth wear: European consensus statement on management guidelines. *J Adhes Dent*. 2017;19(2):111-119.
- Wetselaar P. A PhD completed 6. Tooth Wear Evaluation System: development and applications. *Ned Tijdschr Tandheelkd*. 2016;123(10):492-494.
- Bartlett D, Ganss C, Lussi A. Basic Erosive Wear Examination (BEWE): a new scoring system for scientific and clinical needs. *Clin Oral Invest*. 2008;12(Suppl 1):S65-S68.
- Young A, Amaechi BT, Dugmore C, et al. Current erosion indices—flawed or valid? Summary. *Clin Oral Invest*. 2008;12(Suppl 1):S59-63.
- Olley RC, Wilson R, Bartlett D, Moazzez R. Validation of the basic erosive wear examination. *Caries Res*. 2014;48(1):51-56.
- Vered Y, Lussi A, Zini A, Gleitman J, Sgan-Cohen HD. Dental erosive wear assessment among adolescents and adults utilizing the basic erosive wear examination (BEWE) scoring system. *Clin Oral Invest*. 2014;18(8):1985-1990.
- Khan F, Young WG, Daley TJ. Dental erosion and bruxism. A tooth wear analysis from South East Queensland. *Aust Dent J*. 1998;43(2):117-127.

33. Lavigne GJ, Rompre PH, Montplaisir JY. Sleep bruxism: validity of clinical research diagnostic criteria in a controlled polysomnographic study. *J Dent Res*. 1996;75(1):546-552.
34. Lavigne GJ, Khoury S, Abe S, Yamaguchi T, Raphael K. Bruxism physiology and pathology: an overview for clinicians. *J Oral Rehabil*. 2008;35(7):476-494.
35. Ekfeldt A, Hugoson A, Bergendal T, Helkimo M. An individual tooth wear index and an analysis of factors correlated to incisal and occlusal wear in an adult Swedish population. *Acta Odontol Scand*. 1990;48(5):343-349.
36. Gambon DL, Brand HS, Veerman EC. Dental erosion in the 21st century: what is happening to nutritional habits and lifestyle in our society? *Br Dent J*. 2012;213(2):55-57.
37. Schlueter N, Tveit AB. Prevalence of erosive tooth wear in risk groups. *Monogr Oral Sci*. 2014;25:74-98.
38. Lundegren N. Oral health and self-perceived oral treatment need of adults in Sweden. *Swed Dent J Suppl*. 2012;(223):10-76.
39. Viera AJ, Garrett JM. Understanding interobserver agreement: the kappa statistic. *Fam Med*. 2005;37(5):360-363.
40. Manfredini D, Winocur E, Guarda-Nardini L, Paesani D, Lobbezoo F. Epidemiology of bruxism in adults: a systematic review of the literature. *J Orofac Pain*. 2013;27(2):99-110.
41. Shellis RP, Addy M. The interactions between attrition, abrasion and erosion in tooth wear. *Monogr Oral Sci*. 2014;25:32-45.
42. Heasman PA, Holliday R, Bryant A, Preshaw PM. Evidence for the occurrence of gingival recession and non-carious cervical lesions as a consequence of traumatic toothbrushing. *J Clin Periodontol*. 2015;42(Suppl 16):S237-S255.
43. Addy M, Hunter ML. Can tooth brushing damage your health? Effects on oral and dental tissues. *Int Dent J*. 2003;53(Suppl 3):177-186.
44. Nguyen C, Ranjitkar S, Kaidonis JA, Townsend GC. A qualitative assessment of non-carious cervical lesions in extracted human teeth. *Aust Dent J*. 2008;53(1):46-51.
45. Khan F, Young WG, Law V, Priest J, Daley TJ. Cupped lesions of early onset dental erosion in young southeast Queensland adults. *Aust Dent J*. 2001;46(2):100-107.
46. Johansson AK, Omar R, Carlsson GE, Johansson A. Dental erosion and its growing importance in clinical practice: from past to present. *Int J Dent*. 2012;2012:632907.
47. Macpherson LM, Dawes C. Distribution of sucrose around the mouth and its clearance after a sucrose mouthrinse or consumption of three different foods. *Caries Res*. 1994;28(3):150-155.
48. Bartlett DW, Lussi A, West NX, Bouchard P, Sanz M, Bourgeois D. Prevalence of tooth wear on buccal and lingual surfaces and possible risk factors in young European adults. *J Dent*. 2013;41(11):1007-1013.
49. Ray DS, Wiemann AH, Patel PB, Ding X, Kryscio RJ, Miller CS. Estimation of the rate of tooth wear in permanent incisors: a cross-sectional digital radiographic study. *J Oral Rehabil*. 2015;42(6):460-466.
50. Lambrechts P, Braem M, Vuylsteke-Wauters M, Vanherle G. Quantitative in vivo wear of human enamel. *J Dent Res*. 1989;68(12):1752-1754.

How to cite this article: Gillborg S, Åkerman S, Ekberg E.

Tooth wear in Swedish adults—A cross-sectional study. *J Oral Rehabil*. 2019;00:1-11. <https://doi.org/10.1111/joor.12887>

isbn 978-91-7877-048-9 (print)

isbn 978-91-7877-049-6 (pdf)

MALMÖ UNIVERSITY
205 06 MALMÖ, SWEDEN
WWW.MAU.SE